and sectioned. Most other observers, however, have found that the sebaceous glands, far from being hypertrophied, are actually atrophied.

2.—Cysts of New Formation

Many recent observers (literature by Benecke, 1931) have reverted to the view originally held by Paget that the majority of sebaceous cysts are cysts of new formation and that a true retention cyst of the hair follicle or sebaceous duct is rare. Benecke believed that the majority of the so-called sebaceous cysts met with clinically are more properly termed "epidermoid cysts." These arise by the accumulation of secretion from isolated rests of epidermal cells lying within the corium. Other types of cysts of new formation have been described, arising as naevoid new growths or traumatic implantation cysts.

As evidence against the retention theory attention is drawn to: (i) the rarity of cyst formation in cases of severe acne; (ii) the rarity of comedones in cases of sebocystomatosis; (iii) the difficulty of demonstrating by serial section an occluded or partially obliterated sebaceous duct; (iv) the occurrence of cysts of identical histological structure on the palms of the hands and the volar surface of the fingers, sites devoid of sebaceous glands and hair follicles.

Summary

A family presenting multiple hereditary sebaceous cysts is described. This type of case throws light upon the aetiology of the common sebaceous cyst. The evidence against the widely accepted theory of retention is discussed.

BIBLIOGRAPHY

Familial Multiple Sebaceous Cysts:

- Klausner, E. (1917). *Derm. Wschr.*, **65**, 711.
Paget, J. (1870). *Lectures on Surgical Pathology*, third edition, p. 441.
— (1870a). *Ibid.*, p. 444.
Prakken, J. R. (1933). *Derm. Z.*, **66**, 215.
Weber, F. Parkes, and Schlüter, A. (1936). *Proc. roy. Soc. Med.*, Sect. Derm., **30**, 29.

Multiple Sebaceous Cysts (Sebocystomatosis):

- Arzt, L. (1926). *Zbl. Haut- u. Geschlechtskr.*, **20**, 275.
— (1930). *Ibid.*, **33**, 311.
Benecke, E. (1931). *Frankfurt. Z. Path.*, **42**, 502.
Bossellini, P. L. (1898). *Arch. Derm. Syph.*, Chicago, **45**, 81.
Case report at meeting in Rostock (1929). *Zbl. Haut- u. Geschlechtskr.*, **32**, 415.
Chiari, H. (1890). *Z. Heilk.*, **12**, 189.
Choyce, C. C. (1932). *A System of Surgery*, third edition, **1**, 576.
Cockayne, E. A. (1933). *Inherited Abnormalities of the Skin and its Appendages*, p. 352, London.
Crocker, H. Radcliffe (1888). *Diseases of the Skin*, p. 572.
Gans, O. (1928). *Histologie der Hautkrankheiten*, **2**, 219.
Günther, H. (1917). *Derm. Wschr.*, **64**, 481.
Jamieson, W. A. (1873). *Edinb. med. J.*, **19**, 223.
Maclaren, P. H. (1886). *Trans. Med.-Chir. Soc., Edinb.*, **6**, 77.
Orr, H. (1923). *Proc. roy. Soc. Med.*, Sect. Derm., **17**, 11.
Pringle, J. J. (1899). *Brit. J. Derm. Syph.*, **11**, 381.
Rayer, P. F. O. (1835). *Diseases of the Skin*, p. 1038.

H. R. Hathaway, N. H. Casels, and R. M. Waters (*Anesth. and Analges.*, March-April, 1937, p. 89) review 491 administrations of oxygen in 408 patients. The oro-pharyngeal insufflation technique was used in 92 per cent. of cases. This technique was conveniently and effectively applied with few objections and little discomfort. The authors were able to confirm statistically that early oxygen treatment of anoxic patients improves results. The fact that oxygen was not restricted to dangerously ill patients brought about a relatively low mortality rate.

OSTEOCHONDRITIS DISSECANS

BY

DOUGLAS S. STEVENSON, M.B.E., M.B., Ch.B.,
F.R.F.P.S.Glas.

Assistant Physician, Western Infirmary, Glasgow

AND

FERGUS L. HENDERSON M.B., Ch.B.

Radiologist, Glasgow Public Health Department, etc.

(WITH SPECIAL PLATE)

The differential diagnosis of certain diseases of bones and of joints is often a matter of difficulty; aetiological factors are obscure, and treatment and prognosis rest on insecure foundations. The condition known as osteochondritis dissecans is not an extreme rarity, and has been recognized for many years, but comparatively few cases have been recorded. In the present case we believe we have grounds on which to offer an opinion as to aetiology. The results of treatment will require to be reviewed later. Trauma is still held by one group to be the principal aetiological factor, while the "non-traumatic group," to which we subscribe, disagree and believe that bone disease exists first and that following trauma the condition becomes more obvious and is diagnosed. Even then there is no unanimity of opinion as to the additional factor causing the peculiar features.

Case Record

This case, that of a well-grown lad of 15 years, was seen by one of us in August, 1935, on account of some mild pain in the right elbow-joint with slight limitation of movement. Examination did not show any joint swelling—neither fluid from a synovitis nor synovial thickening—but on moving the joint it was felt that a loose body was present which on certain movements produced pain. The history was that about six to eight weeks earlier the boy had been hit on the elbow with a cricket stump—not a violent blow, and not causing much pain or disability at the time. Clinical examination showed no other abnormality, except that he has a considerable degree of bilateral congenital nerve deafness.

X-ray examination made on August 26, 1935, showed a small bony loose body lying between the head of the radius and the corresponding articular surface of the humerus. In the outer condyle of the humerus was an irregularly shaped area of decalcification. It was found impossible to demonstrate the loose body in a lateral view, and accordingly accurate location could not be given. It was concluded that, as a result of injury, a small piece of bone had been detached from the outer condyle of the humerus and was now lying in the joint. These points are shown in the radiographs reproduced on the photogravure plate.

A surgical opinion was sought; it was decided that the loose body should be removed, and this was done. At operation the synovial membrane appeared to be perfectly healthy; there was no thickening and no excess of fluid. The loose body was composed largely of cartilage with an ossified central portion, was of the size and appearance of a white coat-button—actually 2 cm. in diameter and 0.5 cm. thick at the centre—was smooth on the flat cartilaginous surface, and rough with bony spicules on the concave surface. Some effusion occurred into the joint following operation; passive movements were not commenced for some days and were not well tolerated then. The boy was sent home after ten days with instructions to use the arm freely and to report in three months' time

—his local doctor to inform us if anything required attention sooner.

At home the boy soon played tennis—right-handed, but with difficulty in the overhead service—golfed, shot, and climbed trees. Some limitation of full flexion and especially of full extension remained, but such pain as there may have been did not require any sedatives. In general health he seemed to be very well, and was taking a good full mixed diet and living an open-air life in comfortable surroundings. The earlier signs of puberty had appeared rather more than a year previously, when he was just over 14 years of age.

Late in November, 1935—just three months after the operation—the patient reported to us again; he looked well, was well to clinical examination, and had grown and put on weight. The elbow had still some limitation of full extension; there was slight atrophy of muscles in that arm, but no pain was complained of. Further radiographs were taken; these showed that the area of decalcification in the lower end of the humerus had spread very markedly and that there was a general rarefied condition of the bones, involving the head of the radius, the outer aspect of the olecranon, and the whole of the outer half of the lower end of the humerus. The joint surfaces were involved and a considerable degree of arthritis was present. He was allowed to return to school and indulge in games generally, with the exception of Rugby football. The only incident of note was that he developed measles and was in the sanatorium for five weeks, partly as a precautionary measure and partly because he did not improve as quickly as some cases.

He was not seen by us until April, 1936, when he was again on holiday. The arm had not gained any more movement, but he used it freely and without pain. Muscle atrophy had increased, yet in general he was exceedingly well. On examination it was found that there was now some slight limitation of movement in the left elbow also, and the feeling of a loose body there. Further radiographs were taken; these showed that the active arthritic process had apparently subsided. A considerable degree of decalcification still persisted in the lower end of the humerus, and all the bones involved showed arthritic change, with complete loss of the joint surfaces, accompanied by osteophytic outgrowths.

The left elbow was examined at the same time and revealed a very obvious decalcified area in the outer condyle, similar to what had been seen in the right elbow eight months before, and a loose body was observed in the lateral view on the anterior aspect of the elbow-joint. Careful re-examination of the film of the left elbow taken on August 26, 1935, showed that this area of decalcification was present to a slight degree in the outer condyle of the humerus at that time, but this unfortunately was not noted, attention being fixed on the right elbow, which had received injury.

Further Investigations

A general radiographic examination of the body was then made with the object of discovering any additional bone abnormalities. The spinal column, hands, arms, legs, and feet all seemed normal, but there were areas of decalcification in both ischial regions. The appearances seen there were very similar to those in the elbow-joints, and on the right side there was a distinct suggestion of loose bone formation. The skull appeared to be normal, with the qualification that the sella turcica was small. It is an extremely difficult matter to be sure of any slight deviation from the normal in estimating the size of the sella turcica, but the following measurements taken of

the skull in a lateral position at a distance of thirty inches do, we hold, support the contention that the pituitary fossa is on the small side. The anterior measurement from the occipito-parietal suture to the nasion was 21.5 cm., and that from the fronto-parietal suture to the base of the posterior clinoid process 12 cm., while the greatest breadth of the sella turcica was 1 cm. Apart from these no other x-ray abnormalities were observed.

In consequence of these findings it was evident that some general bone disease existed, and in spite of the non-syphilitic appearances of the bones a Wassermann test was done and proved negative; the nerve deafness had also been suggestive. A diagnosis of osteochondritis dissecans was made: the apparent excellent general health of the boy, the lack of pyrexia, the increase rather than decrease in weight, the fact that no joint swelling existed, all disposed of the idea of tuberculosis. The recurrent loose bodies, the worm-eaten appearance of the bone whence these had come, and the bone decalcification all suggested osteochondritis. A central nervous system lesion—for example, syringomyelia—was ruled out, and no cervical ribs were present to cause nerve involvement. An investigation into the boy's inorganic retention suggested itself and was made, with the following results.

Two specimens of blood—non-oxalated—were taken at an interval of five days. The first revealed a marked lipaemia which was apparently temporary—a previous one taken for the Wassermann test and the second, or rather third, one not showing this. The principal findings were:

Plasma:

Alkali reserve	..	..	57.0% CO ₂
Inorganic phosphorus	..	..	0.005% P
Chloride	..	..	0.60%

Whole blood:

Cholesterol	..	..	0.15%
Serum calcium content	..	..	0.01% Ca (0.014% CaO)

Urine:

Acidity	..	..	22.0 c.cm. N/10 NaOH%
Urea	..	..	1.4%
Ammonia nitrogen	..	..	0.045%
Total nitrogen	..	..	0.698%
Ammonia coefficient	..	..	6.4%
CaO excretion	..	..	0.007% CaO

The principal point here is the low calcium excretion in the urine—with a normal blood calcium (as is usual even in rickets, etc.); it suggested that the absorption of calcium was defective. The boy was having a diet with plenty of lime in it and no lack of vitamin D. The other points of note are the high inorganic phosphorus—but this is usual when growth is active—and the tendency towards an acidosis.

A full examination of blood and of urine otherwise showed 90 per cent. haemoglobin, with 5,200,000 erythrocytes and 5,600 leucocytes per c.mm. The reticulocytes numbered 1.5 per cent. and the platelets 100,000. Films revealed no abnormality of cells. The low platelet count was not associated with an increase in either bleeding or coagulation time, but clot retraction was imperfect. The urine contained a haze of albumin, which was proved to be orthostatic in type, and to be serum albumin and not globulin. No tube casts were present, and the blood pressure was 100/60. The family history was inquired into, with special reference to bone disease and to haemophilia—all with negative result. The patient had not blue sclerotics, nor had any been noted in the family.

A Basis for Treatment

Consideration of all the findings suggested to us that there had been some alteration in the gonad-pituitary balance at puberty which had resulted in defective absorption of lime, with consequent withdrawal from bone

to keep up the blood calcium. No evidence of other endocrine disturbances was present, and there was no suggestion of alteration in thyroid or parathyroids. This gradual withdrawal of lime had rendered the bones more liable to damage from mild trauma, and had been the cause of the unusual sequelae.

Treatment was directed towards increase in lime intake plus vitamin D to increase absorption. Four pints of milk and 2 drachms of cod-liver oil daily were prescribed, with the result that in two weeks' time the urinary lime had risen to 0.008 per cent. CaO. This rise may appear trivial, but it is significant, and when the increase in urinary output is considered in connexion with the four pints of milk ingested, must represent considerable improvement in absorption and perhaps utilization. Three weeks later he was again examined and urinary specimens obtained. His height has increased by three-quarters of an inch in three months and his weight by 1 st. The urinary output of calcium has risen to 0.016 per cent. CaO. Further radiographs in two months' time should show whether this lime is being redeposited in bone.

Commentary

A consideration of the literature of the subject reveals that osteochondritis dissecans always occurs in adolescents, usually males, and that it may be unilateral or bilateral. There are points of similarity between it and Perthes' disease, and, like this, it would appear to become inactive once adolescence is completed and to leave the subject disabled to a lesser or greater extent. This is not in favour of a vascular origin; it might be in favour of a "bland mycotic fungoid" origin, but is much more suggestive of some imbalance of growth factors at a time when these are approaching different stages—the one its greatest activity, the other its termination.

The experimental work of Mirvish and Bosman (1928-9) on the effect of gonad extracts on blood calcium and also that on suprarenal extracts supports us in our theory that osteochondritis dissecans is in reality a temporary hypergonadism, the gonad activity causing a constant lowering of blood calcium, with consequent withdrawal of lime from bone in order to keep up the blood calcium level. The normal food calcium supply is either deficient in the circumstances or is insufficiently absorbed and utilized. The result of this process is that decalcification occurs, and, as shown in one of our radiographs, is present before trivial injury causes a dislodgement of a piece or pieces of bone and attached cartilage. It is admittedly difficult to explain why the decalcification should occur more markedly in the irregular worm-eaten way in which it does, but possibly recurrent trauma may be the explanation.

For the biochemical examination of the above case we are indebted to Dr. S. V. Telfer of the Western Infirmary, Glasgow.

LITERATURE

- King, E. S. J. (1935). *Localized Rarefying Conditions of Bone*, p. 373. Arnold and Co., London.
 Köhler, A. (1929). *Roentgenology*. Fifth edition, translated by Turnbull, p. 168. Baillière, Tindall and Cox, London.
 Brailsford, J. F. (1934). *The Radiology of Bones and Joints*, pp. 68, 138. J. and A. Churchill, London.
 Mirvish, L., and Bosman, L. P. (1928-9). *Brit. J. exp. Biol.*, **6**, 350.
 Fairbank, H. A. T. (1933). *Brit. J. Surg.*, **21**, 67.

In the first two references an extensive bibliography is given.

ADDENDUM.—Clinical examination, and radiographs of November, 1936, show that the boy is now in excellent health. He uses his left arm normally and without pain or restriction, but still has restriction of his right elbow. The radiographs show that recalcification has taken place to an extent that the bones are now normal in appearance.

ENDOMETRIAL BIOPSY

BY

ALBERT SHARMAN, M.D., B.Sc., M.C.O.G.

Assistant Surgeon, Royal Samaritan Hospital for Women, Glasgow

AND

H. L. SHEEHAN, M.D., M.Sc.

Director of Research, Glasgow Royal Maternity Hospital

(WITH SPECIAL PLATE)

For over three years certain investigations into the endometrium in sterility and anovular menstruation (Sharman, 1935) have involved frequent biopsy examinations of uterine mucosa. Much need was felt for a really satisfactory biopsy method which, from the clinical point of view, could be applied to the unanaesthetized patient without causing any real pain, and which, from the pathological point of view, would provide a sufficiently large strip of undistorted mucosa for microscopical examination.

Various techniques have been tried: curette, punch, and suction. Using a specially made narrow curette, similar to that described by Hoffmann (1934), the piece of mucosa removed was sometimes of sufficient size, but it was frequently lost in the uterus. To avoid this the stem was made hollow and suction applied through the stem with the object of holding the curetting, but this was not wholly successful. Suction methods similar to those of Klinger and Burch (1932), Novak (1935), and Tamis (1936) gave better results. The simplest and most satisfactory modification was a segment of gum-elastic catheter with the end cut off; the cannula so formed was attached to a 10 c.cm. syringe. The end of the cannula was pressed firmly against a piece of the fundal mucosa, and the piece was sucked off by means of the syringe. The fragments so obtained were often satisfactory, but were sometimes rather disappointing histologically. In addition, it was sometimes difficult and occasionally impossible to introduce the tube into the uterus, and anaesthesia was found necessary.

The New Technique

To overcome these various imperfections in the technique; naturally the procedure is even easier in multiparous women; naturally the procedure is even easier in multiparous women; naturally the procedure is even easier in multiparous women. The instrument is very simple in construction and use. The basic idea is that, whereas an ordinary curette has to be made so that it can reach the fundus, there is no need in endometrial biopsy for the curetting edge to go into the upper centimetre of the uterine cavity. It is therefore possible to place a small chamber to receive the curetting in the rational place—beyond the cutting edge. The piece of endometrium slides into this chamber as it is cut, and lies in the same position as it did in the uterus. There is no folding back of the curetting as occurs with an ordinary curette, and none of the distortion and tearing apart that is frequently produced by the suction methods. The instrument consists of a 5 cm. length of 3 mm. metal tubing attached to a shaft and handle of the same shape and size as an ordinary uterine sound. The diameter is practically constant throughout head and shaft. The part of the tube near its attachment to the shaft is filed away so that only one-third of its circumference remains. The last 1.8 cm. of the tube at the tip is left intact. The filed edge of this part of the tube, forming two-thirds of the original circumference, is sharpened and slightly undercut at the sides to form the

DOUGLAS S. STEVENSON AND FERGUS L. HENDERSON: OSTEOCHONDRITIS DISSECANS

FIG. 1.—Right elbow-joint;
August, 1935.

FIG. 2.—Right elbow-joint;
November, 1935.

FIG. 3.—Right elbow-joint;
April, 1936.

FIG. 4.—Left elbow-joint;
August, 1935.

FIG. 5.—Left elbow-joint extended;
April, 1936.

FIG. 6.—Left elbow-joint flexed;
April, 1936.

FIG. 7.—Areas of decalcification in both ischii;
April, 1936.

ALBERT SHARMAN AND H. L. SHEEHAN: ENDOMETRIAL BIOPSY

FIG. 4.—Slide with typical
biopsy sections. Five-thirds
actual size.

FIG. 5.—Histological appearances of biopsy material. Case of metrorrhagia. ($\times 80$.)