SURGICAL PROCEDURES IN GENERAL PRACTICE

This is one of a series of articles contributed by invitation

THE USE AND ABUSE OF ANTISEPTICS—II

BY

LAWRENCE P. GARROD, M.D., F.R.C.P.

AND

GEOFFREY L. KEYNES, M.D., F.R.C.S.

Use of Antiseptics on the Body

The intelligent use of antiseptics in this more difficult sphere demands clear thinking, and it is therefore necessary to distinguish between the different aims with which antiseptics are employed, and to decide whether these aims are in fact attainable at all. In the first place there is all the difference in the world between the prevention of infection by means of antiseptics and its treatment with them: treatment takes a number of forms, which may be subdivided into superficial and deep, while prevention may be concerned either with the intact and normal skin or with a recent wound. All these types of use present different problems. We may first consider the normal skin, and here a distinction is to be drawn between that of the surgeon and that of the patient.

Sterilization of the Hands

There are two kinds of bacteria on the skin: its normal saprophytic inhabitants, such as white staphylococci, which live actually in its superficial layers; and extraneous and possibly dangerous bacteria whose presence on the surface is due to accidental contamination. The removal or destruction of the latter is the object of treating the skin. It is plain that roughening of the skin due to repeated scrubbing or to chemical applications favours rather than hinders its colonization by either kind of bacteria. The operating surgeon should therefore confine himself to not more than three minutes' washing with soap and running water. The old-fashioned practice of washing for ten minutes by the clock had its justification as a protest against the carelessness of some of the older surgeons, though it would partly defeat its own purpose by making sodden the superficial layers of the skin and so tending to release the indigenous bacteria. The shorter period of washing is not open to this objection, and the scrubbing-brush should be applied to the nails only, as it will tend to damage the smooth surface of well-cared-for skin. The hands should then be thoroughly rinsed in spirit to dehydrate the surface of the skin, and dried on a sterile towel. An antiseptic is unnecessary, and may tend to injure skin that is at all sensitive. Gloves sterilized by dry heat are preferable to gloves boiled and put on wet, since maceration of the skin inside wet gloves is obviously undesirable. In emergency, such as in midwifery practice when gloves are unobtainable, an antiseptic application may take their place; dettol cream has been shown to be capable of destroying streptococci on the skin, and has the advantage of being easily carried in a form ready for use.

Sterilization of the Patient's Skin

It is generally agreed that before operation the patient's skin should first be thoroughly cleansed with ether soap

and water and then dried by the application of spirit. The addition of biniodide to the spirit does not have any advantage. A further attempt to rid the skin of bacteria may then be made by chemical means. The objection to such treatment repeated on the surgeon's hands does not have quite the same force for a single application to the patient, though it may be harmful. The antiseptic used should be one which penetrates the superficial layers of the skin and has a persistent action, and these qualities are possessed by dyes and mercurial salts. The former have the disadvantage of staining, though Tinker and Sutton's solution (5 per cent. acriflavine in 50 per cent. alcohol and 10 per cent. acetone) is possibly less objectionable from this point of view than Bonney's blue paint (0.5 per cent. brilliant green and 0.5 per cent. crystal violet in 50 per cent. spirit). The best mercurial solution is Harrington's (mercury perchloride 0.8 gramme, hydrochloric acid 60 c.cm., methylated spirit 640 c.cm., water 300 c.cm.); this may be coloured with chrysoidin Y (2 grammes per litre) if an indicator of the area covered by the application is desired.

It may be asked whether such treatment as this should be carried out on the operating table or some time before operation. Some authorities still suggest beginning it one or even two days beforehand. Previous and repeated antiseptic treatment is only indicated when the skin is infected or devitalized, and even then it must be remembered that over-treatment is always harmful. Antiseptic compresses, which make the skin sodden, are to be condemned. If the principle of covering up all skin during all operations is observed the question of applying antiseptics becomes relatively unimportant.

Of other current methods of skin treatment the following may be said. The objections to iodine, so commonly used, are three: the fact that certain skins are sensitive to it and will suffer blistering or peeling afterwards; the fact that it quickly disappears from the skin, being inactivated and destroyed by contact with blood; and its cost. It is said also that bowel which has been in contact with iodine-painted skin is liable to form adhesions. Such contact should never be allowed to occur, for, though we have no proof of this effect, iodine is certainly an irritant which would cause exudative inflammation of the peritoneum. Picric acid, another favourite application, irritates many skins, and occasionally produces such severe symptoms of poisoning as to cause the patient's death. This section may conclude with a protest against the perfunctory dab which often precedes the insertion of a hypodermic needle. If anything is worth doing at all it is worth doing properly, and if ether is used for sterilizing skin it should be rubbed on until the skin is really clean.

Preventive Application of Antiseptics to Wounds

Is it or is it not possible chemically to destroy bacteria in a recent wound and so prevent sepsis? Some surgeons say "No," others "Yes," and many have never seriously asked themselves this question. Some will countenance the use of any of a large variety of antiseptics having quite different properties and suitabilities for such a purpose.

This problem presents itself in a simple, clear, and urgent form when an accidental cut or prick occurs during

a septic operation. Here is a small wound which is known to contain dangerous bacteria. They are in the cavity of the wound, and have not yet invaded the surrounding tissues. Can they be destroyed, and if so, how? This question has been answered by abundant experiments in which both incised and punctured wounds in mice have been infected with virulent streptococci and treated by the subsequent application or injection of antiseptic solutions. Such substances as phenol and mercury perchloride are powerless to prevent the spread of infection, but compounds of the diamino-acridine series will regularly achieve this. They owe this power to an enormous capacity for restraining the growth of *Streptococcus pyogenes*, effective in dilutions of 1 in 100,000 or even higher, and to the fact that this capacity is unimpaired by the presence of serum or blood. They are also comparatively non-toxic: leucocytes retain their motility in a 1 in 1,000 solution of acriflavine in spite of being deeply stained by it. No other class of antiseptic has these properties or even approaches them. The indications in these circumstances are therefore clear. A dressing saturated in 0.2 per cent. acriflavine should be applied to the superficial scratch or cut and retained for two hours. A punctured wound is treated by introducing a hypodermic needle and injecting the solution under pressure. Some degree of permeation of the surrounding tissues is desirable, and this may be achieved by deliberately infiltrating them by subcutaneous injection.

Similar treatment may logically be applied to a recent and possibly infected wound of any kind, thorough irrigation or actual immersion ensuring that the solution reaches all parts of it. Although this is the ideal treatment the trouble and expense involved may be considered excessive for ordinary minor casualties, and the general run of accidental wounds may be cleansed and disinfected together by a solution of one of the more efficient and less toxic antiseptics of coal-tar origin; suitable preparations and strengths are dettol (5 per cent.), izal (0.5 per cent.), and cyllin (0.5 per cent.).

About other antiseptics commonly employed in these circumstances the following facts should be known. Phenol and lysol are highly destructive to tissue in concentrations adequate for antiseptic action, and weaker solutions are ineffective. Iodine is inactivated by blood. Mercury salts are exceedingly slow in action and form inactive compounds with serum proteins. Brilliant green is more actively bactericidal in a short period than acriflavine, but also considerably more toxic, although not actually irritant or caustic. All antiseptics which act by liberating chlorine or oxygen are inactivated in the presence of blood or serum; the patient and persistent use of the former, application being renewed until all visible bleeding has ceased, should, however, contribute to the disinfection of a wound.

We wish it to be understood that we are not concerned in this article with surgical procedures other than the use of antiseptics. Excision is of course the most efficient means of removing infection from a recent wound, when circumstances permit.

Antiseptic Treatment of Established Infections

A serious comparison of the depths to which various antiseptics can penetrate tissue and the depth to which bacterial invasion proceeds in different types of infection has never been made. It may safely be assumed, however, that bacteria which have once invaded the tissues become relatively inaccessible to antiseptic applications,

and these can have only a superficial effect, strikingly beneficial in but a few special cases outside our present survey. Nevertheless the antiseptic treatment of an infected wound may have its advantages in cleansing and deodorizing, in promoting fluid exudation by mild irritant action, and, what is most important, in preventing the access of further infection. These objects can all be secured by the familiar method of continuous irrigation with Dakin's solution. We are not concerned to press strongly the claims of this or any other method of treatment. Whatever is done, strong antiseptic applications to the wound itself are to be avoided, since they hinder phagocytosis and the growth of reparative tissue. It is exceedingly doubtful whether such applications as silver nitrate solutions are ever beneficial, since what may be gained in bactericidal effect is counterbalanced by tissue necrosis, producing a pabulum for further bacterial growth. It is most noteworthy in connexion with wound treatment that not all bacteria are equally susceptible to the action of the same antiseptic. Streptococci and staphylococci are many times more susceptible, as measured in terms of effective dilution, than are some Gram-negative bacilli to the action of acriflavine; *B. proteus* and *B. pyocyaneus* are highly resistant to this and to many other antiseptics. For the treatment of wounds, as opposed to their preventive disinfection, acriflavine and other dyes have the disadvantage of being absorbed by the material of dressings, so that they will only be active in the presence of a large excess of the solution. Their possible action is still further hindered by intimate admixture with an oily basis, as in the acriflavine emulsion of the *B.P.C.*, and this elaborate preparation has no demonstrable antiseptic action whatever. A similar objection applies to the use of "bipp," and we have already implied that in our opinion the claims of iodoform to be regarded as an efficient antiseptic rest on a very insecure basis.

In addition to the treatment of infected wounds, antiseptic irrigation usually plays a prominent part in treating infections of the conjunctiva, the upper air passages, the urethra and other portions of the urinary tract, and the vagina, while antiseptics are also extensively employed for affections of the skin. With these specialized uses it is not our purpose to deal, but it may be pointed out that similar principles apply to them. There is the same necessity to guard against doing damage which outweighs the benefit conferred, and the same unlikelihood of reaching bacteria other than those in surface exudates. The conjunctiva affords a striking example of the fact that prevention by antiseptics is much more certain than cure: Credé's method prevents ophthalmia neonatorum because it attacks the gonococcus before the tissues have been invaded, and it would probably succeed equally well if some other and less irritating antiseptic than silver nitrate were used.

Deep Antisepsis

Hitherto we have considered chiefly those methods of employing antiseptics which involve application to a surface. It must not be forgotten that attempts of many kinds are made by chemical means to destroy bacteria more deeply situated, attempts which consist in administering the antiseptic either by the mouth or by some form of injection. The most ambitious of these aim at disinfection of the blood in septicaemia, others at disinfecting the urinary, intestinal, or biliary tracts.

Circulatory Antiseptics.—It is beyond question that *p*-aminobenzenesulphonamide (sulphanilamide), which is derived from and has superseded the original prontosil,

administered in adequate doses by the mouth, is capable of destroying haemolytic streptococci in the blood stream. This astonishing action, fully attested by both laboratory experiment and clinical experience, stands in strong contrast to that of every reagent previously employed for this purpose. The folly and futility of injecting intravenously such reagents as eusol and mercury salts are now generally recognized. Mercurochrome had a vogue for this purpose, in spite of the repeated demonstration that pyogenic cocci would actually grow in blood containing this drug in a concentration hundreds of times greater than can be attained therapeutically in the circulation. Other, and otherwise more efficient, mercury compounds are useless for this purpose. A few dyes might be expected to exert some effect, but they are probably eliminated too rapidly to attain it: one clue to the success of sulphanilamide is that it remains in maximal concentration in the blood for six hours. It is to be hoped that the discovery of this drug will banish all substances previously used as circulatory antiseptics into the limbo of forgotten things.

Urinary Antiseptics.—Here again the advent first of mandelic acid and then of sulphanilamide has largely superseded previous treatments of doubtful efficiency. Two other antiseptics have a demonstrable action in the urinary tract: hexamine, which is dependent, like mandelic acid, on the maintenance of a low pH; and hexyl-resorcinol, the behaviour of which is unaccountably erratic.

Intestinal Antiseptics.—The coal-tar preparations which are sometimes given to colostomy patients and in various other circumstances in order to suppress bacterial activity in the lower bowel do not reduce the bacterial population of the colon, and have indeed never been shown to reach it. So far as we are aware an antiseptic that is neither absorbed from the bowel nor rendered inert by chemical combination with its contents has yet to be found.

Biliary Antiseptics.—We are not aware of a convincing demonstration that any drug acts as a biliary antiseptic. Hexamine has been much lauded for this purpose; this substance is not itself bactericidal at all, and acts only by liberating formaldehyde under the influence of acid. A degree of acidity adequate for this effect exists nowhere in the body except in the stomach and the urinary tract. For the same reason hexamine is quite inert in the central nervous system.

Conclusion

It has been possible in the present review to deal only with better-known antiseptics and their commoner and more important uses. Essential oils, boric and other acids, soaps, metals, and even the excrement of maggots have been used for their antiseptic properties. For each of the proprietary antiseptics of coal-tar origin which we have named there are a dozen others, similar in composition but in some cases less familiar and unsupported by the same variety of testimony to their value. Since no authority in this country has undertaken the task of examining and adjudicating upon preparations of this kind the medical profession in general has no means other than clinical trial of knowing what their merits are. In this as in other directions which we have indicated the intelligent use of antiseptics is rendered difficult by lack of information. Order is more likely to be brought into the present chaos of this subject if it is more often approached in the manner we have attempted here—that of inquiring into the evidence that the properties of an antiseptic are suited to a particular task.

NATIONAL FITNESS AND THE CARE OF THE INSANE

BY

SIR ROBERT ARMSTRONG-JONES, M.D.,
D.Sc., F.R.C.P.

A striking demonstration of physical exercises combined with musical rhythm was recently given by the Margaret Morris Society before an audience of teachers, nurses, representatives of the Eugenics Society, and medical men, some of the last-named associated with the care and treatment of the insane and coming from a distance so far from London as Cumberland. This most interesting exposition of the value of graded muscular movements, lasting nearly two hours, was presented on December 1 at 31, Cromwell Road, S.W.7, the London Institute of the movement. It was fully explained by Miss Margaret Morris and her staff, who had already directed their remedial treatment classes in several mental hospitals and private homes for the care and treatment of the insane. The attractive demonstration included the use of general and special muscular movements, easy of execution and pleasing to watch, by which to secure the balance of the body, graceful postures, and mental relaxation. They taught the performance of physiological muscular contractions, which were carried out without effort, and a normal poise was maintained which is deplorably wanting in all mental cases. Recently the advocacy of physical fitness has been taken up with great earnestness by the Government in pressing forward the scheme for national fitness. Local authorities on the one hand and a great array of voluntary workers on the other have been engaged in rousing public interest in activities for an improvement in the mental and physical condition of the people, but in no section is this more necessary than in the population of our mental hospitals.

Members of Parliament who desire to urge primarily the question of nutrition—free food and milk—need have no anxiety to-day about the food provided in the mental hospitals, for in these the dietary has been fully considered by the responsible local authorities. During the last twenty years the maintenance rate in the public mental hospitals has been more than doubled, and the nutrition of those under treatment appears to be satisfactory; but the general improvement in bodily conditions through the judicious use of muscular exercise and movements has perhaps not received the special attention required as an aid in remedial treatment. It is not yet realized that the Lunacy Act, 1930, makes it possible to receive into mental hospitals cases in their early stages voluntarily—that is, without the doctor's certificate of insanity. Naturally it is obligatory upon those responsible for the treatment—apart from the question of public duty, personal responsibility, or humanity—to do their very best towards the patient's rehabilitation and recovery.

The Numbers of Mental Patients

The question is a very big one, for in the year 1936—the last for which we have any record—there were under care in our mental hospitals over 130,000 persons suffering from mental disease. Of these more than 25,000 were new cases, a fourth of them—upwards of 6,000—being voluntary patients. Also, of the 82,000 cases of mental deficiency 4,000 were new cases admitted during 1936 (although none of these was a voluntary case). Every effort should be (and is) made for their mental improvement and restoration—quite apart from the urge arising out of a fellow feeling and sympathy, which their condition entitles them to receive.

It is not criticism of the past we require but suggestions for the future; we should begin, therefore, by asking, "What sort of cases are likely to improve by physical treatment?"—that is, by muscular movements, bodily