positive findings in the nervous system. Extending from the coracoid process four and a half inches down the left arm was an old surgical scar. The left arm was two and three-quarter inches shorter than the right, the shortening being limited to the humerus and radius and the ulna. An old scar, two inches long, was seen on the dorsum of the left forefinger, curving proximally to the radial side and extending through the nail, which was deformed.

The patient was admitted to the Gateshead Mental Hospital as a voluntary patient and there given atropine in high dosage, according to the technique of Hall (1935, 1937) after Kleeman. Complete relief from her symptoms was obtained with 4½ mg. daily, spread over three doses. Observation and routine psychological examination produced nothing relevant.

Commentary

Cases of Parkinsonism following poisoning by manganese, gold, cyanide, and carbon disulphide have been recorded, but I have been unable to trace one following poisoning by the substances used in this case for the treatment of the growth at the head of the humerus.

Only upon inquiry was a history obtained of weakness and pain in the interval between the injury and the onset of the tremor, and these had not been regarded by the patient as abnormal. Thus the first feature of Naville and de Morsier is poorly exemplified. But the emphatic story of the tremor beginning soon after the injury, appearing at first only in the injured finger and remaining clearly limited to it for some weeks, is surely significant; and indeed no previously recorded case has shown such a precise local relation between the trauma and the first tremor. Because of this and of such a definite history of excellent general health, and in view of the cumulatively impressive accounts of the condition hitherto published, it would seem unwarranted to postulate an untraceable attack of encephalitis lethargica. Coincidence of occult encephalitis and injury does not explain the previously published cases. Naville and de Morsier (1932) state that "the constant succession of these phenomena—injury, pain, hypertony, and tremor of the injured limb; extension to the homolateral limb, then to the contralateral limbs—enables one to declare that it is not possible for it to be in the nature of coincidence."

The effect of high atropine dosage has not been mentioned in the cases published, and it is pleasant to be able to record its satisfactory action in this case, but it will surely mask any spread of symptoms to the opposite limbs. When the patient was last seen (November, 1937) she was persisting with her atropine treatment, had no complaints, and was able to do a hard day's work.

I am grateful to Dr. C. B. Bamford for his interest and for his permission to publish this case, and to Dr. F. J. Nattrass for confirming the diagnosis of Parkinsonism.

REFERENCES

- Faure-Beaulieu, M. (1932). *Presse méd.*, **40**, 1327.
Hall, A. J. (1935). *British Medical Journal*, **2**, 555.
— (1937). *Ibid.*, **1**, 795.
Hodgson, J. C. (1937). *Ibid.*, **2**, 58.
Naville, F., and de Morsier, G. (1932). *Ann. méd. lég.*, **12**, 169.
Walker, G. F. (1937). *British Medical Journal*, **2**, 65.

W. B. Hubbard (*Arch. Ophthalm.*, Chicago, August, 1937) has experimented with caustic burns in rabbits using 32 per cent. sulphuric acid and 20 per cent. sodium hydroxide. In all cases one eye was irrigated with tap-water, the other being washed out with either 2 per cent. sodium bicarbonate or 2 per cent. acetic acid to neutralize the caustic. In the case of acid burns the best results were obtained with tap-water, while alkaline lesions were best treated with weak acid irrigations. This difference is probably due to the fact that acid proteinates are insoluble and alkaline proteinates are soluble.

RETENTION OF URINE FOLLOWING EXCISION OF THE RECTUM

BY

H. T. SIMMONS, Ch.M., F.R.C.S.

Research Fellow, The Royal Infirmary, Manchester; Honorary
Assistant Surgeon, Christie Hospital and Holt
Radium Institute

Retention of urine following excision of the rectum is so frequent that it is regarded as a usual post-operative complication. It occurs in both sexes, and generally three or four days elapse before the patient is able to empty the bladder voluntarily. Occasionally the resumption of voluntary micturition is delayed for weeks or even months, and this condition appears to be more especially associated with the removal of high and extensive growths by the perineal route. The danger of retention of urine is the supervention of sepsis, which, though not always fatal, causes considerable morbidity and undermines the patient's strength. Minor operations on the perineum and anal region often give rise to a retention of urine which rarely lasts more than twenty-four hours and which responds to simple treatment. This retention is thought to be due to reflex nervous inhibition. After removal of the rectum, however, the condition persists for three or four days, and occasionally for weeks, and must be due to a more definite interference with the nervous mechanism.

Sympathetic and Parasympathetic Innervation

The filling of the bladder is dependent upon its sympathetic nerve supply, which keeps the internal sphincter contracted and so allows the accumulation of urine. Abolition of this nerve supply, or its inhibition, allows the sphincter to relax and the urine can escape. The emptying is due to the action of the parasympathetic nerves, which cause the bladder to contract and at the same time relax the internal sphincter. Destruction of this nerve supply reduces the contractile power of the bladder and leaves the internal sphincter tonically contracted by the action of the sympathetic nerves, which are no longer opposed. Thus retention of urine occurs.

Normally voluntary control of micturition exists because both these nerve supplies have cortical connexions. If, however, both are destroyed the bladder does not remain completely paralysed. After a latent period the musculature contracts to the stretch-stimulus of the accumulating urine; at the same time the internal sphincter is relaxed by the intrinsic nerve plexus. This automatic bladder is an excellent example of how smooth muscle adapts itself to carry on its original function after the loss of its extrinsic nerve supply. The presacral nerve, lying behind the peritoneum over the promontory of the sacrum, brings the sympathetic nerve supply of the bladder into the pelvis from the lumbar ganglionated cords. This nerve divides into the two hypogastric nerves, which, passing round each side of the pelvis, bring the supply to the pelvic plexuses, joining them near the groove between the rectum and the base of the bladder. The fibres pass through these plexuses and are distributed to the trigone and internal sphincter.

The parasympathetic supply to the bladder is brought by the nervi erigentes. In seven pelvises dissected in the anatomy department of the University of Manchester I found that the nervi erigentes arose mainly from the second and third sacral nerves. In only three cases was there any contribution from the fourth sacral nerve. A pelvic plexus is formed on each side of the rectum by

the nervi erigentes, and fibres could be traced down to the anus and as high as the recto-sigmoid junction. The plexuses were attached to the rectum only by their fibres of distribution, and, passing forwards, they communicated with each other and received the sympathetic fibres near the prostate and bladder base. Dissection of the plexuses, and the tracing of the fibres as completely as possible, showed that those arising from the second sacral nerve appeared mainly to supply the bladder, whereas the rectum was chiefly supplied by the lower fibres of the third sacral nerve and the contribution from the fourth nerve.

Operative Injury to Nerves

In the dissecting room the rectum may be removed without the risk of damage to the nervi erigentes or to the pelvic plexuses. The hypogastric nerves which carry the sympathetic supply are in no danger of being injured. The removal of a carcinomatous rectum in a living patient demands, however, a much wider dissection. It was found in the cadaver that the dissection necessary would assuredly destroy the contribution to the pelvic plexuses from the fourth sacral nerves and the lower fibres from the third nerves. During the operation the upper fibres from the third sacral nerve were in constant jeopardy, depending mainly on the height and lateral extent of the excision. Damage to the fibres arising from the second sacral nerve could be avoided, but where a high removal was attempted they were in danger of being divided.

The dissections suggested that in perineal excision of the rectum there is little interference with the sympathetic supply of the bladder, but that the nervi erigentes always suffer some damage. The parasympathetic fibres arising from the fourth, and the lower fibres arising from the third, sacral nerves are usually divided, but as they appear mainly to supply the rectum their division does not affect the bladder to any great extent. The supply to the bladder from the upper fibres of the third sacral nerve suffers from traction, and to a varying degree from actual severance of fibres. The supply from the second sacral nerve is injured usually only by traction, but in the removal of high growths it may be severed. The effect of traction will be to paralyse temporarily, and of severance to paralyse permanently, the bladder parasympathetic supply. Hence the action of the sympathetic nerves will be unopposed, giving in the first case a temporary and in the second case a more or less permanent tonic contraction of the internal sphincter. The resulting retention of urine will be temporary in the first instance and more lasting in the second; but it will hardly become permanent, as the smooth muscle will slowly adjust itself to the altered conditions and give an automatic type of bladder.

Previous attempts to deal with this retention of urine have been mainly concerned with the methods of emptying the bladder when the condition is established. The best results appear to have been obtained by Cuthbert Dukes (1928-9), who suggested the pre-operative administration of two doses, at weekly intervals, of a vaccine prepared from the bacteria present in these urinary infections. He inserted a catheter at the time of operation and afterwards emptied the bladder frequently. The catheter was connected to an apparatus which allowed repeated bladder-washing to be performed, and had a column of anti-septic fluid as a seal for the catheter instead of the customary wooden peg. He demonstrated an immediate improvement in results; but even in his own cases one-third of the males and all the females developed some urinary infection. The danger of urinary infection would

cease to exist if we could avoid the introduction of a catheter into the bladder.

Cause of the Retention

It has been shown that the cause of the retention of urine is a relative overaction of the sympathetic nerve supply of the bladder. This appears to be due to weakening of the parasympathetic supply by traction on its fibres during the operation or, in extreme cases, by actual severance of the nerves. The overaction could be avoided by dividing the sympathetic supply either at the time of excision or at a previous operation. This is easily achieved by the division of the presacral nerve, and we know from the numerous cases in which it has been done for gynaecological pain that it is followed by no untoward effects upon the bladder, even though that organ has an intact parasympathetic supply.

This presacral neurectomy involves no separate intervention. Its performance is very easy and it adds no more than a few minutes to the time taken by the exploratory laparotomy and colostomy which must precede the perineal excision. The presacral nerve is found behind the peritoneum overlying the promontory of the sacrum. An incision two inches in length is made through this peritoneum, the pelvic colon being held well over to the left side. The nerve is formed near the bifurcation of the aorta by the junction of three descending roots, and it passes down into the pelvis either as a single cord or, more commonly, as a loose plexus. Rather more than one inch of the nerve is excised, and to make sure that all the fibres are divided it is better to clear the angle between the right common iliac artery and the left common iliac vein completely of all fatty and fibrous tissue down to but not below the sacral promontory.

Results

A consecutive series of twenty-two cases of perineal excision of the rectum have had their presacral nerves excised at the preliminary laparotomy. Fifteen were males and seven were females, and there was no mortality in the series. In no case was a catheter inserted into the bladder as a guide to the urethra, as it was felt that this was unnecessary and only introduced a risk of sepsis. Of the seven women four passed urine naturally and three had to be catheterized, one for two days and the other two for three days. Of the fifteen men nine passed urine of their own accord; four were catheterized for one or two days, and the other two for five days. In each of these cases the prostate gland was larger than normal. All the patients passing urine normally escaped urinary infection.

These results, although not completely solving the problem of post-operative retention of urine after perineal excision of the rectum, suggest that a preliminary division of the sympathetic nerve supply of the bladder is of considerable value in avoiding this troublesome complication.

Summary

Retention of urine following perineal excision of the rectum appears to be caused by a nervous imbalance—a sympathetic preponderance due to trauma to the parasympathetic supply (the nervi erigentes).

Resection of the presacral nerve at the time of the preliminary laparotomy and colostomy has been followed by a considerable diminution in the incidence of post-operative retention of urine.

REFERENCE

Dukes, Cuthbert (1928-9). *Proc. roy. Soc. Med.*, 22, 259.