A L Tan, Academic Unit of Musculoskeletal Disease, University of Leeds and Chapel Allerton Hospital, Leeds, UK

S Shankaranarayana, Department of Rheumatology, Calderdale Royal Hospital, Halifax, UK

J Madden, Department of Rheumatology, Calderdale Royal Hospital, Halifax, UK

P Emery, Academic Unit of Musculoskeletal Disease, University of Leeds and Chapel Allerton Hospital, Leeds, UK

M F McDermott, Leeds Institute of Molecular Medicine, St James's University Hospital, Leeds, UK

Correspondence to: Professor Dennis McGonagle, Academic Unit of Musculoskeletal Disease, Chapel Allerton Hospital, Chapeltown Road, Leeds LS7 4SA, UK; d.g.mcgonagle@leeds.ac.uk

Accepted 3 June 2007

REFERENCES

- 1 Kaufmann P, Torok M, Hanni A, Roberts P, Gasser R, Krahenbuhl S. Mechanisms of benzarone and benzbromarone-induced hepatic toxicity. *Hepatology* 2005;**41**:925–35.
- 2 Pui CH. Urate oxidase in the prophylaxis or treatment of hyperuricemia: the United States experience. *Semin Hematol* 2001;**38**(4 suppl 10):13–21.
- 3 Chu R, Lin Y, Reddy KC, Pan J, Rao MS, Reddy JK, et al. Transformation of epithelial cells stably transfected with H₂O₂-generating peroxisomal urate oxidase. *Cancer Res* 1996;**56**:4846–52.
- 4 Akahoshi T, Murakami Y, Kitasato H. Recent advances in crystal-induced acute inflammation [review]. *Curr Opin Rheumatol* 2007;**19**:146–50.
- 5 Martinon F, Petrilli V, Mayor A, et al. Gout-associated uric acid crystals activate the NALP3 inflammasome. *Nature* 2006;**440**:237–41.
- 6 Murakami Y, Akahoshi T, Hayashi I, Endo H, Hashimoto A, Kono S, et al. Inhibition of monosodium urate monohydrate crystal-induced acute inflammation by retrovirally transduced prostaglandin D synthase. *Arthritis Rheum* 2003;**48**:2931–41.
- 7 Roberge CJ, Grassi J, De Medicis R, Frobert Y, Lussier A, Naccache PH, et al. Crystal-neutrophil interactions lead to interleukin-1 synthesis. *Agents Actions* 1991;**34**:38–41.
- 8 Di Giovine FS, Malawista SE, Nuki G, Duff GW. Interleukin 1 (IL 1) as a mediator of crystal arthritis. Stimulation of T cell and synovial fibroblast mitogenesis by urate crystal-induced IL 1. *J Immunol* 1987;**138**:3213–18.
- 9 So A, Desmedt T, Revaz S, Tschopp J. A pilot study of IL-1 inhibition by anakinra in acute gout. *Arthritis Res Ther*, 2007;**9**:R28 [Epub ahead of print].
- 10 Tausche AK, Richter K, Grassler A, Hansel S, Roch B, Schroder HE. Severe gouty arthritis refractory to anti-inflammatory drugs: treatment with anti-tumour necrosis factor alpha as a new therapeutic option. *Ann Rheum Dis* 2004;**63**:1351–2.

Tumour necrosis factor receptor 2 (TNFRSF1B) association study in Sjögren's syndrome

Corinne Miceli-Richard*, Philippe Dieude*, Eric Hachulla, Xavier Puechal, François Cornelis, Xavier Mariette

Primary Sjögren's syndrome (pSS) is a complex disease involving both genetic and environmental factors. Among the genetic susceptibility factors, *HLA-DRB1* has been extensively studied as a strong candidate gene and the association of *HLA-DRB1*0301* (DR3) with pSS has been frequently reported among Caucasians.^{1,2} As the Tumour Necrosis Factor α (*TNF α*) gene is located within the *HLA* region, its contribution to disease susceptibility has also been studied. We have previously reported a significant association of *TNF-308A* allele with pSS, in strong linkage disequilibrium with *HLA-DRB1*03*.³ This association was restricted to patients with anti-SSB antibodies. *TNF* exerts its action through two cell surface receptors, *TNF* receptors 1 and 2 (*TNFR1* and *TNFR2*, 55 and 75 kDa respectively). Due to the functional interaction between *TNF* and *TNFR2*, we considered this receptor (also named *TNFRSF1B*) as another interesting candidate gene for the genetic susceptibility to pSS. *TNFR2 T676G* polymorphism replacing a methionine by an arginine (*TNFR2 196 M/R*) is functional as the cytotoxic activity induced by *TNFR2 196R* (*TNFR2 696 G* allele) is increased.⁴ Moreover, *TNFR2 T676G* polymorphism has been previously associated with the genetic susceptibility to other autoimmune diseases: familial, but not sporadic, rheumatoid arthritis (RA)⁵ and systemic lupus erythematosus (SLE).⁶

We carried out a case-control association study of *TNFR2 T676G* polymorphism among 119 unrelated patients with pSS according to European-American consensus group (46 patients without autoantibodies (Ab), 33 patients with anti-SSA Ab only, 40 patients with anti-SSA and anti-SSB Ab) and 95 healthy controls. Both patients and controls were of Caucasian origin. *TNFR2 T676G* polymorphism was genotyped by

polymerase chain reaction–restriction fragment length polymorphism analysis (*Nla* III).

No significant differences in allele and genotype frequencies of *TNFR2 T676G* polymorphism were detected between patients with pSS and controls (Table 1). *TNFR2 T676G* polymorphism was not involved in the genetic predisposition to a specific pattern of autoantibody secretion ($p = 0.61$). No association was found with extraglandular involvement ($p = 0.34$). Interestingly, there was a trend in favour of an association with joint involvement (arthritis and arthralgia) ($p = 0.06$). An epistatic effect between *TNF -308A/G* and *TNFR2 T676G* polymorphisms was then looked for, but not evidenced, among 55 pSS patients.

Table 1 Allelic and genotypic frequencies of *TNFR2 T676G* polymorphism in 119 patients with pSS and 95 controls

<i>TNFR2 T676G</i> Allele frequencies	pSS n = 238	Controls n = 190	p	Odds ratio (95% CI) pSS versus controls
676T (%)	191 (80.3)	149 (78.4)	NS	1.12 (0.7 to 1.79)
676G (%)	47 (19.7)	41 (21.6)		
Genotype frequencies	n = 119	n = 95		
TT	72 (60.5)	56 (58.9)	NS	0.85 (0.62 to 1.15)
TG	47 (39.5)	37 (38.9)	NS	1.17 (0.86 to 1.60)
GG	0	2 (2.2)		

pSS, primary Sjögren's syndrome; CI, confidence interval; NS, not significant.

This study failed to demonstrate an association of *TNFR2* T676G polymorphism with pSS. These results, together with the lack of efficacy of infliximab in pSS⁷ rather suggest a poor involvement of the TNF/TNF receptor system in the susceptibility to pSS.

To date, among rheumatic diseases, *TNFR2* T676G polymorphism association with lupus led to conflicting results.^{4,6,8,9} A recent case-control study performed in an Asian population of patients also failed to demonstrate any association of *TNFR2* T676G polymorphism with SLE.¹⁰ However, this study also provided a metaanalysis of seven case-control studies in eight different ethnic populations. Stratification by ethnicity revealed a significant association among Asians while the effect of *TNFR2* T676G polymorphism on SLE was not significant in two case-control studies involving Caucasians (OR = 0.99, 95%CI = 0.68 to 1.45; *p* = 0.96). Thus, as observed among Caucasian SLE patients, our study provides evidence suggesting that *TNFR2* T676G polymorphism is not involved in the genetic predisposition to pSS in French Caucasian patients.

Authors' affiliations

Corinne Miceli-Richard, Xavier Mariette, Service de Rhumatologie, Institut Pour la Santé et la Recherche Médicale (INSERM) U802, Hôpital Bicêtre, Assistance Publique-Hôpitaux de Paris (AP-HP), Université Paris 11, Le Kremlin Bicêtre, France

Philippe Dieude, François Cornelis, GenHotel, Université d'Evry-Paris 7, Evry Genopole, France

Eric Hachulla, Service de Médecine Interne, Hôpital Claude-Huriez, CHU de Lille, Université Lille 2, France

Xavier Puechal, Service de Rhumatologie, Centre Hospitalier du Mans, Le Mans, France

Competing interests: None.

*These authors contributed equally to the study.

Correspondence to: Corinne Miceli-Richard, Service de Rhumatologie, Institut Pour la Santé et la Recherche Médicale (INSERM) U802, Hôpital

Bicêtre, Assistance Publique-Hôpitaux de Paris (AP-HP), Université Paris 11, Le Kremlin Bicêtre, France; corinne.miceli@bct.aphp.fr

Accepted 22 May 2007

REFERENCES

- 1 Arnett FC, Hamilton RG, Reveille JD, Bias WB, Harley JB, Reichlin M. Genetic studies of Ro (SS-A) and La (SS-B) autoantibodies in families with systemic lupus erythematosus and primary Sjögren's syndrome. *Arthritis Rheum* 1989;**32**:413-9.
- 2 Bolstad AJ, Wassmuth R, Haga HJ, Jonsson R. HLA markers and clinical characteristics in Caucasians with primary Sjögren's syndrome. *J Rheumatol* 2001;**28**:1554-62.
- 3 Gottenberg JE, Busson M, Loiseau P, Dourche M, Cohen-Solal J, Lepage V, et al. Association of transforming growth factor beta1 and tumor necrosis factor alpha polymorphisms with anti-SSB/La antibody secretion in patients with primary Sjögren's syndrome. *Arthritis Rheum* 2004;**50**:570-80.
- 4 Morita C, Horiuchi T, Tsukamoto H, Hattori N, Kikuchi Y, Arinobu Y, et al. Association of tumor necrosis factor receptor type II polymorphism 196R with Systemic lupus erythematosus in the Japanese: molecular and functional analysis. *Arthritis Rheum* 2001;**44**:2819-27.
- 5 Dieude P, Petit E, Cailleau-Moindraut S, Osorio J, Pierlot C, Martinez M, et al. Association between tumor necrosis factor receptor II and familial, but not sporadic, rheumatoid arthritis: evidence for genetic heterogeneity. *Arthritis Rheum* 2002;**46**:2039-44.
- 6 Komata T, Tsuchiya N, Matsushita M, Hagiwara K, Tokunaga K. Association of tumor necrosis factor receptor 2 (TNFR2) polymorphism with susceptibility to systemic lupus erythematosus. *Tissue Antigens* 1999;**53**:527-33.
- 7 Mariette X, Ravard P, Steinfeld S, Baron G, Goetz J, Hachulla E, et al. Inefficacy of infliximab in primary Sjögren's syndrome: results of the randomized, controlled Trial of Remicade in Primary Sjögren's Syndrome (TRIPSS). *Arthritis Rheum* 2004;**50**:1270-6.
- 8 Lee EB, Yoo JE, Lee YJ, Choi YJ, Park KS, Song YW. Tumor necrosis factor receptor 2 polymorphism in systemic lupus erythematosus: no association with disease. *Hum Immunol* 2001;**62**:1148-52.
- 9 Tsuchiya N, Kawasaki A, Tsao BP, Komata T, Grossman JM, Tokunaga K. Analysis of the association of HLA-DRB1, TNFalpha promoter and TNFR2 (TNFRSF1B) polymorphisms with SLE using transmission disequilibrium test. *Genes Immun* 2001;**2**:317-22.
- 10 Horiuchi T, Kiyohara C, Tsukamoto H, Sawabe T, Furugo I, Yoshizawa S, et al. A functional M196R polymorphism of tumor necrosis factor receptor type 2 is associated with systemic lupus erythematosus: A case-control study and a meta-analysis. *Ann Rheum Dis* 2007;**66**:320-4.

International survey on the diagnosis and management of gout

M Doherty, T Bardin, E Pascual

The pathogenesis of urate crystal deposition is reasonably well understood, and with appropriate urate-lowering therapy (ULT) and lifestyle advice, the objective of management is cure. Nonetheless, many patients with gout continue to experience frequent and recurrent episodes of gout and progression of their disease. This is because the condition is often misdiagnosed, or diagnosed late, and treatment is frequently suboptimal.^{1,2} These concerns led to the development of evidence-based recommendations relating to the diagnosis and management of gout by a task force from the European League Against Rheumatism (EULAR). The EULAR recommendations were developed using a combination of research-based evidence and expert consensus, and provide a valuable resource for physicians.^{1,2}

The current diagnosis and management of gout was investigated in a survey conducted among delegates attending EULAR 2006. Delegates visiting the commercial stands were

invited to complete a written questionnaire of 12 questions designed to evaluate awareness of the impact of gout, the goals of treatment, and patterns of clinical practice. A total of 741 respondents (6.7% of attendees), predominantly rheumatologists from Europe, completed the questionnaire.

The results indicate a good level of understanding concerning gout and its management, but also highlight some marked variations in clinical practice.

Respondents reported using both the measurement of serum uric acid (sUA) (97%) and the examination of synovial fluid (91%) to confirm suspected gout. Only 15% said they would prescribe ULT when the diagnosis of gout was made, most reported that they would usually wait for the next gout attack (24%) or for at least two further attacks (32%) to occur before initiating treatment. Whereas 29% said they would initiate ULT during an attack, the remainder confirmed they would wait until after an attack, and most (60%) would then begin therapy

Ann Rheum Dis 2007;**66**:1685-1686. doi: 10.1136/ard.2007.071530