Figure 1 Immunostaining of CNVMs by cell-specific marker and HIF. Immunostaining was performed to confirm the cellular source of expression for HIF-1 α and HIF-2 α . (A–D) Serial sections were immunostained by CD34, CD68, HIF-1 α and HIF-2 α . The vascular endothelium cells were stained by CD34 (A), and some of the endothelial cells were stained by HIF-1 α and similarly by HIF-2 α . (C and D) Macrophages, identified by immunostaining of CD68 (B), were detected abundantly in all CNVMs and were located uniformly in stroma. Most of the macrophages that were immunostained were immunoreactive to HIF-1 α and HIF-2 α (C and D). A partially intact monolayer of retinal pigment epithelium (RPE) cells located on one side of the CNVMs was immunostained by pancytokeratin (E). However, no overlap was observed with HIFs staining (F and G) and RPE cells. Scale bars, 50 μ m.

Each CNVM specimen was immediately embedded in OCT compound (O.C.T. Compound; Sakura Finetek Co., Ltd., Tokyo, Japan), rapidly frozen and prepared for avidin-Biotin complex immunohistochemistry (VECTASTAIN ABC kit, Vector Laboratories, Burlingame, CA).

HIF-1 α and HIF-2 α were detected in 5 and 6 eyes of 5 eyes, respectively, as revealed by immunolabelling with anti-HIF-1 α and anti-HIF-2 α antibody (Novus Biologicals, Littleton, CO). Neovascular vessels immunostained by CD34 (Biomedex, Foster, CA) were detected in all CNVMs and located in stroma, and half of the endothelial cells were immunostained by both HIF proteins (fig 1A–B). CD68-positive (Ylem, Roma, Italy) macrophages were located uniformly in stroma in all CNVMs and almost all macrophages were stained with both HIF antibodies (fig 1C–D). All specimens contained a partially intact monolayer of RPE cells located on one side of the CNVMs and immunostained by pancytokeratin (Sigma, St. Louis, MO) (fig 1E–F). No overlap was observed with HIF proteins and RPE cells (fig 1E–F). The distribution pattern of HIF-1 α and HIF-2 α in the CNVMs was similar in all specimens.

In summary, HIF-1 α and HIF-2 α are detected in the endothelium and macrophages. The HIF transcription factors activate the expression of the *VEGF* gene in response to hypoxic conditions. VEGF, in turn, induces angiogenesis.^{7–8} RPE cells, endothelial cells and macrophages are VEGF-positive in CNVMs from AMD patients,^{9–10} suggesting that VEGF is involved in CNVM formation.¹⁰ Our results, together with previous results, may indicate that HIFs in endothelial cells induce VEGF, thereby contributing to the development of CNVM. In the current study, however, expression of HIF proteins was not detectable in RPE cells, raising the possibility that other mechanisms are involved in the production of VEGF in these cells. Further studies are needed to explore the possible causal relationship between hypoxia and the expressions of HIF proteins.

Yuji Inoue, Yasuo Yanagi, Kyosuke Matsuura, Hidenori Takahashi, Yasuhiro Tamaki, Makoto Araie

Department of Ophthalmology, University of Tokyo School of Medicine, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan

Correspondence to: Yuji Inoue or Yasuo Yanagi, Department of Ophthalmology, University of Tokyo School of Medicine, 7-3-1 Hongo, Bunkyo-ku, Tokyo 113-8655, Japan; yino-tyk@umin.ac.jp or yanagi-tyk@umin.ac.jp

doi: 10.1136/bjo.2006.111583

Accepted 1 March 2007

Competing interests: None declared

References

- 1 Wang GL, Jiang BH, Rue EA, *et al.* Hypoxia-inducible factor 1 is a basic-helix-loop-helix-pas heterodimer regulated by cellular O₂ tension. *Proc Natl Acad Sci USA* 1995;**92**:5510–4.
- 2 Pugh CW, Ratcliffe PJ. Regulation of angiogenesis by hypoxia: Role of the HIF system. *Nat Med* 2003;**9**:677–84.
- 3 Ozaki H, Yu AY, Della N, *et al.* Hypoxia inducible factor-1 α is increased in ischemic retina: Temporal and spatial correlation with VEGF expression. *Invest Ophthalmol Vis Sci* 1999;**40**:182–9.
- 4 Morita M, Ohneda O, Yamashita T, *et al.* HIF/HIF-2 α is a key factor in retinopathy of prematurity in association with erythropoietin. *Embo J* 2003;**22**:1134–46.
- 5 Lambert HM, Capone A Jr, Aaberg TM, *et al.* Surgical excision of subfoveal neovascular membranes in age-related macular degeneration. *Am J Ophthalmol* 1992;**113**:257–62.
- 6 Barbazetto I, Burdan A, Bressler NM, *et al.* Photodynamic therapy of subfoveal choroidal neovascularization with verteporfin: fluorescein angiographic guidelines for evaluation and treatment – TAP and VIP report No.2. *Arch Ophthalmol* 2003;**121**:1253–68.
- 7 Leung DW, Cachianes G, Kuang WJ, *et al.* Vascular endothelial growth factor is a secreted angiogenic mitogen. *Science* 1989;**246**:1306–9.
- 8 Kim KJ, Li B, Winer J, *et al.* Inhibition of vascular endothelial growth factor-induced angiogenesis suppresses tumour growth *in vivo*. *Nature* 1993;**362**:841–4.
- 9 Frank RN, Amin RH, Elliott D, *et al.* Basic fibroblast growth factor and vascular endothelial growth factor are present in epiretinal and choroidal neovascular membranes. *Am J Ophthalmol* 1996;**122**:393–403.
- 10 Oh H, Takagi H, Takagi C, *et al.* The potential angiogenic role of macrophages in the formation of choroidal neovascular membranes. *Invest Ophthalmol Vis Sci* 1999;**40**:1891–8.

'Ab interno' intravitreal suturing of a large traumatic scleral perforation at the posterior pole

Closure of scleral perforations after trauma is frequently achieved by external suturing. Enlarged perforations towards the equator may require temporary removal of the external ocular muscle to visualise and fix the scleral wound.¹ Nakashizuka *et al.* attached 'ab interno' a dislocated intraocular lens by introducing a 10–0 polypropylene loop, held by an intraocular forceps, into the vitreous cavity through a sclerotomy.² Here, we describe the 'ab interno' intravitreal suturing of a large traumatic posterior scleral perforation at the posterior pole.

A 34-year-old patient presented with a paracentral corneal perforation, collapse of the anterior chamber and severe vitreous haemorrhage OD. The patient's visual acuity was light perception. The patient was treated in a two-step fashion according to Kuhn *et al.*³ A primary corneal wound closure and placement of a mild scleral buckle was immediately performed, with the application of topical and systemic corticosteroids. The following day, a computed tomography scan was ordered, which showed a 25-mm-length splinter at the posterior pole (fig 1). On day four, we performed a comprehensive reconstruction under general anaesthesia, including a lensectomy and removal of the vitreous haemorrhage by complete three-port vitrectomy. When the pars plana vitrectomy removed the crystalline lens and vitreous haemorrhage, a large metal foreign body, sticking in the posterior bulbus with a consecutive incarceration of the adjacent retina, became evident. After the splinter was removed from the sclera, a

Figure 1 Orbital computed tomography scan after the primary corneal wound closure and placing an encircling buckle: A splinter of 25 mm length is seen at the posterior pole sticking through the sclera. The location of the cerclage is seen as a dark black area perpendicular to the vitreous base.

hypotonic globe with a Y-shaped scleral rupture at the posterior pole remained. The size of the wound, 3 mm by 4 mm superotemporal to the optic nerve at the posterior pole, prevented suturing from outside or sufficient silicone oil endotamponade.

To avoid an enucleation of the globe, the scleral perforation had to be closed so that 5000 cst silicone oil would remain in the vitreous cavity. We decided to close the large scleral perforation from inside. After lowering the infusion pressure, incarcerated choroidal and retinal tissue was removed from the laceration site with the vitrectomy. A light pipe was introduced via a fourth sclerotomy and fixed with tape,⁴ thus freeing the surgeon's second hand for bimanual intraocular vitreo-retinal manoeuvres. A 10/0 nylon suture with an attached needle was introduced with an intraocular forceps via a sclerotomy into the vitreous cavity. A second intraocular forceps was used to hold scleral tissue, while the needle was passed through the rigid sclera from inside. The knots sealed the scleral wound and a waterproof closure was achieved, completing the vitrectomy. The posterior hyaloid was now engaged at the optic disc and progressively removed up to the vitreous base.

Remaining retina reattached after drainage of the subretinal liquid, endotamponade with silicone oil and endolaser at the edges of the

Figure 2 3 months after the initial trauma: The cornea presents a crescent-shaped wound in the infero-temporal quadrant.

Figure 3 Postoperative finding 6 months after the initial trauma and following silicone oil removal. The globe had a normal shape, normal intraocular pressure and an attached peripheral retina.

retinectomy. The vitreous cutter evacuated subretinal haemorrhages as well as all incarcerated tissues and debris in a ring of 2 mm around the exit wound. This prophylactic 360-degree "chorioretinectomy", combined with three rows of deep endolaser retinopexy around the exit, may reduce the incidence of proliferative vitreoretinopathy significantly.³

Three months after the initial surgery, the cornea section showed an unremarkable peripheral scar (fig 2), the retina remained attached and the intraocular pressure (IOP) was 14 mm Hg, so that silicone oil was removed (fig 3). Visual acuity was limited to hand movements due to a large retinal defect at the posterior pole; however, the patient appreciated the regained peripheral visual field for orientation during 15 months' follow-up.

Comment

Eyes with double perforation are encountered through injuries. Individuals frequently become blind due to retinal detachment, hypotony, endophthalmitis or phthisis. The surgical goal is to maintain limited function and integrity of the globe. Although ocular hypotony secondary to scleral rupture of a choroidal coloboma is treatable by scleral silicone buckling,⁵ we sutured the sclera 'ab interno'.

It is known that the primary wound closure should be performed as early as possible. However, in complex ocular trauma, we prefer a staged approach with an immediate globe closure and a delayed comprehensive globe reconstruction. A delay may diminish the threat of intraoperative haemorrhage and increase the chance of a spontaneous posterior vitreous detachment (*Relja Zivojnovic*, personal communication).

'Ab interno' suturing is feasible for achieving a waterproof closure of large scleral perforation at the posterior pole, leaving an eye with good cosmetic appearance, normal IOP and maintained peripheral visual function.

Acknowledgements

This case was presented in parts at the annual Vitreous Society Meeting in New York, NY, USA, and achieved the First prize of the Video award 2004. This video may be seen at the BJO online video collection at <http://bjophthalmol.com/supplemental>.

Jörg C Schmidt, Stefan Mennel, Steffen Hörle, Carsten H Meyer

Department of Ophthalmology, Philipps-University Marburg, Germany

Correspondence to: Prof Dr Carsten H Meyer, Dept. of Ophthalmology, University of Bonn, Ernst-Abbe-Strasse 2, 53127 Bonn, Germany; meyer_eye@yahoo.com

doi: 10.1136/bjo.2006.112201

Accepted 1 March 2007

This work has been supported by the Fehr Foundation, Marburg, Germany.

Competing interests: None declared

The supplemental video is available at <http://bjophthalmol.com/supplemental>

References

- 1 **Carpinetto P**, Ciancaglini M, Scaramucci S, *et al*. Management of scleral rupture during retinal detachment surgery: a case report. *Eur J Ophthalmol* 2002;12:553-5.
- 2 **Nakashizuka H**, Shimada H, Iwasaki Y, *et al*. Pars plana suture fixation for intraocular lenses dislocated into the vitreous cavity using a closed-eye cow-hitch technique. *J Cataract Refract Surg* 2004;30:302-6.
- 3 **Kuhn F**, Mester V, Morris R. A proactive treatment approach for eyes with perforating injuries. *Klin Monatsbl Augenheilkd* 2004;221:622-8.
- 4 **Schmidt JC**, Rodrigues EB, Meyer CH. A simple technique to fixate the bullet pipe and perform bimanual vitreous surgery. *Ind J Ophthalmol* 2004;52:337-8.
- 5 **Viola F**, Morescalchi F, Gandolfo E, *et al*. Ocular hypotony secondary to spontaneously ruptured sclera in choroidal coloboma. *Arch Ophthalmol* 2004;122:1549-51.

Photodynamic therapy in non-subfoveal choroidal neovascularization secondary to pathological myopia: 1-year outcome

This prospective, open-label, non-comparative, interventional case series evaluates the results of photodynamic therapy (PDT) in juxtafoveal and extrafoveal choroidal neovascularization (CNV) secondary to pathologic myopia in 27 eyes of 27 patients.

The average visual acuity dropped from 58.5 letters to 49.19. The visual acuity remained stable in 13 patients (48.1%), whereas 14 patients (51.9%) lost more than 8 letters due to the extension of the lesion to subfoveal space. A strong association was found between advanced age and a worse visual prognosis ($p = 0.001$).

PDT may stabilise visual acuity in young patients with non-subfoveal CNV secondary to pathologic myopia.

The long-term visual prognosis of myopic CNV is poor;¹ however, favourable prognostic factors such as young age, good initial visual acuity and the extrafoveal location of the lesion have been described.²

The present study aims to analyse the results of PDT in juxtafoveal and extrafoveal CNV secondary to pathologic myopia.

Case report

The study included 27 eyes of 27 patients (21 women and 6 men), with non-subfoveal CNV secondary to pathologic myopia, with a follow-up period of 1 year.

The mean best corrected visual acuity (BCVA; see table 1) dropped nine letters.