- 33 **Cooke HJ**, Reddix RA. Neural regulation of intestinal electrolyte transport. In: Johnson LR, ed. *Physiology of the gastrointestinal tract*. New York: Raven, 1994:2083–132.
- 34 **Sidhu M**, Cooke HJ. Role for 5-HT and ACh in submucosal reflexes mediating colonic secretion. *Am J Physiol Gastrointest Liver Physiol* 1995;**269**:G346–51.
- 35 **Rouzade-Dominguez ML**, Pernar L, Beck S, et al. Convergent responses of Barrington's nucleus neurons to pelvic visceral stimuli in the rat: a juxtacellular labelling study. *Eur J Neurosci* 2003;**18**:3325–34.
- 36 **Valentino RJ**, Kosboth M, Colflesh M, et al. Transneuronal labeling from the rat distal colon: anatomic evidence for regulation of distal colon function by a pontine corticotropin-releasing factor system. *J Comp Neurol* 2000;**417**:399–414.
- 37 **Miampamba M**, Maillot C, Million M, et al. Peripheral CRF activates myenteric neurons by the composition of the commensal flora in the rat. *Am J Physiol Gastrointest Liver Physiol* 2002;**282**:G857–65.
- 38 **Chatzaki E**, Crowe PD, Wang L, et al. CRF receptor type 1 and 2 expression and anatomical distribution in the rat colon. *J Neurochem* 2004;**90**:309–16.
- 39 **Garcia-Lafuente A**, Antolin M, Guarner F, et al. Modulation of colonic barrier function by the composition of the commensal flora in the rat. *Gut* 2001;**48**:503–7.
- 40 **Soderholm JD**, Yang PC, Ceponis P, et al. Chronic stress induces mast cell-dependent bacterial adherence and initiates mucosal inflammation in rat intestine. *Gastroenterology* 2002;**123**:1099–108.
- 41 **Garcia-Rodenas CL**, Bergonzelli GE, Nutten S, et al. Nutritional approach to restore impaired intestinal barrier function and growth after neonatal stress in rats. *J Pediatr Gastroenterol Nutr* 2006;**43**:16–24.
- 42 **Anisman H**, Zaharia MD, Meaney MJ, et al. Do early-life events permanently alter behavioral and hormonal responses to stressors? *Int J Dev Neurosci* 1998;**16**:149–64.
- 43 **Pihoker C**, Owens MJ, Kuhn CM, et al. Maternal separation in neonatal rats elicits activation of the hypothalamic-pituitary-adrenocortical axis: a putative role for corticotropin-releasing factor. *Psychoneuroendocrinology* 1993;**18**:485–93.
- 44 **Shanks N**, Windle RJ, Perks PA, et al. Early-life exposure to endotoxin alters hypothalamic-pituitary-adrenal function and predisposition to inflammation. *Proc Natl Acad Sci USA* 2000;**97**:5645–50.
- 45 **Dinan TG**, Quigley EM, Ahmed SM, et al. Hypothalamic-pituitary-gut axis dysregulation in irritable bowel syndrome: plasma cytokines as a potential biomarker? *Gastroenterology* 2006;**130**:304–11.
- 46 **Kawahito Y**, Sano H, Kawata M, et al. Local secretion of corticotropin-releasing hormone by enterochromaffin cells in human colon. *Gastroenterology* 1994;**106**:859–65.
- 47 **Nilsson O**, Cassuto J, Larsson PA, et al. 5-Hydroxytryptamine and cholera secretion: a histochemical and physiological study in cats. *Gut* 1983;**24**:542–8.
- 48 **Beubler E**, Horina G. 5-HT₂ and 5-HT₃ receptor subtypes mediate cholera toxin-induced intestinal fluid secretion in the rat. *Gastroenterology* 1990;**99**:83–9.
- 49 **Tsukamoto K**, Nakade Y, Mantyh C, et al. Peripherally administered CRF stimulates colonic motility via central CRF receptors and vagal pathways in conscious rats. *Am J Physiol Regul Integr Comp Physiol* 2006;**290**:R1537–41.

EDITOR'S QUIZ: GI SNAPSHOT

Upper gastrointestinal bleeding with a difficult endoscopy

Robin Spiller, editor

Clinical presentation

An 84-year-old man presented at our institution with tarry stools and epigastric pain for 1 day without nausea or vomiting. A medical history included a percutaneous coronary angioplasty (2 years previous) and chronic obstructive airways disease. On physical examination, the patient demonstrated mild tenderness in the epigastrium; however, no mass was palpable. Laboratory data revealed mild normocytic and normochronic anaemia (Hb 11.3 g/dl and Hct 32.9%). A plain abdominal x ray showed a normal gas pattern. An upper gastrointestinal endoscopy revealed a giant fold invaginated into the pyloric orifice originating from halfway down the lesser curvature of the stomach. This prevented the advancement of the endoscope as well as visualisation of the antrum (fig 1). A barium meal examination revealed a dilated stomach and an intramural mass lesion over the bulb of the duodenum (fig 2).

Question

What is the diagnosis?

See page 1556 for answer

Y-Y Chen, H-H Yen, M-S Soon

Department of Gastroenterology, Changhua Christian Medical Center, Changhua, Taiwan

Y-Y Chen

Chungtai Institute of Health Sciences and Technology, Taichung, Taiwan

Correspondence to: Yang-Yuan Chen, Changhua Christian Medical Center, 135 Nanshao Street Changhua, 500 Taiwan; 27716@cch.org.tw

doi: 10.1136/gut.2006.106559

Figure 1 Endoscopy revealed a giant fold invaginated into the pyloric orifice from the lesser curvature side of the middle body.

Figure 2 A barium meal examination revealed a dilated stomach and an intramural mass lesion over the bulb of the duodenum (arrow).