but no substitute for data. In our view, only those patients should undergo interventions of their renal arteries that are included in a randomised, controlled, trial of ARAS such as CORAL. Dear and colleagues have done us a service in drawing attention to "Guideline Mayhem".¹

Authors' affiliations

F C Luft, C M Gross, Medical Faculty of the Charité, Franz-Volhard Clinic, HELIOS Klinikum Berlin-Brandenburg, Berlin, Germany

Competing interests: None declared.

Correspondence to: Dr F C Luft, Experimental and Clinical Research Centre, Lindenberger Weg 80, 13125 Berlin, Germany; Luft@charite.de

Accepted 10 July 2007

REFERENCES

- 1 **Dear JW**, Padfield PL, Webb DJ. New guidelines for drive-by renal arteriography may lead to an unjustifiable increase in percutaneous intervention. *Heart* 2007;**93**:1528–30.
- 2 **Hirsch AT**, Haskal ZJ, Hertzner NR, *et al*. American Association for Vascular Surgery; Society for Vascular Surgery; Society for Cardiovascular Angiography and Interventions; Society for Vascular Medicine and Biology; Society of Interventional Radiology; ACC/AHA Task Force on Practice Guidelines Writing Committee to Develop Guidelines for the Management of Patients With Peripheral Arterial Disease; American Association of Cardiovascular and Pulmonary Rehabilitation; National Heart, Lung, and Blood Institute; Society for Vascular Nursing; TransAtlantic Inter-Society Consensus; Vascular Disease Foundation. ACC/AHA 2005 Practice Guidelines for the management of patients with peripheral arterial disease (lower extremity, renal, mesenteric, and abdominal aortic). *Circulation* 2006;**113**:e463–654.
- 3 **White CJ**, Jaff MR, Haskal ZJ, *et al*. American Heart Association Committee on Diagnostic and Interventional Cardiac Catheterization, Council on Clinical Cardiology; American Heart Association Council on Cardiovascular Radiology and Intervention; American Heart Association Council on Kidney in Cardiovascular Disease. Indications for renal arteriography at the time of coronary arteriography: a science advisory from the American Heart Association Committee on Diagnostic and Interventional Cardiac Catheterization, Council on Clinical Cardiology, and the Councils on Cardiovascular Radiology and Intervention and on Kidney in Cardiovascular Disease. *Circulation* 2006;**114**:1892–5.
- 4 **Balk E**, Raman G, Chung M, *et al*. Effectiveness of management strategies for renal artery stenosis: a systematic review. *Ann Intern Med* 2006;**145**:901–12.
- 5 **Garovic VD**, Textor SC. Renovascular hypertension and ischemic nephropathy. *Circulation* 2005;**112**:1362–74.
- 6 **White CJ**. Catheter-based therapy for atherosclerotic renal artery stenosis. *Circulation* 2006;**113**:1464–73.
- 7 **Scolari F**, Ravani P, Pola A, *et al*. Predictors of renal and patient outcomes in atheroembolic renal disease: a prospective study. *J Am Soc Nephrol* 2003;**14**:1584–90.
- 8 **Cooper CJ**, Murphy TP. Is renal artery stenting the correct treatment of renal artery stenosis? The case for renal artery stenting for treatment of renal artery stenosis. *Circulation* 2007;**115**:263–9.
- 9 **Dworkin LD**, Jamerson KA. Is renal artery stenting the correct treatment of renal artery stenosis? Case against angioplasty and stenting of atherosclerotic renal artery stenosis. *Circulation* 2007;**115**:271–6.

IMAGES IN CARDIOLOGY

doi: 10.1136/hrt.2006.106286

Inappropriate ICD shocks

A patient was implanted with a guidant single-chamber implantable cardioverter-defibrillator (ICD) for episodes of ventricular tachycardia in March 2005. In September, he consulted the outpatient clinic after a shock was delivered. The detailed circumstances, however, remained unclear. The panel shows a print-out of the episode as recorded by the ICD. While the first two beats are of sinus origin, high-frequency discharges (10 Hz) of increasing amplitude are seen before shock delivery. Note that the tracing is compatible with an external electromagnetic source misinterpreted as an episode of ventricular fibrillation. The following day the patient confessed that the physiotherapist he was visiting had asked him to test electro-acupuncture equipment consisting of two metal wands plugged to a 10 Hz–0.9 V electrical generator. The patient reported some tingling sensations before shock delivery. This device, sold by a German company, warns users about its potential side effects in patients with an ICD or pacemaker.

Denis Graf, Etienne Pruvot
denis.graf@chuv.ch