in the α -chain, and of haemoglobin S, which carries its abnormality in the β -chain into α - and β -chains. On recombination they obtained haemoglobin A as well as a hybrid composed of abnormal α - and β -chains. One might therefore expect that when haemoglobin S and another abnormal haemoglobin are found together with haemoglobin A, S being a β -chain mutant, the other abnormal haemoglobin would be an α -chain mutant. We have reason to believe, however, that haemoglobin O is not an α -chain mutant, and indeed the same applies to haemoglobin P. A full account of this will be given by two of us (B.R. and H.L.) in collaboration with Drs. E. M. Shooter and E. R. Huehns, of University College, London, and a tentative explanation for this apparent paradox will then be put forward.

Summary

A new sickling condition—sickle-cell haemoglobin O trait—has been demonstrated in two children of an Arab family. This is the first occasion on which haemoglobin O has been seen outside Indonesia and that its mode of inheritance could be examined.

REFERENCES

- Aksoy, M., Ikin, E. W., Mourant, A. E., and Lehmann, H. (1958). *Brit. med. J.*, **2**, 937.
 Dherte, P., Lehmann, H., and Vandepitte, J. (1959). *Nature (Lond.)*, **184**, 1133.
 Ingram, V. M. (1959). *Brit. med. Bull.*, **15**, 27.
 Itano, H. A., and Robinson, E. (1959). *Nature (Lond.)*, **183**, 1799.
 Lie-Injo Luan Eng and Sadono (1958). *Brit. med. J.*, **1**, 1461.
 Singer, K., Chernoff, A. I., and Singer, L. (1951). *Blood*, **6**, 413.
 Smith, E. W., and Torbert, J. V. (1958). *Bull. Johns Hopk. Hosp.*, **102**, 38.

A CASE OF SUBACUTE RICKETTSIAL ENDOCARDITIS; WITH A SURVEY OF CARDIAC PATIENTS FOR THIS INFECTION

BY

B. P. MARMION, M.D.

Public Health Laboratory Service, Leeds

F. E. HIGGINS, M.B., M.R.C.P.

Department of Medicine, General Infirmary, Leeds

J. B. BRIDGES, F.I.M.L.T.

Public Health Laboratory Service, Leeds

AND

A. T. EDWARDS, M.B., Ch.B.

Department of Pathology, Halifax General Hospital

A few reports during the last 10 years suggest that Q fever in man sometimes pursues a chronic course with infection of the heart valves. Marmion *et al.* (1953) reviewed three possible cases of the condition—two Californian patients and one English.

More recently it was possible to observe the course of the disease during a two-year period in a patient of Dr. F. H. Scadding's at the Middlesex Hospital, London. A brief description of the findings was given at the Sixth International Congress of Tropical Medicine and Malaria (see Marmion, 1958), and a detailed account has been published together with a review of the literature on persistent Q fever infections (Robson and Shimmin, 1959; Andrews and Marmion, 1959). The patient was

a man aged 47, who had pneumonia clinically resembling Q fever, recovered, and was apparently healthy for six months, then became chronically ill with a swinging fever. Aortic valve incompetence developed and gradually increased until his death some 18 months later. *Rickettsia burneti* was isolated from his bloodstream at the beginning of the chronic febrile illness and again just before death; it was also shown to be present in large numbers in aortic valve vegetations collected after death.

Evans, Powell, and Burrell (1959) have also reported the isolation of *R. burneti* from the aortic valve vegetations and from the spleen of a patient dying five months after an influenza-like illness.

Although subacute rickettsial endocarditis is probably a rare sequel to an acute infection with *R. burneti*, it seemed desirable to examine a number of cardiac patients for rickettsial antibodies so as to obtain an estimate of its frequency.

This paper describes the results of these examinations so far, together with the clinical state, history, and laboratory diagnosis of another patient with subacute Q fever endocarditis who was detected as a result.

Selection of Patients and Methods

Sera were obtained from suspected cases of subacute endocarditis in Leeds and certain other centres. The sera were tested for complement-fixing antibodies to the phase 1 and 2 antigens of *R. burneti* (see Andrews and Marmion, 1959), and also for those against murine typhus and rickettsial-pox group-specific antigens. The complement-fixation test used was either that described by Stoker, Page, and Marmion (1955) or a locally developed micro-method using essentially the same reagent concentrations and incubation times.

Results

Fifty-six patients were tested in the period between April and December, 1959. A diagnosis of subacute bacterial endocarditis, with isolation of *Streptococcus viridans*, enterococci, or other bacteria, was made in 13 of the patients. In 35 of the remainder there was evidence of valvular damage and a strong suspicion of an infective process, but blood cultures were sterile. The remaining eight patients had valvular damage but no evidence of infection.

No patient had C.F. antibody to the group-specific typhus antigen. One patient had Q fever C.F. antibodies and a case report is given below.

Another patient had a C.F. antibody titre of 1/128 to rickettsial-pox group-specific antigen (Lederle). Dr. J. Colquhoun (Department of Bacteriology, Royal Infirmary, Sheffield) kindly informed us about his investigations, which indicate that this patient's endocarditis was of bacteriological origin. No antigenic relationship was shown between the causative bacterium and the rickettsial-pox antigen, and it seems probable that the rickettsial-pox antibody resulted from some past experience. Dr. C. C. Shepard (Montgomery, Ala.) kindly confirmed the specificity of the serological findings by C.F. tests with their rickettsial-pox antigen and by staining smears of *Rickettsia akari* with 1/10 and 1/100 dilutions of the patient's serum, using the indirect fluorescent antibody technique. The patient has not been out of the country, and local investigations into the possible presence of a member of the spotted-fever

group of rickettsiae—or some other organism with a related antigen—are in progress.

Case Report

The patient with Q fever antibody was a window-cleaner aged 48, and was first admitted to hospital (St. Luke's Hospital, Bradford) on June 18, 1957. On the day before admission he had a frontal headache for about three hours, and had sweated a lot and taken plentiful fluids. During the night he shivered and slept fitfully. On rising he noticed weakness of the right arm and leg, together with pain on the left side of his neck. A slight frontal headache was still present.

His medical history revealed that in 1953 he had suffered from an iridocyclitis of the right eye accompanied by extensive retinal haemorrhages and secondary optic atrophy (diagnosed as Eales's disease). At that time the urine was free from albumin and the Kahn reaction was negative, but a chest x-ray examination had revealed some cardiac enlargement. His blood-pressure was then 140/60 mm. Hg, but no observations were made on the presence or absence of cardiac murmurs. There was no history of chorea or acute rheumatism. His general practitioner recorded attacks of right-sided pain (? pleurisy) at the age of 23 and again at the age of 41. The family history was not significant except that his elder brother had an attack of chorea at the age of 10 and was known to have had mitral stenosis and aortic incompetence without infective endocarditis at the age of 44. He died in congestive heart failure five years later.

On examination the patient was afebrile and drowsy, but was able to give a coherent history. He had a spastic right hemiparesis without sensory loss. The heart was in regular rhythm with no murmurs and was apparently of normal size. Blood-pressure was 125/65 mm. Hg. The pulsation in the left carotid artery was diminished. He appeared to be anaemic. A lumbar puncture yielded a clear fluid not under increased pressure, with normal chemical readings and with no excess of cells.

During the first few days after admission the patient's drowsiness and the right-sided weakness increased and he became incontinent of urine. A provisional diagnosis of left carotid occlusion was made.

Numerous rhonchi were found in the chest, but, apart from a few temperature recordings of 99° and 99.6° F. (37.2 and 37.6° C.), he was afebrile. A chest x-ray film taken with portable x-ray equipment gave a poor result but suggested pneumonic changes at the right base. Between June 19 and 24 he was given seven million units of penicillin. He gradually improved except for an unexplained effusion into the right knee-joint on August 17, and he was discharged home in November, 1957. By April, 1958, he was able to do light work in a bakery.

In October, 1958—some 16 months after the first admission—he had several right-sided convulsions and was readmitted to St. Luke's Hospital. He was afebrile. Finger-clubbing was present and the heart was enlarged. An aortic diastolic murmur was now audible at the left sternal edge. No mitral murmurs were detected, but radiological screening revealed a large left auricle.

Investigations revealed a normochromic anaemia and an increased sedimentation rate (57 mm./hour Wintrobe). The urine contained a trace of albumin and there were numerous red cells in the centrifuged deposit. Four blood cultures were sterile and the patient was consistently afebrile. Other negative investigations included examination of the peripheral blood and marrow preparations for L.E. cells, estimation of total proteins and albumin/globulin ratio, a muscle biopsy, and eosinophil counts.

Nevertheless the clinical evidence for infective endocarditis was strong enough to justify antibiotic therapy, and during the next six weeks he was given one million units of penicillin six-hourly.

He failed to improve, and the albuminuria, the anaemia, and the raised sedimentation rate persisted. After six weeks' treatment, prednisone, 10 mg. daily, was substituted for antibiotics, and was continued until mid-February, 1959, without objective evidence of improvement. This steroid was eventually discontinued when the patient developed oedema of the ankles.

In May, 1959, he was transferred to the General Infirmary at Leeds, under the care of Dr. D. Taverner, for further investigation. At this stage he was in congestive heart failure, with auricular fibrillation and unmistakable evidence of both mitral stenosis and aortic incompetence. Inequality of carotid pulsation was confirmed, but there was neither splenomegaly nor evidence of emboli. The urine contained albumin varying in amount from 2 to 5 g./litre. Further investigations revealed a serum albumin of 1.8 g./100 ml., globulin 3.5 g./100 ml., and serum cholesterol of 290 mg./100 ml. Although his cardiac failure responded to digitalis and diuretics his nephrotic manifestations persisted. A biopsy of right and left kidneys in June, 1959, was reported (Dr. C. K. Anderson) as showing the changes of chronic lobular glomerulonephritis (Ellis type 2)—an unusual feature, although it has been described in subacute bacterial endocarditis (Kark *et al.*, 1958; Spain and King, 1952).

During the course of six weeks in hospital the patient was afebrile save on two occasions when temperatures of 99 and 101°F. (37.2 and 38.3°C.) were recorded. Six further blood cultures were sterile, and renewed search for evidence of polyarteritis nodosa or disseminated lupus erythematosus was unrewarded.

As part of the survey of endocarditis patients a sample of serum was then examined for Q fever antibodies and found to be positive.

The patient suffered progressive renal impairment, the blood urea rising to 100 mg./100 ml. He was subsequently transferred to Scotton Banks Sanatorium, where he gradually deteriorated, dying of congestive heart failure on August 26, 1959.

Pathological Investigations

At necropsy significant internal findings were confined to the chest and abdomen. Approximately 2 pints (1,140 ml.) of serous fluid was present in each pleural cavity.

Both lungs were oedematous and the left lung was collapsed, fibrosed, and covered with a layer of fibrous tissue which partially obliterated the left pleural cavity.

The heart showed moderate hypertrophy and dilation of the right ventricle and gross hypertrophy of the left ventricle. Pulmonary and tricuspid valves were normal but the aortic and mitral valves showed thickening and fusion of the cusps and small nodules of collagen—probably healed vegetations—along their points of opposition. The mitral valve was the more obviously diseased, and in addition to incompetence there was a mild sub-valvular type of stenosis due to fibrosis and adhesion of the tips of the papillary muscles with the valve cusps. There was also a small haemorrhagic friable area at the point where the anterior mitral cusp fused with the papillary muscles; possibly a more active vegetation.

In the abdominal cavity there was gross ascites. The liver was slightly enlarged, with a typical chronic congestive ("nutmeg") appearance. No striking abnormality was found in the other abdominal organs, and, in particular, there was no sign of old embolic episodes. There were no significant changes in the brain, in the meninges, or in any other system.

The cause of death was thought to be congestive heart failure due to mitral and aortic valvular disease with healed infective endocarditis.

Histological Investigations

In the anterior mitral cusp there was evidence of a chronic lesion which had largely healed. The deeper parts of the cusp consisted of dense collagen with a very scanty cellular infiltrate of lymphocytes and plasma cells. Among this tissue there were two localized areas which in a haematoxylin- and eosin-stained section appeared to consist of compact masses of basophilic debris. In sections stained by the Giemsa method these aggregations were seen to be collections of rickettsia-like organisms (see Fig.). The microcolonies were elongated, and were composed of numerous organisms 0.5μ or less in size and staining blue-purple with the Giemsa method and a faint red with Gram's stain. As with the case described

Photomicrograph of colonies of rickettsia-like organism found in mitral valve. (Giemsa. $\times 1,000$.)

by Andrews and Marmion (1959), they were not clearly inside or associated with cells. They were closely invested by the collagen and there was little or no inflammatory response in the immediately surrounding tissue.

More superficially in the cusp there were patchy layers of loose granulation tissue in which the mononuclear cellular reaction was more intense. No colonies of bacteria could be identified in these areas, using Giemsa or Gram stains. A small number of cells with cytoplasmic granules varying in size from 0.5 to 2μ or more and staining purple with Giemsa was seen in this area. These were thought to be mast cells rather than cells with intracytoplasmic colonies of rickettsiae. The posterior mitral cusp and the aortic valve, although scarred, showed no signs of active disease and organisms were not seen.

Sections from the lower lobe of the left lung confirmed the macroscopic appearances of old collapse and fibrosis, and there was widespread intra-alveolar oedema and accumulations of "heart-failure" macrophages. No microorganisms were seen in numerous Giemsa-stained sections. The liver showed chronic venous congestion and the splenic pulp was congested with reactive hyperplasia. Sections from the brain, and mediastinal and mesenteric lymph nodes were normal. Sections of the kidney showed a chronic nephritis (Ellis type 2) and there was no evidence of focal embolic nephritis.

Bacteriological Investigations

Tests on Specimens Taken During Life.—The first positive Q fever complement-fixation finding was obtained with a serum specimen collected on May 11, 1959—that is, about three and a half months before death. There was an antibody titre of $1/256$ to the phase 1 C.F. antigen of *R. burneti* and a titre of $1/64$ to the phase 2 antigen. Essentially the same titres were obtained in tests carried out on sera collected on May 21 and June 6, 1959. Attempts were made to isolate *R. burneti* from seven samples of heparinized blood (or blood clots) and three

samples of urine taken at intervals during May and June but without success.

Tests on Specimens Taken at Necropsy.—Specimens of the lung, liver, kidney, spleen, brain, mediastinal and mesenteric lymph nodes, and the mitral and aortic valves were collected. Bacteriological culture of all these yielded a mixed growth of *Proteus* and an enterococcus—probably post-mortem contaminants. Ten per cent. (w/v) suspensions of the specimens were prepared and 2-ml. quantities were inoculated into guinea-pigs by the intraperitoneal route—two guinea-pigs each for the lung, liver, kidney, spleen, lymph nodes, and brain specimens, and four guinea-pigs each for the aortic- and mitral-valve suspensions. The rectal temperatures of the guinea-pigs inoculated with the heart-valve suspensions were taken daily, and two animals in each group which showed slight elevations of temperature between the 12th and 20th days after inoculation were killed and suspensions of their spleens passed to fresh guinea-pigs. These animals ("second-passage animals") and all those inoculated directly with the specimens ("first-passage animals"), were tested for Q fever C.F. antibody (phase 2) five to seven weeks after inoculation. The guinea-pigs inoculated with lung, liver, kidney, spleen, brain, lymph nodes, and aortic valve were negative. However, one out of two of the second-passage animals in the series inoculated with the mitral-valve suspension had Q fever (phase 2) C.F. antibody at a titre of $1/512$. The first-passage animals and the serologically negative second-passage animal in this series were then challenged with known positive material in the manner of Derrick (1937). They did not become febrile, and so appeared to be immune even though they had not developed C.F. antibody.

Consequently a fresh suspension of the mitral valve was prepared and inoculated into eight guinea-pigs and four hamsters. Four of the guinea-pigs were given cortisone acetate, 12.5 mg. subcutaneously, just before the inoculation of the specimen and the same dose 10 and 20 days after inoculation. One cortisone-treated guinea-pig died 25 days after inoculation, and a suspension of its liver and spleen was passed to fresh guinea-pigs. The surviving animals were tested for Q fever antibody four to five weeks after inoculation. The accompanying Table shows the results

Results of Serological Examination of Guinea-pigs and Hamsters Inoculated with a Suspension of the Mitral Valve

Animal	No.	Reciprocal of Titre of C.F. Antibody to the Nine Mile (phase 2) Antigen of <i>R. burneti</i>	
		Pre-inoculation Serum	Post-inoculation Serum
Guinea-pigs:			
No cortisone	132	<4	32
	133	<4	64
	134	<4	64
	135	Anti-complementary	32
Cortisone-treated	136	<4	Not tested*
	137	<4	32
	140	<4	64
	141	<4	16
Hamsters	63	Not tested	32
	64	"	64
	65	"	64
	66	"	32

* Died 25 days after inoculation: a suspension of liver and spleen produced Q fever infection in two other guinea-pigs.

obtained, and it is evident that all animals were infected with *R. burneti* and that the rickettsia had been isolated once more from the mitral valve.

Discussion

It seems from the results so far that rickettsial infection, in particular with *R. burneti*, is a rare cause of subacute infective endocarditis—at least in the areas surveyed. If both a pre-existing valvular disease and a Q fever infection are required, then obviously their coincidence in one patient will be rare indeed.

The sequence of events in our present case is not as clear as that observed in the case at the Middlesex Hospital—largely because the serological diagnosis was made only three and a half months before death and at a time when the infection was no longer active. *R. burneti* was isolated from the mitral valve—and no other organ—and rickettsia-like organisms were seen in the valve. It is of some interest that there was no typical attack of Q fever in the patient's history. The hemiplegia in 1957 seems to have resulted from a left carotid occlusion which presumably was either thrombotic or embolic in origin. If the former, then the episode may have been an atypical attack of Q fever. The headache, sweating, and ill-defined signs in the chest would fit with this, and thrombosis of peripheral vessels has been described in Q fever (for discussion and references see Robson and Shimmin, 1959). On the other hand, if the episode was embolic then the cardiac lesion may have existed before 1957, even though there was no detectable evidence of valvular disease at that time.

Despite the absence of a history of rheumatic illness the heart did appear to be enlarged on x-ray examination in 1953–4 and occasional blood-pressure readings had always revealed a wide pulse pressure. These findings, coupled with the occurrence of the fatal rheumatic heart disease without evidence of subacute infective endocarditis in the patient's brother, raise the suspicion of a silent rheumatic carditis in early life subsequently aggravated by a later and superimposed Q fever endocarditis.

After the hemiplegic attack, and during the subsequent two years of chronic illness, the picture was one of subacute infective endocarditis with the development and increase of cardiac murmurs, anaemia, and finger-clubbing, with, however, little or no fever. Repeated blood cultures in several hospitals were sterile and several courses of penicillin were ineffective.

A diagnosis of Q fever infection would not have been made but for the routine serological testing. It is suggested, therefore, that a Q fever complement-fixation test might well be done as a routine on patients with suspected subacute infective endocarditis—especially when there is a sterile blood culture, or no response to penicillin, or when treatment with corticosteroids is to be given.

It is of particular interest that the patient's serum, unlike that of an uncomplicated case of Q fever, contained antibody against the phase 1 C.F. antigen of *R. burneti*. This finding reinforces the suggestion (Andrews and Marmion, 1959) that the presence of phase 1 C.F. antibody may be a useful indicator of chronic Q fever, but obviously more cases will have to be examined before a firm conclusion can be reached.* In this connexion it may be noted that the sera from the patient described by Evans *et al.* (1959), when tested in Leeds, did not contain antibody against phase 1 C.F. antigen. This and some other differing features may perhaps be the result of the relatively short period—five months—between the presumptive attack of Q fever and death in their patient.

*Since this paper was completed Dr. N. R. Grist (Glasgow) has told us about another case of Q fever endocarditis, and we are also investigating one seen by Dr. Louis Glick (Halifax General Hospital). Both patients had high titres of antibody to Phase 1 antigen.

Summary

A small series of patients with suspected or proved subacute infective endocarditis have been examined for Q fever, murine typhus, and rickettsial-pox complement-fixing antibodies.

The results indicate that subacute rickettsial endocarditis is not a numerically important cause of the syndrome of subacute infective endocarditis with a sterile blood culture.

Fifty-six patients were examined. None had typhus C.F. antibody. One had rickettsial-pox C.F. antibody, a finding unconnected with his endocarditis. Another had Q fever antibody and was found to have a healed Q fever endocarditis.

The last patient, a middle-aged man with aortic- and mitral-valve disease, had symptoms and signs suggesting subacute infective endocarditis for over a year before his death. *R. burneti* was isolated from the mitral valve after death and rickettsia-like organisms were seen in sections of the valve.

C.F. antibody to the phase 1 antigen of *R. burneti* was again demonstrated, and a wider use of the Q fever complement-fixation test in the investigation of subacute infective endocarditis is suggested.

We are indebted to the following persons for permitting and arranging the collection of specimens from the cardiac patients. Professor C. L. Oakley and the staff of the department of bacteriology, University of Leeds; the consultant physicians of the General Infirmary at Leeds; Dr. W. Goldie and the staff of the department of pathology and the consultant physicians at St. James's Hospital, Leeds. Thanks are also due to physicians and pathologists in various parts of the country who sent sera from their patients. In particular we are grateful to Dr. D. Taverner for his help and for allowing us to publish details of the patient with Q fever who was under his care. We are also grateful to Dr. L. Smith (Bradford), Dr. R. K. MacCuish (St. Luke's Hospital, Bradford), and Dr. K. M. Barran (Scotton Banks Sanatorium) for information about the patient. Dr. C. C. Shepard (Communicable Disease Centre, Montgomery, Ala.) gave much help with advice, rickettsial-pox antigens, control sera, and confirmatory tests. Dr. H. R. Cox (Lederle), Dr. D. Lackman (Rocky Mountain Laboratory, Montana), and Dr. J. Tonge (Brisbane) also kindly provided certain antigens and control sera.

REFERENCES

- Andrews, P. S., and Marmion, B. P. (1959). *Brit. med. J.*, **2**, 983.
 Derrick, E. H. (1937). *Med. J. Aust.*, **2**, 281.
 Evans, A. D., Powell, D. E. B., and Burrell, C. D. (1959). *Lancet*, **1**, 864.
 Kark, R. M., Pirani, C. L., Pollak, V. E., Muehrcke, R. C., and Blainey, J. D. (1958). *Ann. Intern. Med.*, **49**, 751.
 Marmion, B. P. (1958). *Proceedings of 6th International Congress of Tropical Medicine and Malaria*, Lisbon, September, 1958, p. 197.
 —, Stoker, M. G. P., McCoy, J. H., Malloch, R. A., and Moore, B. (1953). *Lancet*, **1**, 503.
 Robson, A. O., and Shimmin, C. D. G. L. (1959). *Brit. med. J.*, **2**, 980.
 Spain, D. M., and King, D. W. (1952). *Ann. Intern. Med.*, **36**, 1086.
 Stoker, M. G. P., Page, Z., and Marmion, B. P. (1955). *Bull. Wld Hlth Org.*, **13**, 807.

The Reticuloendothelial Society (New England Institute for Medical Research, Ridgefield, Connecticut, U.S.A.) will shortly be publishing a symposium entitled "Reticuloendothelial Structure and Function." Investigators who live outside the United States may claim a 20% discount on the volume, provided they have an academic affiliation and the order is placed through the institution to which they belong.