cytosis, and the type of skin rash being similar in the reactions to both drugs.

The white-cell count in this case was not significantly lowered, though a transient eosinophilia was observed; this was not as pronounced as that of Makous and Vander Veer's case (22% of a total count of 22,500/c.mm.). Neutropenia has been a notable feature in reported severe reactions to phenindione, having been recorded in 10 of the 13 cases reviewed by Bingle and Shine (1959). It was present in three cases which terminated fatally (Brown and MacMillan, 1954; Kirkeby, 1954; Burns and Desmond, 1958), and there is general agreement that it constitutes an indication for treatment with steroids and antibiotics.

The skin rash was late in appearing: it may occasionally be the only manifestation of sensitivity, and in view of the possible development of agranulocytosis and hepatitis its occurrence merits withdrawal of the drug, if necessary with replacement by another anticoagulant. The management of patients exhibiting sensitivity to phenindione has been discussed by Bingle and Shine (1959), who have suggested a scheme of treatment.

Summary

A further case of hepatitis occurring as a result of sensitivity to phenindione is reported; the clinical and laboratory findings are described and previously reported cases briefly reviewed.

I thank Brigadier A. P. Trimble, consulting physician to the British Army of the Rhine, and Group Captain J. C. Bowe for their help and advice, and the laboratory staff of the Military Hospitals at Rinteln and Munster for the biochemistry.

Addendum.—Since this paper was completed another case of hepatitis occurring in a patient receiving phenindione has been reported by R. H. Brooks and H. B. Calleja (*Ann. intern. Med.*, 1960, 52, 706); it was associated with fever, a rash, and nephritis. Recovery followed withdrawal of the drug.

REFERENCES

- Bingle, J., and Shine, I. (1959). *Lancet*, 2, 377.
 Breneman, G. M., and Priest, E. M. (1955). *Amer. Heart J.*, 50, 129.
 Brown, K. W. G., and MacMillan, R. L. (1954). *Amer. J. med. Sci.*, 227, 526.
 Burns, C., and Desmond, F. B. (1958). *N.Z. med. J.*, 57, 283.
 Chesney, W. M. (1954). *Brit. med. J.*, 2, 591.
 East, E. N., and Beamish, R. E. (1957). *Canad. med. Ass. J.*, 77, 1028.
 Hanger, F. M., jun., and Gutman, A. B. (1940). *J. Amer. med. Ass.*, 115, 263.
 Kirkeby, K. (1954). *Lancet*, 2, 580.
 Makous, N., and Vander Veer, J. B. (1954). *J. Amer. med. Ass.*, 155, 739.
 Medical Research Council (1959). *Brit. med. J.*, 1, 803.
 Sherlock, S. (1955). *Diseases of the Liver and Biliary System*. Blackwell, Oxford.
 Toohey, M. (1953). *Brit. med. J.*, 1, 650.
 Werner, S. C., Hanger, F. M., and Kritzer, R. A. (1950). *Amer. J. Med.*, 8, 325.

The number of early infectious syphilis cases in America increased from 6,661 in 1958 to 8,178 in 1959, or 22.8%. Gonorrhoea rose from 220,191 in 1958 to 237,318 in 1959, or 7.8%. During 1959 venereal diseases continued to increase in the teenage population: the increase on the 1958 figure was 14.3% in the 10-14-year age group and 11.4% in the 15-19-year-olds. (*Social Health News*, American Social Health Association, February, 1960.)

IDIOSYNCRASY TO PHENINDIONE

BY

GORDON MATHER, M.A., M.D., M.R.C.P.
 Physician, Bristol Clinical Area

AND

CLIFFORD RILEY, M.D.
 Pathologist, Royal Sussex County Hospital, Brighton

Anticoagulants are being used on an increasing scale in the treatment of cardiac infarction and thrombotic and embolic disorders. The following case reports are presented so that others may be aware of some of the untoward reactions which occur.

Case Reports

Case 1.—A man aged 47 was admitted to hospital on account of sudden right-sided chest pain, dyspnoea, and haemoptysis. Four weeks previously he had had thrombosis of a right calf vein. Pulmonary embolism was diagnosed, and treatment started with phenindione. He was stabilized on a dose of 125 mg. daily, which kept the prothrombin level at about 30% of normal. After 16 days fever developed (Fig. 1) together with nausea, vomiting, and diarrhoea. Culture of urine and stools yielded no pathogens. The white-cell count was 11,400, with a normal differential. The fever and abdominal symptoms subsided when the anti-coagulant was stopped three days later, and he made an uneventful recovery.

FIG. 1.—Case 1. Temperature chart.

Case 2.—A man of 48 was admitted to hospital because of severe central chest pain, vomiting, and sweating. Cardiac infarction was diagnosed and confirmed by E.C.G. Treatment with heparin and phenindione was instituted; on a dose of 75 mg. daily the prothrombin level was about 20%. After 21 days he developed fever, diarrhoea, and an ache in the chest. Chest x-ray examination showed nothing abnormal, W.B.C. 9,800/c.mm. with a normal differential count; culture of blood, sputum, urine, and stools yielded no pathogens. He developed ulcers in the mouth which

FIG. 2.—Case 2. Temperature chart, showing relationship of fever to treatment with phenindione.

resembled pemphigus. These symptoms and the fever persisted (Fig. 2) until the phenindione was discontinued 11 days later. He made an uneventful recovery and left hospital seven weeks after admission. Ten days later anginal pain recurred, and his general practitioner gave him phenindione 50 mg. b.d. Fever and generalized aching in the muscles and eyes returned, and he was readmitted two days later. There was no clinical or E.C.G. sign of further infarction. Culture of blood, sputum, and urine again revealed no pathogens. The W.B.C. numbered 6,800/c.mm., with a normal differential count. The anticoagulant was stopped and the fever subsided rapidly. The patient made an uneventful recovery.

Case 3.—An 82-year-old woman was admitted to hospital on account of severe retrosternal pain, which was shown to be due to cardiac infarction. She was treated with phenindione, and a satisfactory control of the prothrombin level was achieved. After three weeks she developed a fever up to 101.8° F. (38.8° C.) for which no cause could be found. The white-cell count was normal. After a week of fever the anticoagulant was stopped, with immediate remission of the fever. She made an uneventful recovery.

Case 4.—A man of 67 was admitted to hospital on account of central chest pain which was shown to be due to a cardiac infarct. He was treated with phenindione, but he developed a fever on the 19th day. This persisted, and was accompanied by numerous mouth ulcers and a disproportionate tachycardia. After 14 days of fever he was seen by one of us and was thought to be suffering from a drug fever. His fever abated promptly when the phenindione was stopped. The mouth ulcers healed rapidly with the help of hydrocortisone lozenges, and he recovered.

Case 5.—A 69-year-old man who had previously been healthy suddenly became breathless and faint while walking home from a dinner. Three days later he had left-sided pleuritic pain. These symptoms persisted and he developed fever up to 102.6° F. (39.2° C.). He was admitted to hospital on October 4, 1956, a fortnight after the onset. Clinical and radiographic signs of a pulmonary infarct were found. He was treated with phenindione and penicillin; the fever, pain, and dyspnoea gradually disappeared. The penicillin was discontinued after 10 days. After three weeks in hospital he became febrile, the temperature rising to 104° F. (40° C.). On October 28 his white cells numbered only 3,000, with 8% neutrophils. Blood and urine cultures were sterile, and the Widal test was negative. On November 6 he developed a pharyngitis, and the next day his white-cell count was down to 2,800/c.mm., with 7% neutrophils. Further bacteriological examinations were negative, but he was treated with penicillin 250,000 units six-hourly. On November 9 the anticoagulant was discontinued, and the temperature quickly returned to normal. The penicillin was continued, and in addition cortisone 25 mg. daily was given because of the agranulocytosis. There was a prompt rise in the white-cell count, with healing of the mouth and pharynx. He was discharged on November 29 quite fit, and subsequent blood examinations were normal.

Case 6.—A man aged 55 developed substernal pain which was shown by E.C.G. to be due to a cardiac infarct. He was treated at home with phenindione and stabilized on a dose of 100 mg. daily. Twenty-three days after the start of the anticoagulant he developed ulcers in the mouth, a generalized urticarial eruption, and a temperature of 101° F. (38.3° C.). The white-cell count was normal. All the symptoms subsided within two days of stopping the phenindione, and he made an uneventful recovery.

Comment

Febrile reactions may occur occasionally to most drugs, and it is therefore not surprising to find such reactions to phenindione. Febrile reactions to phenindione have so far been reported in 13 patients (Bingle and Shine, 1959). We are drawing attention to this

idiosyncrasy as early recognition may prompt the physician to withdraw the drug and prevent more serious complications. It is possible that neutropenia in Case 5 could have been avoided if the fever had been attributed to a drug reaction at an earlier date. Only one fatal case of agranulocytosis due to phenindione treatment has so far been reported (MacMillan and Brown, 1953). Thus both febrile and neutropenic reactions must be an extreme rarity considering the large scale on which the drug is used. Our experience with these reactions has not altered our opinion that phenindione is in general a safe and reliable anticoagulant.

Summary

Six examples are reported of drug fever occurring during phenindione ("dindevan") treatment. In one of them profound neutropenia developed, but the patient recovered.

We thank Dr. R. Kemball Price and Dr. C. Barrington Prowse for permission to refer to patients under their care.

REFERENCES

- Bingle, J., and Shine, I. (1959). *Lancet*, 2, 377.
MacMillan, R. L., and Brown, K. W. G. (1953). *Canad. med. Ass. J.*, 69, 330.

DANGEROUS POTENTIATION OF PETHIDINE BY IPRONIAZID, AND ITS TREATMENT

BY

J. CHARLES SHEE, M.D., M.R.C.P., D.T.M.&H.

Honorary Consultant Physician, Bulawayo General
Hospital, Bulawayo, S. Rhodesia

Iproniazid, a close chemical relation of isoniazid, was first introduced for its antituberculous action. It was noted to have a euphoriant effect, and for this reason has been used in psychiatry, in the treatment of depressive states. This antidepressive effect is possibly due to its being an inhibitor of mono-amine oxidase. It has recently become popular in cardiology, for use in cases of intractable angina pectoris (Cossio, 1958; Towers and Wood, 1958).

Mitchell (1955) reported the case of a patient, receiving treatment for tuberculosis with iproniazid, who developed nausea and muscle twitching 20 minutes after being given 100 mg. of pethidine. He became cyanosed, was sweating, and had a rapid pulse, exaggerated tendon reflexes, ankle clonus, and a bilateral extensor plantar reflex. The attack subsided within 36 hours. Papp and Benaim (1958) described the case of a woman receiving iproniazid, 50 mg. t.d.s., for decubital angina. She had a severe attack of pain, not relieved by the injection of pethidine, 100 mg.; this was repeated after half an hour, and within minutes the woman had passed into a state of semiconsciousness with uncontrollable hyperactivity, and became completely unmanageable. Paraldehyde, 5 ml., various barbiturates, and promethazine, 50 mg. by injection, were given without effect. Finally, chlorpromazine, 50 mg. by injection two hours after the attack started, quieted her. The iproniazid was stopped. The next day, when she had a further bout of angina, suppressed by 200 mg. of pethidine, a similar state recurred, easily controlled by an injection of 50 mg. of chlorpromazine.