ance. I thank the many patients who have been willing and ready to co-operate with us on every occasion on which they have been asked. I am also indebted to the President of the Royal College of Physicians of London for my election to the Bradshaw Lectureship.

REFERENCES

- Banting, W. (1864). *Letter on Corpulence*, 3rd ed. Harrison, London.
- Bernhard, K., and Schoenheimer, R. (1940). *J. biol. Chem.*, **133**, 737.
- Brody, T. M. (1955). *Pharmacol. Rev.*, **7**, 335.
- Buchanan, J. M., and Hastings, A. B. (1946). *Physiol. Rev.*, **26**, 120.
- Chalmers, T. M., Kekwick, A., Pawan, G. L. S., and Smith, I. (1958). *Lancet*, **1**, 866.
- Pawan, G. L. S., and Kekwick, A. (1960a). *Amer. J. clin. Nutr.* In press.
- (1960b). *J. Endocr.*, **20**, 21.
- Cohn, C., and Joseph, D. (1959). *Amer. J. Physiol.*, **196**, 965.
- Dole, V. P. (1960). *Amer. J. clin. Nutr.* In press.
- Fredrickson, D. S., and Gordon, R. S., jun. (1958). *J. clin. Invest.*, **37**, 1504.
- Gersh, I., and Still, M. A. (1945). *J. exp. Med.*, **81**, 219.
- Hausberger, F. X., and Milstein, S. W. (1955). *J. biol. Chem.*, **214**, 483.
- Henriques, V., and Hansen, C. (1901). *Skand. Arch. Physiol.*, **11**, 151.
- Hollifield, G., and Parson, W. (1958). *Metabolism*, **7**, 179.
- Kekwick, A., and Pawan, G. L. S. (1956). *Lancet*, **2**, 155.
- (1957). *Metabolism*, **6**, 447.
- and Chalmers, T. M. (1959). *Lancet*, **2**, 1157.
- Keys, A., and Brozek, J. (1953). *Physiol. Rev.*, **33**, 245.
- Lancet*, 1959, **2**, 276.
- Mackay, E. M., and Sherrill, J. W. (1941). *Endocrinology*, **28**, 518.
- MacLagan, N. F., and Sheahan, M. M. (1950). *J. Endocr.*, **6**, 456.
- Mayer, J. (1957). *Metabolism*, **6**, 435.
- Mirski, A. (1942). *Biochem. J.*, **36**, 232.
- Newburgh, L. H. (1942). *Arch. intern. Med.*, **70**, 1033.
- Plummer, W. A. (1940). *Trans. Amer. Ass. Goiter*, p. 88.
- Portman, O. W., and Hegsted, D. M. (1957). *Ann. Rev. Biochem.*, **26**, 307.
- Putten, L. M. van, Bekkum, D. W., and Querido, A. (1955). *Metabolism*, **4**, 68.
- Reynolds, E. L., and Asakawa, T. (1950). *Amer. J. phys. Anthropol.*, **8**, 343.
- Rittenberg, D., and Schoenheimer, R. (1935). *J. biol. Chem.*, **111**, 169.
- Salcedo, J., and Stetten, D. (1943). *Ibid.*, **151**, 413.
- Scow, R. O. (1951). *Endocrinology*, **49**, 522.
- (1956). *Metabolism*, **5**, 79.
- Shapiro, B., and Wertheimer, E. (1943). *Biochem. J.*, **37**, 102.
- Stetten, D., jun., and Grail, G. F. (1943). *J. biol. Chem.*, **148**, 509.
- Strang, J. M. (1959). In *Diseases of Metabolism*, edited by G. G. Duncan, 4th ed., p. 529. Saunders, Philadelphia.
- Tuerkischer, E., and Wertheimer, E. (1942). *J. Physiol. (Lond.)*, **100**, 385.
- Wells, R. G. (1940). *J. Amer. med. Ass.*, **114**, 2177, 2284.
- Wertheimer, E. (1945). *J. Physiol. (Lond.)*, **103**, 359.
- and Shapiro, B. (1948). *Physiol. Rev.*, **28**, 451.
- Widdowson, E. M., McCance, R. A., and Spray, C. M. (1951). *Clin. Sci.*, **16**, 113.
- Wilmer, H. A. (1940). *Proc. Soc. exp. Biol. (N.Y.)*, **43**, 386.

"There is prevalent amongst all Africans the injection craving." This perhaps originated from the fact that at the turn of the century there were outbreaks of diseases that could be cured or prevented by making scarification marks or by giving prophylaxis by injection and also curative measures by injections, e.g., yaws and N.A.B. or bismuth. This form of curative approach came near to the African's way of effecting cure for all sorts of diseases, which was by making scarification marks and breaking the surface of the skin and rubbing in medicine. The belief was that by so doing the medicine entered the blood and through this reached all the system. Thus an injection inserted into the buttock or into a vein carries a notion as near to the old belief as possible." (Dr. W. Ouko: 3rd Conference of East African Branch of the Society of Medical Officers of Health, *E. Afr. med. J.*, March, 1960.)

WHY THIOPENTONE INJECTED INTO AN ARTERY MAY CAUSE GANGRENE*

BY

J. H. BURN, M.D., F.R.S.

Oxford

In 1948 the Swedish worker Schmitterlöw showed that if he made an extract of arteries or of veins from a horse or an ox the extract contained noradrenaline. His observations, however, did not lead him to suggest that the presence of noradrenaline in the artery wall had any importance, and no further attention was paid to it until 1957, when Burn and Rand showed that the day after rabbits were injected with reserpine the aorta no longer contained appreciable amounts of noradrenaline. Whereas the thoracic aorta from normal rabbits contained a mean quantity of 0.52 $\mu\text{g./g.}$, the aorta from rabbits treated with reserpine contained less than 0.08 $\mu\text{g./g.}$ It was also shown that while a spiral strip of aorta from a normal rabbit placed in an isolated organ bath contracted on addition of nicotine to the bath, a similar strip from a rabbit treated with reserpine did not contract. Burn and Rand (1958a) suggested that the constrictor action of nicotine was due to the release of noradrenaline from the store in the artery wall.

Another substance which was found to be inactive in the animal treated with reserpine was tyramine. Carlsson, Rosengren, Bertler, and Nilsson (1957) showed that while tyramine caused a rise of blood-pressure in an anaesthetized cat it failed to cause a rise of blood-pressure in a similar cat previously treated with reserpine. This observation was confirmed (Burn and Rand, 1958b), and several other amines related to tyramine, among them amphetamine and ephedrine, were shown to behave in the same way. While they were able to raise the blood-pressure in a normal animal, they were unable to do so in an animal which had been previously treated with reserpine.

These observations showed that substances which cause constriction may do so in two ways. They may constrict the blood-vessels themselves like adrenaline and noradrenaline, or they may act indirectly by liberating noradrenaline from the store in the artery wall.

Action of Thiopentone

There has been a series of recorded accidents in which an anaesthetist has injected thiopentone into an artery by mistake, probably because of the aberrant course of the artery. The incidence has been variously estimated from 1 in 8,000 (Lundy, 1942) to 1 in 55,000 (Cohen, 1948). Out of 24 cases, nine developed gangrene, and, of these, six required amputation of the forearm and three lost one or more digits. Professor Sir Robert Macintosh pointed these accidents out to me and asked whether the action of thiopentone could be investigated. It seemed possible that thiopentone might prove to be a substance which released noradrenaline from the store in the artery wall, and experiments were carried out to see if this was so. Observations were first made on spiral strips of rabbit aorta. When 2 mg. of thiopentone was added to a 5-ml. bath the strips of aorta contracted; they relaxed again when the solution in the bath was changed. Observations were then made by perfusing

*A lecture to the North of England Society of Anaesthetists, May 13, 1960.

the isolated ear of the rabbit with a modified Ringer's solution. A cannula was tied in the central artery, and the solution entering the cannula then passed through the ear vessels and a record of the outflow was obtained. It was found that the injection of 25–50 μ g. of thiopentone into the solution before it entered the artery caused a constriction of the vessels of the ear similar to that caused by small doses of noradrenaline. Our first thought was that the vasoconstriction might be due to the reaction of the solution; the solution of thiopentone which we injected had a pH of 8.65. When, however, we injected the same volume of Ringer-Locke's solution adjusted to the same pH there was no change in outflow.

In order to see if the vasoconstriction caused by thiopentone was due to the release of noradrenaline we applied three tests. The first was to repeat the injections of noradrenaline and of thiopentone after adding cocaine to the perfusion fluid. Cocaine is known to potentiate the vasoconstrictor action of noradrenaline. We found that cocaine also potentiated the effect of thiopentone and to the same extent. The second was to determine the effect of tolazoline ("prisol"). Tolazoline is known to abolish the constrictor action of noradrenaline, and we found that it abolished the constrictor action of thiopentone also.

Finally, we tested the action of thiopentone on the vessels of the ears of rabbits which had been treated with reserpine. The injection of reserpine removed the extractable noradrenaline from the ears, and we found that thiopentone had little or no constrictor action in them. All three tests gave results consistent with the idea that thiopentone caused constriction by releasing noradrenaline from the artery walls (Burn and Hobbs, 1959).

A substance with similar properties to thiopentone is hexobarbitone. There are, however, no records of gangrene after the injection of hexobarbitone into patients. We therefore tested its action on the vessels of the perfused rabbit ear. We found that it had almost no constrictor action—less than one-sixtieth of that of thiopentone. This finding supported the idea that the constrictor action in the rabbit ear was related to the production of vascular thrombosis and gangrene.

Effect of Sympathectomy

Some months later observations were described by Kinmonth and Shepherd (1959) in which they injected thiopentone into the central artery of the rabbit's ear while the animal was anaesthetized. They found that in the course of the next two weeks the terminal portion of the ear became gangrenous. They standardized their procedure, and, having injected a group of 20 rabbits, found that the mean area of gangrene produced was 14.5 sq.cm. Among other experiments, they removed the superior cervical ganglion and divided the great and posterior auricular nerves at the base of the ear. Having allowed time for the sympathetic fibres to degenerate, they injected thiopentone as before, and found that the area of gangrene produced was much less. The mean area in 20 rabbits was 8.8 sq. cm. Now when the sympathetic nerves to an organ have degenerated, Euler and Purkhold (1951) found that the noradrenaline which could be extracted from the organ was very greatly reduced. These results of Kinmonth and Shepherd's were therefore a further indication that the occurrence of gangrene was related to the presence of noradrenaline in the walls of the arteries, and was due to the release of this noradrenaline by the thiopentone.

B

Production of Gangrene in Tail of Mouse

The next step which was indicated was to study the occurrence of gangrene in animals treated with reserpine. I intended to do this in rabbits, but my colleague Dr. David McDougal suggested that we should try white mice. We found that if we injected a small volume, 0.04 ml. of a 10% solution of thiopentone under the skin of the tail, then signs of gangrene followed. We found it necessary to inject into the ventral surface of the tail, for injections on the lateral or dorsal aspects did not cause gangrene. It seemed likely that the injection had to be on the ventral side in order that the thiopentone should make contact with the artery.

Twenty-four hours after thiopentone injection into the tail of a mouse, examination of the tail showed necrosis at the site of injection, and the necrotic tissue often sloughed off, leaving an ulcer. Distal to the injection site a large part of the tail was cyanosed in almost all the mice. The region of cyanosis beyond the site of injection often made it difficult to distinguish the presence of a necrotic region. However, the ulcer following necrosis was easily seen and was recorded. At the end of 72 hours the cyanosed part of the tail showed signs of becoming stiff, while the ulcer remained as on the previous day. At the end of 96 hours the cyanosed part of the tail was gangrenous, being like a piece of wire.

Almost all control mice developed a gangrenous tail at the end of 96 hours. About 70% of the control mice developed an ulcer at the site of injection. In a small number of mice in which cyanosis was slight and in which there was no ulceration, the tail appeared almost normal at 96 hours.

Gangrene in Tail

The actual numbers and percentages of mice are recorded in Table I, in which the experiments have been divided into two groups according to whether reserpine

TABLE I.—*Proportion of Mice in Which Tails were Gangrenous 96 Hours After Injection of Thiopentone*

No. of Doses of Reserpine before Thiopentone. Each Dose = 10 μ g.	Control Mice	Mice Treated with Reserpine Previously
1	55/61 = 90%	38/73 = 52%
2	56/68 = 85%	27/93 = 29%

was given once (24 hr.) or twice (24 and 6 hr.) before the injection of thiopentone. At 48 and 96 hours after thiopentone injection the differences between controls and reserpine-treated mice was significant (P less than 0.01) in both groups. Further, the difference between the percentage of gangrenous tails at 96 hours after one injection of reserpine and that after two injections of reserpine was also significant at the 1% level. The effect of reserpine was to delay the appearance of gangrene by about 24 hours in animals given two doses of reserpine, and by a somewhat shorter time in those given one dose. However, all tails that became cyanosed probably became gangrenous eventually, though we did not observe them after 96 hours. The number of mice without cyanosis was not different in the control and treated groups.

Ulceration

The results for the occurrence of ulceration at the site of injection are given in Table II. The necrosis and ulceration were not so uniform in the control mice as

TABLE II.—*Proportion of Mice with Necrosis and Ulceration*

Time after Thiopentone	No. of Doses Reserpine	Control Mice	Mice Treated with Reserpine Previously
48 hours	1	24/42 = 57%	28/59 = 47%
72 "	1	32/60 = 53%	33/79 = 42%
96 "	1	30/61 = 49%	40/79 = 50%
48 "	2	49/85 = 58%	13/82 = 16%
72 "	2	57/85 = 67%	32/103 = 31%
96 "	2	54/72 = 75%	27/82 = 33%

was the occurrence of gangrene. Thus ulceration was recorded in 50–60% of the first group, whereas in the second group ulceration was recorded in 75%. One injection of reserpine before thiopentone gave no protection against ulceration, as Table II shows, but two injections gave protection, the difference between treated and control mice being significant (P less than 0.01) at 48 hours, 72 hours, and 96 hours. There was no significant difference in the incidence of necrosis or gangrene between the mice which were given no reserpine after thiopentone injection and those which were given three injections of reserpine after thiopentone injection.

These results show that the changes produced by thiopentone at the end of 96 hours affected a much smaller percentage of mice which received beforehand two injections of reserpine. So far as gangrene was concerned it seemed likely that the development was only delayed, and that it would have appeared in a higher percentage of mice at the end of 120 hours. But there was no similar progressive development of necrosis and ulceration, for the number of mice affected did not significantly increase after 72 hours. Thus two injections of reserpine had a protective action (Burn and McDougal, 1960).

Discussion

There are now three pieces of evidence concerning the production of gangrene by thiopentone. First, thiopentone has a vasoconstrictor action, which is due to the release of noradrenaline from artery walls (Burn and Hobbs, 1959). Second, partial protection against gangrene is obtained after degeneration of the sympathetic fibres (Kinmonth and Shepherd, 1959). Protection obtained in this way may be due to the disappearance of noradrenaline from the vessels when the sympathetic fibres degenerate. Third, considerable protection against gangrene is obtained by treating the animal with reserpine beforehand. This protection may also be due to the disappearance of noradrenaline from the vessels (Burn and Rand, 1957, 1958a).

That noradrenaline when liberated by thiopentone can produce gangrene is made likely by the evidence that noradrenaline infusions have led to necrosis of the skin. Kepes, Haimovici, and Simon (1954) and Shapiro and Perlow (1956) have described skin necrosis and ulceration in patients after intravenous infusion of noradrenaline (4 μ g./ml.) for periods of 48 hours. The necrosis was not around the site where the needle was introduced, but at some distance from it. Skin necrosis has also been produced in dogs and rabbits. Finally, it may be said that the liberation of noradrenaline by thiopentone is probably a local release in the vessel-wall rather than a release into the blood-stream.

Summary

If thiopentone is by accident injected into an artery, gangrene may occur in the part supplied by the artery.

Thiopentone has been shown to cause vasoconstriction in the perfused rabbit ear, and this appears to be due to the release of noradrenaline from the artery walls. The constriction is increased by cocaine, abolished by tolazoline, and is very small or absent in preparations from reserpine-treated animals.

Gangrene has been produced in the tail of the mouse by injecting thiopentone. It was accompanied by necrosis and ulceration at the site of injection. These changes were much less frequent in mice pretreated with reserpine. The administration of reserpine causes the disappearance of noradrenaline from the artery walls. Hence thiopentone appears to cause gangrene because of the local release of noradrenaline.

REFERENCES

- Burn, J. H., and Hobbs, R. (1959). *Lancet*, 1, 1112.
 — and McDougal, D. B. (1960). *J. Pharmacol. exp. Ther.* In press.
 — and Rand, M. J. (1957). *Lancet*, 2, 1097.
 — (1958a). *Brit. med. J.*, 1, 903.
 — (1958b). *J. Physiol. (Lond.)*, 144, 314.
 Carlsson, A., Rosengren, E., Bertler, A., and Nilsson, J. (1957). *Psychotropic Drugs*, edited by Garattini and V. Ghetti, p. 363. Elsevier Publishing Company, Amsterdam.
 Cohen, S. M. (1948). *Lancet*, 2, 361, 409.
 Euler, U. S. von, and Purkhold, A. (1951). *Acta physiol. scand.*, 24, 218.
 Kepes, E. R., Haimovici, H., and Simon, B. (1954). *Surgery*, 36, 822.
 Kinmonth, J. B., and Shepherd, R. C. (1959). *Brit. med. J.*, 2, 914.
 Lundy, J. S. (1942). *Clinical Anesthesia*, p. 542. Saunders, Philadelphia and London.
 Schmiterlw, C. G. (1948). *Acta physiol. scand.*, 16, Suppl. 56.
 Shapiro, R. A., and Perlow, S. (1956). *Amer. J. Surg.*, 92, 566.

A SURVEY OF THE EPILEPSIES IN GENERAL PRACTICE

A REPORT BY THE RESEARCH COMMITTEE OF THE COLLEGE OF GENERAL PRACTITIONERS*

Epilepsy has been recognized as a disease from the dawn of history. It is a common and often a dramatic malady, and yet accurate figures of its incidence are difficult to collect. Parents and relatives are only too willing to forget such happenings in their children. The admission of "blackouts" debars one from a driving licence, and may even prevent emigration. Insurance policies may be loaded, and the history of fits may make employment difficult to find. Personal factors such as these tend to make people deceptive about the illness, both in themselves and in their kinsfolk. Lord Cohen (1958) suggested that there were between 2 and 4 epileptics per 1,000 of the population; a figure borne out by Logan and Cushion (1958), who estimated the figure at 3.3 persons per 1,000. Lesser and Hunt (1954) suggest that the figure in the U.S.A. lies between 4 and 5, but there is a dearth of accurate figures; indeed, Brain (1951) quoted the findings of Gowers published some 50 years before. Tizard (1955) in a chapter on the subject gives no figures of incidence.

Another difficulty in the assessment of epilepsy is that patients in the various age-groups are usually referred to different consultants. Children are sent to the paediatrician, the aged to the geriatric specialist, and

*Members of Working Party: Dr. D. L. Crombie, Dr. K. W. Cross, Dr. J. Fry, Dr. R. J. F. H. Pinsent, Dr. C. A. H. Watts (recorder).