- Liebow, A. A. (1952). *Atlas of Tumor Pathology*, Sect. V, Fasc. 17, p. 22. Armed Forces Institute of Pathology, Washington.
- Lubarsch, O. (1888). *Virchows Arch. path. Anat.*, 111, 280.
- MacDonald, R. A. (1956). *Amer. J. Med.*, 21, 867.
- Mattingly, T. W., and Sjoerdsma, A. (1956). *Mod. Con. cardiov. Dis.*, 25, 337.
- Mohler, D. N. (1957). *J. Amer. med. Ass.*, 163, 1138.
- Rosenbaum, F. F., Santer, D. G., and Claudon, D. B. (1953). *J. Lab. clin. Med.*, 42, 941.
- Sandler, M., and Snow, P. J. D. (1958). *Lancet*, 1, 137.
- Schnecko, R. E., McIsaac, W. M., and Page, I. H. (1959). *J. Amer. Med. Ass.*, 170, 1143.
- Sjoerdsma, A., Weissbach, H., and Udenfriend, S. (1955). *J. Amer. med. Ass.*, 159, 397.
- (1956). *Amer. J. Med.*, 20, 520.
- Stanford, W. R., Davis, J. E., Gunter, J. U., and Hobart, S. G. (1958). *Sth. med. J. (Bham, Ala.)*, 51, 449.
- Thorson, A. H. (1958). *Acta med. scand.*, 161, Suppl. 334.
- Thorson, A., Björck, G., Björkman, G., and Waldenström, J. (1954). *Amer. Heart J.*, 47, 795.
- and Nordenfekt, O. (1959). *Brit. Heart J.*, 21, 243.
- Warner, R. R. P., and Southren, A. L. (1958). *Amer. J. Med.*, 24, 903.

HYPOTHERMIA AND THE HEART-LUNG MACHINE

CLINICAL APPLICATION OF THE TECHNIQUE

BY

D. N. ROSS, B.Sc., F.R.C.S.

Consultant Thoracic Surgeon, Guy's Hospital, London

The concept of combining the heart-lung machine with hypothermia is by no means new. Many workers have combined a degree of hypothermia with their conventional heart-lung circuits, usually with the object of increasing their ability to handle larger patients.

A quite distinct development has been the use of a heart-lung circuit to achieve deep ranges of hypothermia. Notable in this field have been Gollan *et al.* (1955), who first reported extracorporeal cooling to 0° C. in dogs, and more recently there has been the experimental work of Kenyon and Ludbrook (1957). Drew and Anderson's (1959) clinical work has demonstrated the safety and advantages of temperatures around 15–20° C. The use of mechanical pumps in deep hypothermia is to maintain the circulation at temperatures below 30° C. At this temperature and below, the efficiency of the heart as a pump becomes progressively

impaired until ventricular fibrillation supervenes.

The use of moderate degrees of hypothermia with the conventional heart-lung machine adds a complication to the apparatus without fully exploiting the

FIG. 1.—Diagram of the circuit used. In clinical use a coronary sinus reservoir and two suction pumps are added.

advantages of hypothermia. On the other hand, a deep or profound reduction of body temperature, while providing the theoretical ideals of a quiescent heart and dry operating conditions, is hardly applicable in every case in which open heart surgery is contemplated.

Hypothermia offers the possibility of lower flows and reduced priming volumes. In this way blood trauma can be diminished and the ever-increasing burden on the blood banks is relieved.

With the object of providing a more flexible instrument and technique, a hypothermic by-pass apparatus

FIG. 2.—Photograph of the apparatus assembled for use.

has been designed. It is equally applicable to children and full-sized adults, and provides for short or long periods of by-pass with complete arrest of the circulation if required.

Apparatus.—Basically the circuit consists of a disk oxygenator (rather smaller than the Kay-Cross variety) and a heat-exchanger made up of thin-walled stainless-steel tubes within an insulated casing (Fig. 1). The priming volume of the machine has been kept to a minimum, and is 3 pints (1.7 litres) for the average case. It is hoped to reduce this further. All blood spilt from the heart and operative field is returned to the circuit by two separate pump lines (Fig. 2). The disks normally rotate at 120 revolutions a minute, but a slower speed-setting is provided. Cooling and rewarming fluids at –4° C. and +45° C. respectively are supplied to the heat-exchanger by a conventional hypothermic apparatus.

Technique

As soon as the chest has been opened, the venae cavae and the femoral artery are cannulated simultaneously. Cooling is then started by a partial by-pass technique—it is in effect similar to the venous cooling method of hypothermia (Ross, 1954), but a proportion of the venous return is taken from the cavae, oxygenated, cooled, and returned to the femoral artery instead of to the venous system.

The initial cooling to 30° C. takes about five minutes. When the myocardium is beginning to flag and the body's metabolic requirements are down to about 50% of normal, the patient's heart is completely by-passed by tightening the vena caval snares. At this temperature it is easy to achieve what amounts to full physiological flows even in adults. A number of possibilities are now available as follows:

1. Further active cooling can be stopped, and a conventional by-pass is then carried out with the advantages of the reduced flow requirements and the added safety factor

of hypothermia should a technical fault occur—for example, a loose connexion. Under these circumstances one has at least 10 minutes of circulatory arrest available or one can deliberately arrest the circulation to achieve a dry field. This conventional by-pass at 30° C. has been used for the closure of atrial septal defects and for open pulmonary valvotomy (Fig. 3).

As soon as the septal defect is closed or the valve has been opened, rewarming is started. The cardiectomy is then closed and the circulation through the heart is re-established. Further rewarming is carried out by a partial bypass technique while the heart carries an increasing load of the circulation.

FIG. 3.—Temperature chart recorded during closure of an atrial septal defect in a 47-year-old woman.

again proceeds to 30° C., at which stage the right ventricle is opened. While the infundibular resection is carried out, active cooling is continued to the region of 17-20° C. The ventricular septal defect can then be closed, with the advantage of a relatively quiescent heart. At this temperature, flows can be completely arrested as in Drew and Anderson's technique, but more often a very low (10 ml./kg.) flow has been used. This flow avoids the embarrassment of an excessive bronchial return and avoids a pressure build-up in the great veins. It is also likely that a partial cerebral flow is better than no flow at all even at these temperatures. To provide for this low flow a slower setting of the pump is calibrated and a slow rate of rotation of the oxygenator disks (30 r.p.m.) is maintained. Increased flows and rewarming are started as soon as the defect has been closed (Fig. 4).

3. In dealing with calcific aortic stenosis, up to an hour's work on the valve is sometimes necessary. During this time coronary flow has to be maintained with awkward perfusion techniques at normal temperatures. However, with the aid of deep hypothermia adequate protection is afforded to the myocardium, and the heart can be restarted without difficulty once the circulation is re-established.

4. In the case of complex anomalies like transposition of the great vessels it has been possible to cool to a very low temperature and to arrest the circulation completely for re-deployment of the caval cannulae. Even using total

FIG. 4.—Fallot's tetralogy. The heart was open for about 60 minutes and the circulation was maintained on a low flow for 30 minutes at a temperature between 15 and 20° C.

circulatory arrest, it is reassuring to have the ability to switch on a circulation at any stage and without delay.

Results

Twenty-five cases have been operated upon by me, though the machine and technique have been used on a number of additional patients. There has been no attempt to select the cases on the basis of age, weight, or type of lesion. Four of the six deaths were in bad-risk patients—two of these cases being reoperations for Fallot's tetralogy. One death was in a high-resistance ventricular septal defect (11 units), and one was a case of transposition. The two additional deaths were in a case of Fallot's tetralogy in which the septal defect reopened and in a case of calcific aortic stenosis where death resulted from a basilar artery embolus.

Ages have varied from 2 to 47 years and weights from 11 to 81 kg.

Summary

The method described aims at tailoring the needs of the patient to the capacity of a smaller machine. Low but physiologically full flows and low priming volumes are achieved. There is the added safety factor of hypothermia allowing for periods of complete circulatory arrest in case of mishap or where a dry field is essential.

Deep hypothermia permits operations on the aortic valve without the need for coronary perfusion methods.

The full range of hypothermia is available from moderate degrees of cooling to deep hypothermia, together with a circulation of oxygenated blood available in any quantity and at any stage during the procedure.

Total circulatory arrest has been avoided where possible. Instead, a low flow has been maintained in order to prevent rises of venous pressure and the accumulation of metabolites. Under these flow conditions there has been no need to combat metabolic acidosis, and recovery in all survivors has been gratifyingly prompt.

REFERENCES

- Drew, C. E., and Anderson, I. M. (1959). *Lancet*, 1, 748.
- Gollan, F., Phillips, R., Grace, J. T., and Jones, R. M. (1955). *J. thorac. Surg.*, 30, 626.
- Kenyon, J. R., and Ludbrook, J. (1957). *Lancet*, 2, 171.
- Ross, D. N. (1954). *Ibid.*, 1, 1108.

The first car—a Morris Miniminor—has been issued under the Government's scheme to provide better transport for those war pensioners disabled in the service of their country who are eligible for Government-issued power-propelled tricycles. Cars have been available for some of the most severely disabled war pensioners since 1948, when a scheme was introduced by the former Ministry of Pensions. On March 31 last 1,802 war pensioners in the United Kingdom and Ireland had cars, but for other severely disabled war pensioners, not eligible for cars, power-propelled tricycles have been supplied, and the number of these issued in England and Wales at the same date was 1,577. Where cars are supplied the Ministry pays the insurance and road tax, and the pensioner gets an annual allowance towards maintenance. This allowance, which increases as the car gets older, averages £65 a year over the first eight years. There is also a grant towards engine replacement once during the pensioner's use of the car. If he cannot drive he may have a friend nominated to drive for him. The Ministry pays for the insurance, tax, and maintenance of Government tricycles. There is also an annual allowance in the case of power-propelled tricycles to compensate for the increases in petrol duty of 1951 and 1952.