REFERENCES

- Andrewes, F. W., and Horder, T. J. (1906). *Lancet*, 2, 708, 775, 852.
- Beattie, J. W. (1954). *Brit. med. J.*, 2, 25.
- Brainerd, H., Lennette, E. H., Meiklejohn, G., Bruyn, H. B., and Clark W. H. (1949). *J. clin. Invest.*, 28, 992.
- Cates, J. E., Christie, R. V., and Garrod, L. P. (1951). *Brit. med. J.*, 1, 653.
- Creevy, C. D., and Feeney, M. D. (1949). *Bull. Univ. Minn. Hosp.*, 20, 314.
- Finn, J. J., and Kane, L. W. (1952). *J. Urol. (Baltimore)*, 68, 933.
- Geraci, J. E., and Martin, W. J. (1954). *Circulation*, 10, 173.
- Hager, R. P., Heitzman, E. J., and Young, R. M. (1951). *Ann. intern. Med.*, 28, 510.
- Harvey, J. C., Mirick, G. S., and Schaub, I. G. (1949). *J. clin. Invest.*, 28, 987.
- Havard, C. W. H., Garod, L. P., and Waterworth, P. M. (1959). *Brit. med. J.*, 1, 688.
- Horder, T. J. (1909). *Quart. J. Med.*, 2, 289.
- Hunter, T. H. (1947). *Amer. J. Med.*, 2, 436.
- James, T. N. (1952). *A.M.A. Arch. intern. Med.*, 90, 646.
- Loewe, L., Candel, S., and Eiber, H. B. (1951). *Ann. intern. Med.*, 34, 717.
- McCoy, W. C., and Mason, J. M. (1951). *J. med. Ass. Ala.*, 21, 162.
- Mathews, F. P. (1948). *Northw. Med. (Seattle)*, 47, 581.
- Merritt, W. A. (1951). *J. Urol. (Baltimore)*, 65, 100.
- Parmley, L. F., jun., Orbison, J. A., Hughes, C. W., and Mattingly, T. W. (1954). *New Engl. J. Med.*, 250, 305.
- Rantz, L. A., and Kirby, W. M. M. (1943). *Arch. intern. Med.*, 71, 516.
- Robbins, W. C., and Tompsett, R. (1951). *Amer. J. Med.*, 10, 278.
- Sirota, J. H., Gerber, I. E., and Baehr, G. (1947). *J. Mt Sinai Hosp.*, 94, 846.
- Waisbren, B. A. (1954). *A.M.A. Arch. intern. Med.*, 94, 846.

AORTIC OCCLUSION (LERICHE'S SYNDROME) IN MITRAL STENOSIS

REPORT OF SIX CASES

BY

F. STARER, M.R.C.P.Ed., D.M.R.D.

AND

DAVID SUTTON, M.D., F.F.R., M.R.C.P.

From the Radiological Department, St. Mary's Hospital, London

Sudden occlusion of the aortic bifurcation by a "saddle" embolus is an uncommon, but well-recognized, complication of mitral stenosis. It usually presents as an acute surgical emergency (Lodin *et al.*, 1955). Peripheral emboli are a more frequent hazard to the patient with mitral stenosis and auricular fibrillation. Gradual occlusion of the abdominal aorta leading to a typical Leriche syndrome does not seem to have been described as a complication of mitral stenosis. This syndrome is characterized by bilateral intermittent claudication, wasting and weakness of the muscles in the lower limbs, and, in the male, loss of sexual potency. Mild nutritional disturbances of the skin may also be noted. Gangrene is a late and surprisingly uncommon complication. Before our previous paper (Starer and Sutton, 1958) none of the recent larger series on insidious aortic thrombosis mentioned mitral stenosis as an aetiological factor (Milanés *et al.*, 1952; Beaconsfield and Kunlin, 1953; DeBakey *et al.*, 1954; Pataro *et al.*, 1954).

In the diagnostic radiological department at St. Mary's Hospital six patients have been seen during the last few years in whom a typical Leriche syndrome developed during the course of mitral stenosis with auricular fibrillation. These cases have occurred among some 60 patients with aortic thrombosis, 32 of which

were reported in some detail in our previous communication (Starer and Sutton, 1958). However, only passing reference was made in that article to the aetiological significance of mitral stenosis, and the evidence is now presented in more detail.

Case 1

This patient had rheumatic fever in 1931, when aged 23. During 1941 she became short of breath and developed swelling of the ankles. From 1947 onwards she had several episodes of left submammary pain with cough and frothy sputum. In 1948 her right arm suddenly became white and cold. She felt pain in the arm when she used it vigorously; this pain was relieved by rest. Gradually the symptoms abated. During 1953 she was confined to bed for a few days with a mild gastro-intestinal upset. On rising she noticed coldness of both legs, gradually creeping up to the knees. This gave way to pain and later to numbness, and her toes became blue. On admission to her local hospital both femoral arteries were found to be pulsate. Her condition improved, but two weeks later the femoral pulses could no longer be felt and walking was limited to 50 yards (45.7 metres) because of bilateral claudication.

On admission to St. Mary's Hospital on December 31, 1953, signs of mitral stenosis and auricular fibrillation were observed. No pulses could be felt in the right brachial artery or at the wrist, and no pulses were palpable in the lower limbs. B.P. was 150/100 mm. Hg. No nutritional changes were present in the legs.

Aortography showed complete occlusion of the lower abdominal aorta (Fig. 1).

On February 3, 1954, valvotomy was performed. The auricle was moderately enlarged and contained a single clot. The degree of stenosis was slight, the valve easily admitting an index finger.

Comment.—From the present point of view this case is of particular interest. After undoubted peripheral emboli in both legs the femoral pulses disappeared during observation in hospital without, apparently, any further embolic episodes.

FIG. 1.—Case 1. Complete occlusion of the lower abdominal aorta opposite L3.

Case 2

In 1936, at the age of 34, this patient had an attack of rheumatic fever. Nine years later she began to complain of shortness of breath on exertion. One morning in 1945 she awoke from sleep to find both legs numb and cold. Two days later she experienced severe pain in the buttocks, thighs, and legs. The feet became cold and blue. On attempting to walk, pain in the lower limbs immediately became worse, but some pain persisted at rest. Gradually this improved until she could walk 50 yards.

Examination on July 1, 1954, showed her to have mitral stenosis and auricular fibrillation. No nutritional changes were observed in the legs, though the feet were cold to touch. No pulses were palpable in the lower limbs.

A percutaneous lumbar aortogram showed occlusion of the aorta beginning at the level of the inferior mesenteric artery; its lower limit was at the common iliac bifurcation on the right but well down in the thigh on the left.

Comment.—In this patient, also, maximal development of symptoms occurred over a period of several days, during which there was no evidence of acute aortic embolization.

Case 3

This patient, a woman aged 45, had rheumatic fever at the age of 9. She remained well until she was 31, when she had a sudden onset of pain in the right leg; this was made worse by exercise and relieved by rest. After a short time the pain disappeared. When 42 she suffered a cerebral embolus, causing temporary loss of consciousness and transient left-sided hemiparesis; she made a perfect recovery from this. Six months later she suddenly developed pain in the right foot, gradually spreading to both legs. It was aggravated by exercise, walking being limited to 100 yards (91 metres). By this time she was also complaining of increasing shortness of breath, and mitral stenosis with auricular fibrillation was diagnosed. At the age of 44 a

mitral valvotomy was performed elsewhere, with marked relief of dyspnoea.

An aortogram performed at St. Mary's Hospital showed occlusion of the aorta from just below the level of the renal arteries. The lower limits of the block were difficult to define, but both femoral arteries filled faintly.

Comment.—Here again the full development of symptoms occurred gradually after an embolus into the right leg.

Case 4

This patient began to complain of increasing shortness of breath in 1942, when she was 29. Eight years later this had become disabling and mitral stenosis was diagnosed. During the next year (1951) auricular fibrillation was first noted. In 1952, while having a bath, she experienced sudden cramp in the right leg and temporary loss of power in that limb, which also became white and cold. She gradually recovered over the next few weeks, but was left with pain in the calf on exercise. In the following year, while convalescing from lobar pneumonia, she suffered a similar episode involving the left leg. Since then she has had cramp in the left calf, both thighs, and hips on walking, her walking distance being reduced to 15 yards (13.7 metres). The cramp is eased by rest, but she also has it during the night. The legs and thighs became very thin, and she suffered from persistent backache. There was marked weakness of the legs.

On examination the presence of mitral stenosis and auricular fibrillation was confirmed. B.P. was 110/85 mm. Hg. There was marked wasting of both legs and thighs, and the feet were cold. Pulses could not be felt in the legs, but there were no nutritional disturbances.

Two aortograms were carried out. The first (in 1952) between the two embolic episodes, demonstrated obstruction of the right common iliac artery only. The second aortogram (on July 2, 1954, after the second embolus, when symptoms were fully developed) showed complete obstruction of the lower abdominal aorta and common iliac arteries,

FIG. 2.—Case 5. Serial films from aortogram showing complete occlusion of the aorta opposite L3. The lower limits of the block are at the iliac bifurcations. The upper pole of the right kidney shows evidence of old infarct.

and on the left there was probably a separate popliteal block. In February, 1954, valvotomy was performed. The auricular appendage was unusually large, but no clot was present. The valve orifice was estimated at 1.5 sq. cm., and a good split was obtained. She made a good recovery from her operation, but walking was still limited by claudication. Two years later there was no change in her condition.

Case 5

As a child this patient, a man aged 37, had three attacks of rheumatic fever. During the war he was rejected from military service because of mitral stenosis, and at that time began to suffer from occasional shortness of breath. In 1948, at the age of 32, he had two attacks of cerebral embolism, both causing transient left hemiplegia. He was admitted to St. Mary's Hospital and found to have auricular fibrillation. Recovery was rapid and complete. In 1949 he had the first of a series of embolic episodes in the lower limbs. They were associated with sudden pain, coldness, and numbness of the legs, with return to normal in about two weeks. Gradually, however, he began to complain of intermittent claudication. Eventually, by 1953, he was left with severe bilateral claudication, numbness over the coccyx, and impotence.

On admission to hospital in August, 1953, the presence of mitral stenosis and auricular fibrillation was confirmed. B.P. was 120/66 mm. Hg. A minimal left hemiparesis was present. The legs and buttocks were wasted, the feet were cold, but no nutritional changes were noted. No pulses could be felt in the lower limbs.

An aortogram showed complete occlusion of the aorta, extending from the lower border of the third lumbar vertebral body to the bifurcations of the common iliac arteries on both sides. Above the upper limit of the block the aortic lumen was narrowed up to the level of the renal arteries. There was also evidence of old infarction of the upper pole of the right kidney (Fig. 2).

In October, 1953, mitral valvotomy was performed (Mr. Lance Bromley). The auricular appendage was moderate in size, and no clot was present in the auricle. The mitral orifice was about 1 cm. in length. A good split was obtained, and the auricular appendage was amputated.

The patient made an excellent recovery. Because of the danger of involvement of the renal arteries in the thrombotic process, it was decided to graft the aorta. An operation was carried out in April, 1954 (Professor Rob). A freeze-dried arterial homograft was inserted. Unfortunately the patient died after the operation.

Post-mortem examination showed the cause of death to have been leakage from the graft above the aortic bifurcation. The remaining aorta and other great vessels showed no evidence of atheroma.

Case 6

This man, aged 49, had been perfectly well until three weeks before admission, when he experienced pain in the left calf of sudden onset, beginning during exercise. It was followed by pins and needles in both feet. During the next few days the left foot became colder than the right, and he began to complain of pain in the calves on walking. This was relieved by rest. There was no history of rheumatic fever, and heart disease had not been previously diagnosed.

On examination he was found to have auricular fibrillation and the classical signs, clinically and radiologically, of mitral stenosis. No pulses could be felt in the lower limbs. The feet were cold, but no nutritional skin changes were present. No abnormal neurological signs were elicited, and no convincing evidence of muscle wasting was noted.

Aortography showed occlusion of the abdominal aorta from the level of the third lumbar vertebra, the block extending to the common iliac bifurcations on both sides. The collateral circulation was not as abundant as is usually seen in cases of aortic occlusion, possibly because the lesion was very recent.

Discussion

It is clear from consideration of the case-histories that the patients presented in this series had in fact suffered from peripheral emboli and that the subsequent development of occlusion of the abdominal aorta was related to these. It is equally clear, however, that these were not "saddle" emboli remaining impacted at the aortic bifurcation. After the original peripheral emboli the subsequent course of events which led to aortic occlusion is not clear. Two possibilities require consideration.

The first is that an extensive thrombosis involving the aorta was built up by the accumulation of repeated small emboli, most of which were too small to produce major symptoms by themselves. The patients under review here did in fact give histories of multiple emboli. More may have occurred which were too small to give rise to immediate symptoms but which finally led to the established picture of aortic occlusion.

The second possibility is that there was progressive clotting in the vessel proximal to the peripheral embolic occlusion. This may have occurred slowly and have led eventually to the Leriche syndrome. When clot becomes impacted in a vessel the flow of blood distally and proximally ceases, and this favours thrombosis in the stagnant segments. As a collateral circulation develops the blocked vessel becomes filled in its more distal portions by retrograde flow through collaterals. Thrombosis will spread distally to the mouth of the nearest important collateral. The resulting appearance can readily be seen on arteriograms. Proximal spread of thrombosis continues for the same reason. In the case of the common iliac artery, a swift flow is encountered again at the aortic bifurcation; nevertheless, clot is able to build up along the wall of the aorta—a state of affairs than can also be observed in aortograms. Hence, clot spreads or is carried down the opposite iliac artery. Once both common iliac arteries are occluded blood-flow in the lowermost aorta becomes greatly diminished, the only effluents being relatively small vessels. Thrombosis is again favoured and continues up to the level of the renal arteries. These vessels are large and receive a large volume of blood, and flow in the aorta above that level is again considerable. It is presumably for this reason that thrombosis commonly stops at the level of the renal arteries. Some evidence for this hypothesis can be found in the literature. Baldwin and Thomas (1954) describe a case of spreading thrombosis after an embolus, and Burt *et al.* (1955) also discuss emboli as a source of spreading thrombosis.

On present evidence it is not possible to be certain which of the views discussed here is likely to be correct. It could also be argued that patients with mitral stenosis and auricular fibrillation have an increased liability to thrombosis either as a result of changes in the blood or because of their abnormal circulatory dynamics. This is certainly true of the pulmonary circulation, where these patients are liable to spontaneous thrombosis.

If any undue tendency to thrombosis does exist, it would favour the propagation of clot, though such assumption is in no way necessary for our thesis.

The course of events in these patients may be further illustrated by the following case history.

Case 7.—A spinster aged 62 was confined to bed for 14 days with an attack of gastro-enteritis. On getting up

she noticed her right leg to be numb, and, when she attempted to walk, found that she was unsteady because of weakness of both legs. She had been getting increasingly short of breath in recent months, but was not known to have suffered from heart disease. On admission she was found to have auricular fibrillation and signs of mitral stenosis. B.P. was 180/90. The femoral pulses could not be felt on either side. An aortogram showed occlusion of the left common iliac artery at its origin, the lower limit of the block lying near the origin of the external iliac artery. On the right side the external iliac artery was occluded above Poupart's ligament, the lower limit of the block lying above the origin of the profunda artery.

In this patient, as in Case 6, peripheral vascular disease was the factor which first drew attention to the underlying mitral stenosis. We believe it likely that the extent of this patient's arterial occlusion may progress and eventually involve the lower aorta. Her history is described as it represents a stage through which the other six patients (Cases 1-6) had passed before developing complete aortic occlusion.

Incidentally, it may be pointed out that it would be quite impossible to differentiate the vascular lesions in the last patient from complete aortic occlusion on clinical grounds alone. This can be done only by means of an aortogram.

In this series four out of six patients were women. This reflects the higher incidence of mitral stenosis in women. It is in sharp contrast to the heavy male preponderance in aortic thrombosis of atheromatous origin. This sex distribution also suggests that these patients fall into a separate aetiological category.

The evidence available suggests, therefore, that emboli in peripheral vessels may sometimes result in occlusion of the abdominal aorta; once this has occurred the clinical picture differs in no way from the typical Leriche syndrome.

Summary and Conclusion

Six patients are described in whom mitral stenosis, auricular fibrillation, and aortic occlusion of gradual onset were observed. There is evidence that the Leriche syndrome may be an unusual end-result of peripheral emboli in patients with mitral stenosis.

Our thanks are due to Dr. J. Litchfield, Professor Charles Rob, and Mr. Lance Bromley for their helpful co-operation, and for referring cases for examination.

REFERENCES

- Baldwin, R. B. T., and Thomas, D. F. (1954). *Brit. med. J.*, **2**, 399.
 Beaconsfield, P., and Kunlin, J. (1953). *A.M.A. Arch. Surg.*, **66**, 356.
 Burt, C. C., Learmonth, J., and Richards, R. L. (1952). *Edin. med. J.*, **59**, 65, 113.
 DeBaKey, M. E., Creech, O., and Cooley, D. A. (1954). *Ann. Surg.*, **140**, 290.
 Lodin, H., Rudström, P., and Törnblom, N. (1955). *Acta med. scand.*, **151**, 171.
 Milanés, B., Bustamante, R., Guerra, R., Núñez, A. N., Hernández, A. L., Pérez-Stable, E., McCook, J., and Inigo, J. R. (1952). *Angiology*, **3**, 472.
 Pataró, V. F., Perretta, A., and Navarret, E. E. (1954). *Angiology*, **5**, 1.
 Starer, F., and Sutton, D. (1958). *Brit. med. J.*, **1**, 1255.

The *Disabled Persons (Employment) Act (Northern Ireland), 1960*, includes sections on the minimum age for attendance at courses, amendments as to registration, the provision of sheltered employment by local authorities, and preference for ex-service women in selection for vocational training and industrial rehabilitation courses and for possible employment.

D

IDIOPATHIC STEATORRHOEA AND HAEMORRHAGE DUE TO MALABSORPTION OF VITAMIN K

BY

SIDNEY SHAW, M.D.

Senior Lecturer and Consultant in Clinical Pathology,
 Charing Cross Hospital and Medical School, London

Fanconi (1927) described a haemorrhagic diathesis in three patients with coeliac disease. Later he suggested that this might be due to vitamin-K deficiency (Fanconi, 1938). Kark *et al.* (1940) noted haemorrhages associated with idiopathic steatorrhoea; there was a prolonged prothrombin time that was shortened by administration of vitamin K. A patient with sprue, complicated by haemoptysis and haematuria, was found to have a prolonged prothrombin time (Collins and Hoffmann, 1943). Cooke *et al.* (1953) reviewed 100 cases of idiopathic steatorrhoea and reported an incidence of bleeding in 10%. Haemorrhage was found most often in the skin, but haematuria, haemarthrosis, and bleeding from the gastro-intestinal tract were sometimes present.

On occasion the malabsorption may be discovered in retrospect; Butler and Young (1955) reported a case presenting with severe subcutaneous haemorrhages and menorrhagia. The coagulation time and prothrombin time were both greatly prolonged; the patient responded rapidly to blood transfusion and intravenous vitamin K₁. Subsequent investigation revealed an idiopathic steatorrhoea.

Case Report

A man aged 43 was admitted to hospital on February 3, 1959, with a history of haematuria for five days and extensive bruises and petechiae, especially on the legs but also on the face, chest, and back. He had been treated for diarrhoea, diagnosed as amoebic dysentery, in 1942 while serving in India; after that his bowels had acted normally until 1954, when he had occasional attacks of diarrhoea. This had become worse since 1957, and investigations revealed the presence of steatorrhoea.

On admission there was evidence of an acute haemorrhagic illness, a smooth red tongue, and gaseous abdominal distension. The Hess test was positive and the one-stage prothrombin time was prolonged to 150 seconds (control 18 seconds). Blood examination showed a moderate macrocytic anaemia with leucopenia: haemoglobin, 75% (11.1 g./100 ml.); red cells, 3,020,000/c.mm.; colour index, 1.24; the red cells showed marked anisocytosis, poikilocytosis, and polychromasia; they appeared normally haemoglobinized, and many were macrocytic; white cells, 2,700/c.mm. (neutrophils 75%, lymphocytes 14%, monocytes 11%); platelets, 406,000/c.mm.; serum calcium, 8.1 mg./100 ml. A sternal marrow aspiration showed active erythropoiesis which was chiefly normoblastic but with a moderate incidence of transitional megaloblasts; granulopoiesis and platelet formation appeared normal. Stool microscopy was negative for cysts and amoebae, and cultures were negative for pathogens; the total faecal fat was 45% of the dry weight of the stool (split fat 33% and unsplit fat 12%).

Subsequent barium follow-through radiographs showed some dilatation of the ileum with increase in size of the mucosal folds—appearances consistent with an idiopathic steatorrhoea. Biopsy of mucosa of the upper jejunum showed marked flattening and atrophy of the villi—appearances very suggestive of idiopathic steatorrhoea. The diagnosis was therefore confirmed as idiopathic steatorrhoea.