There is little doubt that tuberculosis could be eliminated quite quickly if everyone were prepared to take sufficient trouble. Meantime the only proper attitude, at least in economically developed countries, is that every death from tuberculosis and every new notification is a failure which ought to have been prevented.

Summary

With 12 to 25 million infectious cases of pulmonary tuberculosis in the world the disease is far from being defeated. Even in economically developed countries the elimination of tuberculosis could be proceeding much more rapidly if present knowledge were more widely and more intensively applied.

The main weapons for the defeat of tuberculosis are miniature radiography to detect cases, chemotherapy to cure them, and B.C.G. to raise the individual's resistance to the disease.

It is suggested that B.C.G. has an important, if accessory, part to play in economically developed countries, and might be even more useful in under-developed countries provided the efficacy of the vaccine is kept constantly under review, both in the laboratory and in the field.

The proper use of miniature radiography is reviewed. It is suggested that this weapon could be used with much greater intelligence and effectiveness.

Chemotherapy should be able to ensure the arrest of virtually all newly diagnosed cases of the disease, and there should be little or no relapse. Evidence for this is adduced. Methods for testing for drug resistance are at present chaotic. Failures of therapy are often due to the use of wrong combinations of drugs or to misleading drug-resistance tests.

With good team-work and proper treatment the elimination of tuberculosis could be greatly accelerated. The Edinburgh figures for mortality and morbidity, though by no means perfect, have shown a dramatic change for the better over the last few years and now compare very favourably with other cities in the United Kingdom, Europe, and the United States. This is held to demonstrate the value of team-work, and of scrupulous chemotherapy, in the elimination of the disease.

I am indebted to many people for much of the work on which this lecture is based. I would like particularly to mention my clinical colleagues, Drs. I. W. B. Grant, N. W. Horne, J. D. Ross, and J. Williamson; my bacteriological colleagues, Drs. Sheila Stewart and A. T. Wallace; Mr. Andrew Logan and the Edinburgh thoracic surgeons; Drs. H. E. Seiler and J. M. Mair, of the Edinburgh Public Health Department; and many able past and present assistants. I must thank Drs. J. E. Geddes, R. N. Johnston, Ian M. Macgregor, W. P. Phillips, and V. H. Springett for providing me with certain British figures, and Dr. B. Pirc, of the World Health Organization, for the figures for cities outside Great Britain. I am most grateful to Miss Una Mackie, of the medical illustration department of Edinburgh University, for preparing the diagrams. The Edinburgh research was supported by generous grants from the Royal Victoria Hospital Tuberculosis Trust.

Note.—In the diagrams all Edinburgh figures for 1959 are calculated from the returns for the first 10 months of the year.

REFERENCES

- Chief Medical Officer (1958). *Report of the Ministry of Health, Part II.* H.M.S.O., London.
- Cowan, K., Horne, W. A., Geddes, J. E., Tait, A. W., and Macgregor, I. M. (1958). "Glasgow's X-ray Campaign Against Tuberculosis." Glasgow Corporation, Glasgow.
- Crofton, J. (1959). *Brit. med. J.*, **1**, 1610.
- (1960). *Brit. med. Bull.*, **16**, 55.
- Daniels, M., and Hill, A. B. (1952). *Brit. med. J.*, **1**, 1162.
- Frimodt-Møller, J. (1959). Address at XVth Conference of International Union Against Tuberculosis.
- Geddes, J. E. (1959). *Hlth Bull. (Edinb.)*, **17**, 14.
- Holm, J. (1959). Address at XVth Conference of International Union Against Tuberculosis.
- Indian Council of Medical Research (1959). "Tuberculosis in India: A Sample Survey, 1955-58," *Spec. Rep. Ser.*, No. 34. New Delhi.
- Logan, W. P. D., and Benjamin, B. (1957). "Tuberculosis Statistics for England and Wales, 1938 to 1955," General Register Office *Studies on Medical and Population Subjects*, No. 10. H.M.S.O., London.
- Medical Officer of Health, Edinburgh (1958). Annual Report of the Public Health Department.
- Medical Research Council (1948). *Ibid.*, **2**, 769.
- (1950). *Ibid.*, **2**, 1073.
- (1953a). *Lancet*, **2**, 217.
- (1953b). *Brit. med. J.*, **2**, 1005.
- (1955). *Ibid.*, **1**, 435.
- (1959). *Ibid.*, **2**, 379.
- Ministry of Health (1959). *Radiological Hazards to Patients. Interim Report of the Committee.* H.M.S.O., London.
- Palmer, C. E., Shaw, L. W., and Comstock, G. W. (1958). *Amer. Rev. Tuberc.*, **77**, 877.
- Pollock, T. M. (1959). *Tubercle (Lond.)*, **40**, 399.
- Rist, N., and Crofton, J. (1960). *Bull. Un. int. Tuberc.*, **30**, 2.
- Ross, J. D. (1959). *Ibid.*, **29**, 710.
- Horne, N. W., Grant, I. W. B., and Crofton, J. W. (1958). *Brit. med. J.*, **1**, 237.
- Seiler, H. E., Welstead, A. G., and Williamson, J. (1958). *Tubercle (Lond.)*, **39**, 339.
- Springett, V. H. (1959). *Ibid.*, **40**, 432.
- Stewart, Sheila, M., Turnbull, F. W. A., and Macgregor, Agnes R. (1956). *Ibid.*, **37**, 388.
- Sutherland, I. (1959). *Ibid.*, **40**, 413.
- United States Public Health Service (1959). *Tuberculosis Chart Series.* United States Department of Health, Education and Welfare, Washington.

A NEW SYNTHETIC PENICILLIN

BY

A. H. DOUTHWAITE, M.D., F.R.C.P.

Senior Physician, Guy's Hospital, London

AND

J. A. P. TRAFFORD, M.B., B.S.

Medical Registrar, Guy's Hospital, London

Since the original communications on the therapeutic value of penicillin G (Chain *et al.*, 1940; Abraham *et al.*, 1941) search has been continued for new effective antibiotics. It has become evident that the penicillin-resistant staphylococcus is an organism whose control presents a major problem. Recent papers have emphasized these difficulties, and in a trial at the Postgraduate Hospital, Hammersmith (Barber *et al.*, 1960), penicillin G was actually withdrawn from routine use in an endeavour to reduce the number of staphylococci in the hospital which were penicillin-resistant. The later introduction of erythromycin (McGuire *et al.*, 1952), novobiocin (Smith *et al.*, 1956; Wallick *et al.*, 1956), and vancomycin (McCormick *et al.*, 1956), which are effective against penicillin-resistant staphylococci, have their own disadvantages, such as the rapid development of resistance on the part of the organism or toxic effects on the eighth nerve.

Batchelor *et al.* (1959) announced the isolation of 6-aminopenicillanic acid. An opportunity presented for the development of synthetic penicillins effective against penicillin-resistant staphylococci. The first of these new synthetic penicillins—namely, BRL 1241 ("celbenin")—is the subject of this trial. The chemical formula of this compound is sodium 6-(2,6 dimethoxybenzamido) penicillanate monohydrate.

Material and Methods

The trial of the compound's effectiveness was directed solely against penicillin-resistant staphylococci.

In the selection of cases the following criteria were adopted: (1) the patients were infected with *Staphylococcus aureus*; (2) the staphylococcus was resistant to penicillin G and did not respond to treatment with other antibiotics; (3) the use of BRL 1241 was confined to cases where an antibiotic was clearly indicated—the mere presence of *Staph. aureus* in carriers was not taken as an indication for treatment; and (4) the production of bacteriological evidence of the presence of the staphylococcus, its sensitivities before treatment, and the examination of the appropriate material—for example, swabs or sputum—after treatment.

The drug itself was kept at -2°C . until use and was administered in the dosage indicated in each case. A dose of 1 g. entailed the injection of 2 ml. intramuscularly. Concentrations in serum and urine were assayed at intervals after injections.

Case 1

A man aged 59 was admitted to hospital in January, 1960, with severe right-upper-lobe pneumonia which followed influenza. White blood count, 16,400. Sputum culture on January 4 showed heavy growth of coagulase-positive *Staph. aureus*, resistant to penicillin, streptomycin, tetracycline, and erythromycin. His temperature ranged between 99 and 102°F . (37.2 and 38.3°C). He was treated, before the results of the sputum culture were returned, with penicillin, streptomycin, and tetracycline without improvement. He then received a five-day course of BRL 1241, 1 g. four-hourly from January 8 to 13. His clinical improvement was striking, and his temperature returned to normal and remained normal thereafter. No staphylococci could subsequently be obtained from the sputum. He was discharged on January 19.

Case 2

A woman aged 60 was admitted to hospital for surgical treatment of carcinoma of the floor of the mouth. Glands were found in the neck and she underwent a two-stage operation on November 17 and December 1, 1959. After operation she developed an intractable wound-infection of the cervical incision. Swab culture on January 12 showed *Staph. aureus* resistant to penicillin, streptomycin, tetracycline, and erythromycin. She had already received penicillin and tetracycline with no effect. Before treatment with BRL 1241 was started her temperature fluctuated between 99 and 102°F . (37.2 and 38.9°C). She was treated with BRL 1241, 1 g. four-hourly for five days. Her temperature became normal in 24 hours and remained so. Culture was negative; she underwent further operation without ill effect on January 21, and was eventually discharged on February 19, after her second operation.

In this case severe cervical cellulitis and cervical suppuration were cleared by BRL 1241.

Case 3

A youth of 18 developed a trophic ulcer on his right leg after poliomyelitis. For two years no healing was achieved, and as the leg was useless he had a below-knee amputation on February 1. Ten days later he developed a wound infection and his stump became red, swollen, and painful. Swab culture on February 16 of discharging pus showed penicillin-resistant *Staph. aureus*. He was treated with BRL 1241, 1 g. four-hourly for five days. Swab on February 23 showed only a few colonies of *Staph. albus*. From this time the wound healed.

This is an example of treatment of wound infection with BRL 1241 and demonstrates its value in this type of case.

Case 4

A man aged 61 was admitted to hospital for amputation of the left leg for peripheral vascular disease. He had chronic bronchitis and emphysema, and after operation a right lower lobe pneumonia occurred. He developed congestive cardiac failure and was extremely ill, without fever but with a pulse rate of 100 – 110 and a respiratory rate of 30 – 40 . Sputum culture of February 29 grew coagulase-positive *Staph. aureus* resistant to penicillin, streptomycin, tetracycline, and erythromycin. He was treated with BRL 1241, 1 g. three-hourly for 10 days, and though the staphylococci disappeared from his sputum he showed no clinical improvement. Chloramphenicol, streptomycin, and "crystamycin" also had no effect. He died on March 15 in congestive cardiac failure.

This patient was suffering from cor pulmonale. BRL 1241 cleared the staphylococci but of course made no difference to the underlying cardiac condition.

Case 5

A man aged 71 was admitted to hospital with a carcinoma of the left bronchus. Left pneumonectomy was performed. After this procedure his condition was not satisfactory in that he continued to have an intermittent pyrexia and was unwell in himself. He had a wound infection which grew *Staph. aureus* on culture, but pneumonectomy-space aspiration showed a clear fluid. He was observed for a period. Aspiration of the pneumonectomy space later revealed thick creamy pus from which *Staph. aureus* resistant to penicillin and streptomycin was grown. He was treated with BRL 1241, 1 g. three-hourly. Ten days later, on March 17, culture of aspirated fluid was sterile.

It was recognized that surgery was indicated to drain this empyema, but it is of note that treatment with BRL 1241 sterilized the fluid.

The most noticeable clinical effect prior to drainage was the disappearance of constitutional disturbance and the rapid reduction in respiratory rate after treatment with BRL 1241. Antibiotics are not advocated as a substitute for surgical drainage of pus, and this patient was thus treated only until surgery was performed.

Case 6

A man aged 44 was admitted to hospital for laryngectomy for carcinoma and block dissection of the left side of the neck. His wound became infected with *Staph. aureus* resistant to penicillin and streptomycin. He was treated with BRL 1241, 1 g. four-hourly for five days. The discharge was much reduced and his wound healed.

In this patient there was considerable risk of spread of a penicillin-resistant staphylococcal infection. After treatment was begun, no spread, in fact, occurred.

Case 7

A man aged 57 admitted to hospital for prostatectomy was found to have an infection of the urinary tract with penicillin-resistant *Staph. aureus*. Prior to prostatectomy he was treated with BRL 1241, 1 g. four-hourly for three days, a catheter specimen then being found sterile. After retropubic prostatectomy he developed a wound infection with *Staph. aureus*. It was mild, and cleared up without treatment.

Case 8

A man aged 40 was admitted to hospital for assessment of his bronchiectasis for surgery. He produced a quantity of sputum daily and had an intractable cough. Sputum culture showed *Staph. aureus* resistant to penicillin. The surgeon was unwilling to perform a thoracotomy in the

presence of penicillin-resistant staphylococci. BRL 1241 was given for five days, 1 g. four-hourly. Sputum cultures were then negative. Bronchoscopic suckings had not shown any pathogens.

Case 9

A man aged 33 was admitted to hospital in March, 1960, with a history of cough and dyspnoea. A diagnosis of bronchiectasis was made, confirmed by bronchogram, and this was found to be localized to the lingula of the lung. On April 5 lingular resection was carried out. This procedure was soon followed by the production of much purulent sputum. Bronchopneumonia was diagnosed. Culture of the sputum grew penicillin-resistant *Staph. aureus*, and he was treated with BRL 1241, 1 g. four-hourly for seven days. After treatment the amount of sputum was greatly reduced and culture showed no evidence of staphylococci. He was discharged one week later.

This was a case of rather severe post-operative staphylococcal pneumonia which responded rapidly and effectively to BRL 1241.

Case 10

A woman aged 38 was admitted to hospital in April, 1960, for exploration of the mitral valve. At operation it was thought that mitral regurgitation was predominant, and in view of her condition it was decided to attempt reconstruction of the mitral valve at a later stage.

Prior to a second operation she developed pneumonia, as evidenced by fever, cough, sputum 2½ oz. (70 ml.) a day, signs of pneumonic changes in the lungs, and radiographic evidence of the same. Sputum grew penicillin-resistant *Staph. aureus*. She was treated with BRL 1241, 1 g. four-hourly, with clearance of the chest signs and radiographic appearances, and the sputum was reduced to negligible proportions. Culture after treatment showed no staphylococci.

This is another case of resistant staphylococcal lung infection which was successfully treated.

Case 11

A woman aged 26 was admitted with a history of infection for two years. In 1957 and 1958 she had a breast and perinephric abscess, but unfortunately no record of the organism or the treatment could be obtained. In 1959 she had a right otitis media complicated by mastoiditis and labyrinthitis, and in June, 1959, she was suspected of developing a right cerebellar abscess; this was not confirmed. In November, 1959, exploration of the right posterior fossa yielded pus in the right cerebellar hemisphere which cultured *Staph. aureus*, resistant to penicillin but sensitive to chloramphenicol. She was treated with chloramphenicol, between 1 and 2 g. daily from November, 1959, to April, 1960, punctuated by short courses of erythromycin, oxytetracycline, and achromycin. Between November, 1959, and January, 1960, the abscess was repeatedly tapped, with growth of the same staphylococcus with the same sensitivities.

In January 26, 1960, a right cerebellar abscess was excised from which the same organism was again cultivated. This *Staph. aureus*, with the same sensitivities, was also cultured from the ventricular and lumbar C.S.F. On March 18 a cutaneous abscess over the occiput developed, growing the same staphylococcus. On March 25 a ventricular tap showed a C.S.F. with the same staphylococcus. She had a hyperpyrexial meningeal reaction with signs of brain-stem involvement and a purpuric rash. The fever and meningitis were not affected by an augmented dose of chloramphenicol. On March 30, BRL 1241, 1 g. four-hourly, was started. Within seven days the C.S.F. white cells fell from 3,380/c.mm. before treatment to 7/c.mm., and her pyrexia settled. On April 15 40 ml. of pus was aspirated from the posterior fossa, but for the first time no organism was grown from this. Treatment with BRL 1241 was continued, but so severe was the residual neurological damage that it was thought

quite likely that she would die. However, she began to improve, and up to the time of writing has continued to do so without any evidence of further staphylococcal infection, although she has marked residual neurological damage.

This woman was suffering from a severe staphylococcal meningitis and cerebellar abscess which did not appear to respond to surgical measures or standard antibiotic treatment. Systemic administration of BRL 1241 apparently cleared the staphylococcal meningitis and sterilized the abscess in the posterior fossa.

Case 12

Two weeks before admission this patient, a woman aged 29, had had an attack of chicken-pox. This had been associated with much scratching and consequent secondary infection. She developed pain in the back, which became worse and was associated with high fever. She was tender over the fifth to ninth dorsal spines and the pains became girdle in type. Paralytic ileus and bladder paralysis developed. Lumbar puncture showed a slow rise of pressure, a protein content of 1,000 mg./100 ml., and 59 white cells. A myelogram showed a block from T 3 to L 1 spines. An extradural abscess was diagnosed. On May 15 extradural penicillin and chloramphenicol were introduced and a "polythene" cannula was left *in situ* after laminectomy, T 9 to 11. Culture of the abscess showed a penicillin-resistant *Staph. aureus* insensitive to chloramphenicol and streptomycin. These were grown repeatedly.

On May 18 BRL 1241 was administered, 3 ml. four-hourly intramuscularly. Culture on May 26 showed persistence of the organism. On May 27 further exploration still showed an extensive abscess. Her legs were paralysed after the second operation. On the evening of operation BRL 1241 was introduced locally through the cannula, 500 mg. being put in every day. Culture on May 30 was sterile. From this date onwards recovery ensued. Her pyrexia settled completely and movement returned to her legs. On June 8 injections of BRL 1241 were discontinued. The only local reaction was severe pain and swelling at the site of injection.

The severity of this woman's condition rendered it inadvisable to treat her with one drug alone, and therefore she had chloramphenicol throughout the period in which BRL 1241 was given. It would seem, however, that its local instillation had sterilized an abscess cavity when chloramphenicol had failed. To make any further claims might well exaggerate the significance of this case.

Case 13

A man aged 41 was admitted with pleuritic pain, high fever, cough, and sputum. Examination revealed consolidation of the right lower lobe, confirmed on chest x-ray examination. Culture of the sputum showed penicillin-resistant *Staph. aureus*. Treatment for five days with BRL 1241, 1 g. four-hourly, resulted in resolution of the chest condition, and no further staphylococci were cultured. This was an acute staphylococcal pneumonia which responded rapidly to treatment.

Comment

Toxic Effects.—A patient outside this series who was sensitive to penicillin G showed no reaction to a test dose of BRL 1241. However, 36 hours after receiving full dosage she developed a widespread erythematous rash which was controlled with large doses of antihistamines. Stewart (1960) quotes an interesting case of a volunteer sensitive to benzylpenicillin who was found to be sensitive to BRL 1241 in comparable dosage.

In common with the experience of Stewart *et al.* (1960) it has been found that the injections are painful and prolonged treatment has caused local reactions. Patients who have received both benzylpenicillin and

BRL 1241 consider the pain of injections is comparable. No other toxic effects were encountered.

Blood and Urine Levels.—These were assayed on patients who were being treated with BRL 1241 whenever this was possible. In all cases assayed, blood levels showed a range of concentration between 18 and 21 $\mu\text{g./ml.}$ at half an hour, 13 to 20 $\mu\text{g.}$ at one hour, 5 to 9 $\mu\text{g.}$ at two hours, and 1.5 to 2.7 $\mu\text{g.}$ at three hours. By the fourth hour serum concentrations were invariably below 2 $\mu\text{g.}$ Urinary assay showed that 75% of the dose administered in one day was excreted in the urine—for example, six injections of 1 g. were given in 24 hours and 4,372 mg. were excreted in the urine over the same period, which is 73% of the injected dose. Knudsen and Rolinson (1960), in their studies of absorption and excretion, showed similar results.

Discussion

The real test of effectiveness of this drug is whether it failed to eradicate staphylococci in any instance or caused such severe side-effects that it had to be withdrawn. So far in our limited experience no case failed to respond to treatment.

In Case 4, with staphylococcal infection of the lung, the organism was eliminated but the patient died of his underlying cor pulmonale. It was not thought that any antibiotic treatment would have made much difference to his cardiac condition.

Stewart *et al.* (1960) report one patient (Case 14, aged 2) who relapsed after treatment of her multiple boils with BRL 1241. They report that staphylococcal infection was eradicated in all their other cases but was not necessarily extirpated from the anterior nares.

Summary and Conclusions

A clinical trial of BRL 1241, a derivative of 6-aminopenicillanic acid, is reported.

Thirteen cases of staphylococcal infection resistant to penicillin G and/or other antibiotics are recorded.

No serious toxic effects were encountered. The drug was successful in eradicating staphylococci in all.

The results reported suggest strongly that the drug is effective against the resistant staphylococcus.

We are grateful for supplies of BRL 1241 to Beecham Research Laboratories Limited, whose registered trade name for this drug is "celbenin." Our thanks are due to the physicians and surgeons at Guy's Hospital who allowed us to treat their cases.

REFERENCES

- Abraham, E. P., Chain, E., Fletcher, C. M., Florey, H. W., Gardner, A. D., Heatley, N. G., and Jennings, M. A. (1941). *Lancet*, 2, 177.
- Barber, M., Dutton, A. A. C., Beard, M. A., Elmes, P. C., and Williams, R. (1960). *Brit. med. J.*, 1, 11.
- Batchelor, F. R., Doyle, F. P., Nayler, J. H. C., and Rolinson, G. N. (1959). *Nature (Lond.)*, 183, 257.
- Chain, E., Florey, H. W., Gardener, A. D., Heatley, N. G., Jennings, M. A., Orr-Ewing, J., and Sanders, A. G. (1940). *Lancet*, 2, 226.
- Knudsen, E. T., and Rolinson, G. N. (1960). *Brit. med. J.*, 2, 700.
- McCormick, M. H., Stark, W. M., Pittenger, G. E., Pittenger, R. C., and McGuire, J. M. (1956). *Antibiot. Ann.*, 1955-6, p. 606.
- McGuire, J. M., Bunch, R. L., Anderson, R. C., Boaz, H. E., Flynn, E. H., Powell, H. M., and Smith, J. W. (1952). *Antibiot. and Chemother.*, 2, 281.
- Smith, C. G., Dietz, A., Sokolski, W. T., and Savage, G. M. (1956). *Ibid.*, 6, 135.
- Stewart, G. T. (1960). *Brit. med. J.*, 2, 694.
- , Nixon, H. H., and Coles, H. M. T. (1960). *Ibid.*, 2, 703.
- Wallick, H., Harris, D. A., Reagan, M. A., Ruger, M., and Woodruff, H. B. (1956). *Antibiot. Ann.*, 1955-6, p. 909.

A NEW PENICILLIN (BRL 1241) ACTIVE AGAINST PENICILLIN- RESISTANT STAPHYLOCOCCI

BY

R. KNOX, M.D., F.R.C.P.

Professor of Bacteriology, Guy's Hospital Medical School,
London

The discovery of 6-aminopenicillanic acid, which has been described as the penicillin "nucleus" (Batchelor, Doyle, Nayler, and Rolinson, 1959), has led to the synthesis of new penicillins, some of which show antibacterial activity very different from that of penicillin G. One of these, BRL 1241 ("celbenin") shows remarkable activity against penicillin-resistant staphylococci (Rolinson, Stevens, Batchelor, Cameron-Wood, and Chain, 1960). The present paper describes some laboratory studies of this compound and some comparisons between it and three other penicillins. BRL 1241 is sodium 6-(2,6-dimethoxybenzamido)penicillanate monohydrate.

Methods

BRL 1241 and "broxil" (phenethicillin) (Knudsen and Rolinson, 1959) were obtained from Beecham Research Laboratories. Penicillin G and penicillin V were commercial preparations from Glaxo Laboratories Ltd. Commercial penicillinase from Burroughs Wellcome and Co. was used. Antibacterial activity of the different antibiotics was measured, usually in nutrient broth, by the serial tube dilution test, using young broth cultures undiluted or in various dilutions. Cultures were incubated at 37° C. and minimum inhibitory concentrations read at intervals. Bactericidal tests were carried out as described in the text.

Antimicrobial Activity of BRL 1241

The serial tube dilution test was used to compare the antimicrobial activity of BRL 1241, penicillin G, penicillin V, and broxil against some common micro-organisms of medical importance. The results are summarized in Table I. These confirm and extend the

TABLE I.—Antibacterial Activity of Four Different Penicillins

	Minimum Inhibitory Concentration ($\mu\text{g./ml.}$)			
	Pen. G	Pen. V	Broxil	BRL 1241
"Oxford" staphylococcus	0.02	0.01	0.01	1.0
<i>Staph. pyogenes</i> (E3) ..	125	62	>250	2.0
<i>Str. pyogenes</i> (group A) ..	0.005	0.01	0.01	0.2
<i>Str. viridans</i> ..	0.01	0.01	0.02	0.1
* <i>N. meningitidis</i> ..	0.2	0.2	0.2	6
<i>Proteus vulgaris</i> ..	15	>250	>250	>250
<i>Ps. pyocyanea</i> ..	>250	>250	>250	>250
<i>Salm. typhi</i> ..	7.5	62	>250	>250
" <i>paratyphi</i> A ..	7.5	125	>250	>250
" " B ..	31	>250	>250	>250
" " C ..	0.2	7.5	62	62
<i>E. coli</i> ..	15	125	>250	>250

Inoculum: 0.02 ml. of 1/100 dilution of overnight broth cultures inoculated into 1 ml. of broth containing serial doubling dilutions of antibiotic. Minimum inhibitory concentrations read after 18 hours, incubation at 37° C.

* This was tested in serum broth and the inoculum was 0.02 ml. of undiluted serum broth culture.

observations of Garrod (1960), who pointed out that there may be significant differences in the action of different penicillins and that it is unwise now to assume that a micro-organism sensitive to one penicillin will necessarily be sensitive to another. It can be seen that BRL 1241 was still inhibitory in a concentration of about 2 $\mu\text{g./ml.}$ against a penicillin-resistant staphylococcus, which under the conditions of the test was