thyroid replacement therapy (Thompson *et al.*, 1929). The "paretic" Lange curve in the C.S.F. of Case 2, however, requires further explanation. It suggests a relative increase in C.S.F. gamma-globulin due to the combination of hypothyroidism and hyperglobulinaemia (found in many cases of autoimmune thyroiditis, including non-goitrous myxoedema). Such an increase was demonstrated electrophoretically in myxoedema by Bronsky *et al.* (1958).

Summary

Antibody to thyroglobulin was detected in the spinal fluid of five patients with autoimmune thyroiditis. In two out of three patients who had complement-fixing antibody in the serum, this antibody could also be detected in the C.S.F. In three patients who were hypothyroid the total C.S.F. protein was raised.

We are grateful to Professor G. A. Smart and Dr. T. H. Boon for allowing us to study their patients. We also wish to acknowledge the able technical assistance of Mrs. J. Johnson.

REFERENCES

- Boyden, S. V. (1951). *J. exp. Med.*, **93**, 107.
 — and Sorkin, E. (1955). *J. Immunol.*, **75**, 15.
 Bronsky, D., Shrifter, H., de la Huerza, J., Dubin, A., and Waldstein, S. S. (1958). *J. clin. Endocr.*, **18**, 470.
 Gries, G., Aly, F. W., and v. Oldershausen, H. F. (1953). *Klin. Wschr.*, **31**, 644.
 Owen, S. G., and Smart, G. A. (1958). *Lancet*, **2**, 1034.
 Roitt, I. M., and Doniach, D. (1958). *Ibid.*, **2**, 1027.
 Thompson, W. O., Thompson, P. K., Silveus, E., and Dailey, M. E. (1929). *Arch. intern. Med.*, **44**, 368.
 Tiselius, A. (1937). *Trans. Faraday Soc.*, **33**, 524.

SIMULTANEOUS OCCURRENCE OF INTRACRANIAL ANEURYSM AND ARTERIOVENOUS MALFORMATION REPORT OF CASE SUCCESSFULLY TREATED

BY

RONALD FINE, M.R.C.P.Ed.

Senior Registrar in Neurology

ALASTAIR PATERSON, M.B., Ch.B., F.R.F.P.S.

Consultant Neurosurgeon

LESLIE STEVEN, M.A., M.B., Ch.B., D.M.R.D.

Consultant Radiologist

AND

**J. B. GAYLOR, M.A., M.B., Ch.B., B.Sc., F.R.C.P.Ed.
F.R.F.P.S.**

Consultant Neurologist

*From Department of Neurology, Western Infirmary,
Glasgow, and Neurosurgical Unit, Killearn Hospital,
Glasgow*

Both cerebral aneurysms and arteriovenous malformations are well-recognized causes of spontaneous subarachnoid haemorrhage. However, the simultaneous occurrence of both lesions is a great rarity, and up to 1959 only 14 examples had appeared in the literature (Boyd-Wilson, 1959). Recently a 49-year-old woman was investigated at Killearn Neurosurgical Unit for subarachnoid haemorrhage; she was found to have an aneurysm arising from the right posterior inferior cerebellar artery and also an arteriovenous malformation situated in the posterior upper surface of the right cerebellar hemisphere.

Because of the paucity of such cases in the literature, and especially as both lesions were successfully treated, this case is reported with a brief review of the previous literature.

Case Report

A 49-year-old widow was admitted to Knightswood Hospital on December 26, 1959. For the previous five years she had suffered from headaches, which were occipito-vertex and often unilateral, though either side might be affected. At first the headaches were very severe in the afternoons after hard work, but recently they had been occurring in the early mornings. On the day of her admission she had wakened with headache, which she regarded at first as her usual attack of migraine. On bending down to wash her face, however, the headache became intense. She then noticed stiffness of her neck and heard peculiar noises inside her head; a little later she started vomiting and had diarrhoea.

On admission she complained of severe occipital headache, blurring of vision, and occasional double vision.

On examination she was found to be grossly obese. The blood-pressure was 180/110 and the apex beat was displaced 1 in. (2.5 cm.) outside the midclavicular line. The aortic second sound was accentuated. She was irritable and mildly confused. There was definite nuchal rigidity, with Kernig's sign. No localizing signs, however, were discovered: the deep tendon reflexes were brisk, but both plantar reflexes were flexor. A lumbar puncture disclosed a heavily blood-stained fluid at a pressure of 350 mm. of water. Gradually over the next few days the headache became less severe, but she occasionally felt dizzy and continued to see double. Three weeks later, on January 18, 1960, she suddenly became comatose; after lumbar puncture the cerebrospinal fluid was found to be uniformly blood-stained and under increased pressure. By the next day she had recovered consciousness, but was mildly confused and disorientated: the left plantar reflex was noted to be extensor.

These findings necessitated her transfer for further investigation at Killearn Neurosurgical Unit on January 22. On examination there, in contrast to her condition four days previously, she was alert and fully orientated. The nuchal rigidity persisted. Ophthalmoscopic examination revealed marked papilloedema with subhyaloid haemorrhages and attenuated arteries. With some defect of fixation the visual fields showed a left homonymous cut, complete on the right and involving the lower quadrant on the left. On deviation of the eyes to the left a few nystagmoid movements were observed in the initial phase of the deviation. Other cranial-nerve functions were normal. Muscular tone and power were undisturbed and sensation was intact throughout. Both plantar reflexes were flexor and the deep reflexes were symmetrical. No bruit was audible over the orbits or over the parieto-occipital areas.

Bilateral carotid angiography on January 22 showed no deformity apart from some spasm of the terminal portions of both internal carotid arteries. Four days later (January 26) a right vertebral angiogram revealed a small aneurysm arising from the right posterior inferior cerebellar artery, and an arteriovenous malformation probably situated on the upper surface of the right cerebellar hemisphere (see Figs. 1 and 2).

On January 28—a month after the first haemorrhage—exploration of the right posterior fossa was carried out by Mr. A. Paterson. The right posterior inferior cerebellar artery was identified, and was noted to be very tortuous and sclerotic. The artery was followed distally in an anterior direction around the anterior aspect of the tonsil and then posteriorly on to the hemisphere where it broke into two small branches. No aneurysm could be found and eventually the posterior inferior cerebellar artery was clipped a considerable distance from its origin. The impression at the time was that the clip was distal to the aneurysm. The upper dural flap was then raised until the upper surface of the cerebellum could be seen, and close

to the midline lay a small arteriovenous malformation fed by two large arterialized veins, one coming through the tentorium and the other running medially. Both vessels were clipped and the angioma was completely excised.

After the operation the patient was confused: this state gradually diminished and by February 16 she was alert and fully orientated. A further vertebral angiogram showed that the small aneurysm, previously noted arising from the posterior inferior cerebellar artery, was not filling and that

the artery was apparently clipped proximal to and near the origin of the aneurysm. The arteriovenous malformation was shown to be occluded with several silver clips (see Fig. 3).

There was no residual neurological abnormality, except for the resolving subhyaloid haemorrhages in both fundi, papilloedema of the right disk, and a residual left hemianopic defect.

Discussion

In a recent survey of nearly 3,000 consecutive post-mortem examinations Chason and Hindman (1958) found an incidence of almost 5% of intracranial aneurysms, and therefore it may be assumed that 5% of cases of arteriovenous malformations also have intracranial aneurysms. However, from a survey of the literature on cerebral angiomas, Paterson and McKissock (1956) are the only authors who reported the coexistence of these two lesions. In only four of their 110 cases of cerebral angiomas were aneurysms present. It is possible that the simultaneous occurrence of these two lesions will become more apparent as complete routine arteriography comes to be performed in all cases of subarachnoid haemorrhage where the exact source of bleeding is in doubt. Certainly the existence of multiple aneurysms in the one patient has been realized with the wider application of arteriography; and, similarly, it is likely that more cases of both arteriovenous malformations and aneurysm will become apparent, especially as both probably share the same congenital aetiology.

A further interesting feature about this case was the presence of the aneurysm on the posterior inferior cerebellar artery. This is a most unusual site for aneurysms, and is the first seen in this unit. Walton (1956) lists only one such case in a series of 140 intracranial aneurysms. Again, however, it is felt that lesions in this site will become more obvious with the wider application of percutaneous vertebral angiography.

Summary

A case is reported with an aneurysm on the right posterior inferior cerebellar artery and an arteriovenous malformation supplied by branches from the superior cerebellar artery and the posterior inferior cerebellar artery. Both lesions were successfully treated, and the patient made an uneventful recovery save for residual ocular signs. It is suggested that the coexistence of intracranial aneurysms and cerebral angiomas will become more obvious with the wider application of complete arteriography in subarachnoid haemorrhage.

REFERENCES

- Boyd-Wilson, J. S. (1959). *J. Neurol. Neurosurg. Psychiat.*, **22**, 218.
Chason, J. L., and Hindman, W. M. (1958). *Neurology (Minneapolis)*, **8**, 41.
Paterson, J. H., and McKissock, W. (1956). *Brain*, **79**, 233.
Walton, J. N. (1956) *Subarachnoid Haemorrhage*. Livingstone, Edinburgh and London.

FIG. 1.—Right lateral vertebral angiogram, showing a small bilobed aneurysm on the posterior inferior cerebellar artery and a small angiomatous malformation on upper surface of cerebellum.

FIG. 2.—Antero-posterior view of right vertebral angiogram, showing the aneurysm and the angiomatous malformation.

FIG. 3.—Post-operative right lateral vertebral angiogram, showing the silver clip proximal to the site of the aneurysm on the posterior inferior cerebellar artery. The angiomatous malformation has been occluded by silver clip.

The major cause of death in Australia in 1959 was heart disease. Official statistics show that heart diseases caused more than one-third of the 89,000 deaths in the year. The principal causes of death were: heart diseases 30,000, cancer 13,000, strokes nearly 12,000, accidents 5,000, and pneumonia 3,000. The infant mortality rate was 21.5 per 1,000, one point higher than last year. The lowest infant mortality rate in Australia was 20.5 in 1958 (Radio Australia News, July 29).