chromosome abnormality. The presence of Hb-F could have been caused by an alteration in the genetic pattern of the marrow cells occurring either at the time of onset of the leukaemia or subsequently. There is, however, the other possibility that a pre-leukaemic chromosomal abnormality which caused persistence of Hb-F had been present from early life.

It will be interesting to discover whether this association of Hb-F and leukaemia is fortuitous or whether it will prove to be an important observation from an aetiological as well as from a diagnostic point of view.

Summary

The case of a girl with leukaemia presenting as haemolytic anaemia associated with 19% foetal haemoglobin is described. Evidence suggestive of a defect in her red cells was found.

The possible significance of these findings is discussed in relation to the genetic hypothesis for the aetiology of leukaemia.

We wish to acknowledge our indebtedness to Professor H. Scarborough and Professor J. Gough for their advice and encouragement, to Dr. H. Lehmann for measurement of Hb A₂, and to Mr. R. G. Wood for his help with the isotope studies.

ADDENDUM.—Marrow and peripheral blood have now been cultured from this patient during a second remission induced by busulphan. There were no morphological abnormalities in the chromosomes of cells cultured from the peripheral blood, and the chromosome number was 46. The chromosomes of the few intact marrow cells which could be examined in detail were morphologically normal. This investigation was done by Dr. C. M. Clarke and Mr. G. Breckon, of the M.R.C. Microbiological Research Unit at Harwell, to whom we are greatly indebted.

REFERENCES

- Baikie, A. G., Court Brown, W. M., Jacobs, P., and Milne, J. S. (1959). *Lancet*, **2**, 425.
 Beaven, G. H., and White, J. C. (1953). *Nature (Lond.)*, **172**, 1006.
 Chanarin, I., Dacie, J. V., and Mollin, D. L. (1959). *Brit. J. Haemat.*, **5**, 245.
 Hickling, R. A. (1953). *Lancet*, **1**, 57.
 Kilpatrick, G. S., Withey, J. L., and Hardisty, R. M. (1960). In preparation.
 Lehmann, H. (1959). *Brit. med. Bull.*, **15**, 40.
 Mollison, P. L. (1959). *Brit. med. J.*, **2**, 1035.
 Singer, K., Chernoff, A. I., and Singer, L. (1951). *Blood*, **6**, 413.
 Valentine, W. N., and Beck, W. S. (1951). *J. Lab. clin. Med.*, **38**, 39.
 Wetherley-Mein, G., Epstein, I. S., Foster, W. D., and Grimes, A. J. (1958). *Brit. J. Haemat.*, **4**, 281.
 — Hutt, M. S. R., Langmead, W. A., and Hill, M. J. (1956). *Brit. med. J.*, **1**, 1445.
 White, J. C., and Beaven, G. H. (1959). *Brit. med. Bull.*, **15**, 33.
 Wintrobe, M. M. (1956). *Clinical Hematology*, 4th ed., p. 990. Kimpton, London.

This year a home-printer's course for instructing spastics in the printing of Christmas cards was organized with the help of a retired master printer. It was designed to enable those who were too severely handicapped to be placed either in open industry or sheltered employment to earn money by working at home. As most of those who attended the course were too handicapped to handle small type a system was devised whereby all printing could be done by means of printers' blocks. The course was sponsored by the National Spastics Society's employment department. Christmas cards printed by spastics can be obtained from Sundew Cards, 32 The Mall, Ealing, London W.5.

DYSPHAGIA LUSORIA PRESENTING AS A SUPERIOR MEDIASTINAL SYNDROME

BY

ROY MORRIS, M.D., M.R.C.P.Ed.

Johannesburg, South Africa

In 1787 Dr. David Bayford carried out a necropsy on a woman who died of starvation, after having suffered from dysphagia from the age of 13 to 60. At first he could find no obvious cause for her symptoms, but ultimately his dissection led him to the discovery of an aberrant subclavian artery, which arose from the left side of the aortic arch and coursed between the oesophagus and the trachea to supply the right arm. He concluded that this anomalous vessel had pressed upon the oesophagus, causing dysphagia. He applied to this disorder the name of dysphagia lusoria, *lusus naturae*, meaning a freak of nature (Asherson, 1952).

Dysphagia lusoria in infancy signifies oesophageal compression by any congenital arterial abnormality, such as right aortic arch, any aberrant great vessel, vascular ring, double aortic arch, and right aortic arch with an aberrant left subclavian artery. In adults the term usually denotes oesophageal compression due to an aberrant right subclavian artery, this usually being the only abnormality.

Incidence

In the adult type of disorder the right subclavian artery arises from the left side of the aortic arch beyond the origin of the left subclavian artery. This vessel then courses to the right in one of three ways. It may run behind the oesophagus (80%), between the oesophagus and the trachea (15%), or in front of the trachea and oesophagus (5%) (Neuhauser, 1946). It will be seen from this that classical dysphagia lusoria as described by Bayford is probably extremely rare. Aberrant subclavian artery has been estimated to occur in about 1% of otherwise normal people. Some anatomical studies place this figure as high as 2%, but radiological studies obtain about 0.18% of positives (Palmer, 1955). The abnormality therefore is common, but the production of symptoms is very rare. I was able to find only seven references to dysphagia lusoria in the English language over the past ten years.

FIG. 1.—A. The caudal portion of the right aortic arch normally disappears. B. The cephalic portion disappears and the caudal part persists, forming the aberrant right subclavian artery.

If there were any point in doing so, I feel that the radiological diagnosis could be made as often as the anatomical diagnosis, using the method of subclavian-artery catheterization as described by Raphael *et al.* (1952). Symptoms are usually so mild and non-progressive that this procedure is not warranted.

FIG. 2.—Origin and course of the aberrant right subclavian artery, with the dilated diverticulum of Kommerell. Adapted from Gross (1946).

Anatomy

Embryologically, an arteria lusoria represents the persisting distal descending portion of the fourth right aortic arch, there having been obliteration of the proximal arch in reverse of the normal situation. Normally the fourth right aortic arch forms the right aortic arch, which disappears. Its persistence leads to the formation of an aberrant subclavian artery (see Fig. 1). At the origin of the abnormal vessel there may be an aneurysmal dilatation known as the diverticulum of Kommerell (see Fig. 2).

Symptomatology

The commonest and mildest symptom is a feeling of food sticking under the manubrium sterni. Regurgitation is rare, but has caused a fatality (Felson *et al.*, 1950). The dysphagia may be mild and chronic or mild and recurrent. It may become progressively worse or, as in the case described below, it may be fairly severe with acute exacerbations. Palmer (1955) mentions the symptoms of disability, fear, and fatigue, but it is difficult to see how these arise. The dysphagia is mainly for solids, but also for liquids if taken fast. There is no pain, and the swallowing-time is usually normal. In infants respiratory symptoms usually predominate; in adults, dysphagia. Certain signs have been described, but they are rare. For example, there may be inequality of the radial pulses (Copleman, 1945), and, rarely, trophic changes in the affected limb (Felson *et al.*, 1950). I have not encountered in the literature anything approaching the symptoms experienced by our patient.

Diagnosis and Treatment

Many cases of so-called globus hystericus must have been referred to psychiatrists under the mistaken impression that no organic cause existed for their symptoms. Greater awareness of the symptomatic possibilities of aberrant subclavian artery will lead to the diagnosis in an increasing number of cases, and, in fact, this disorder may come to be rather commonplace.

Apart from the clinical presentation, there are four main diagnostic measures which may be employed to prove the origin of the patient's symptoms.

1. Barium swallow: If the typical appearances are encountered there is usually no need to proceed further with diagnostic measures. In the postero-anterior view, if the oesophagus contains too much barium no indentation will be seen (Fig. 3). If, however, there is merely a trace of barium present one will notice an oblique radiological defect about 0.5 cm. wide passing upwards from left to right on the posterior aspect of the oesophagus just above the aortic arch, which is usually in its normal position (Fig. 4). On the postero-lateral aspect of the oesophagus the abnormal vessel produces a typical notch (Fig. 5). If the vessel runs in front of the oesophagus a similar P.A. appearance is noted, but in this case the notching will be anterior. Should the vessel course in front of the trachea, no oesophageal defect will be encountered. In the P.A. view, above the aortic knuckle, there may be a prominence of the left mediastinal shadow of pulsatile nature, probably due to Kommerell's diverticulum (Stauffer and Pote, 1946).

2. Angiography may be employed to demonstrate the aberrant vessel, but this form of investigation carries a risk and is better avoided.

3. Oesophagoscopy may reveal the presence of a pulsatile oesophageal defect, either anteriorly or posteriorly, in about 40% of cases (Palmer, 1955).

4. The right subclavian artery may be catheterized without the injection of dye, and, in conjunction with a barium swallow, the defect can be proved to be due to an aberrant right subclavian artery.

FIG. 3.—Oesophagus overfilled with barium. The defect is not apparent. P.A. view. FIG. 4.—Oesophagus now contains little contrast medium, and the obliquely running defect is clearly visible. FIG. 5.—Notching of the postero-lateral border of the oesophagus is evident in the left anterior oblique projection.

In infants surgical intervention usually becomes imperative, because of the serious nature of the underlying abnormalities (Gross, 1946; Gross and Neuhauser, 1951). There is no record of an adult patient having been operated on. Palmer described 11 cases of dysphagia lusoria. He claims that "detailed explanation of the x-ray films to the patient proved to be an effective therapeutic gesture." There is no follow-up available on these patients.

Case Report

A married woman aged 54 was first seen on July 16, 1959. About 25 years previously she had noticed some dysphagia on swallowing pills, which had to be crushed, or on attempting to swallow large or hard pieces of food. Her dysphagia was constantly present, but it fluctuated in intensity. About 14 years ago she found it impossible to lie flat in bed and had become used to having six to eight pillows. When lying flat she felt as if something in the chest moved up into her throat and caused swelling of the neck. This swelling would rise as far as the ears, but no higher, and it produced an intense sensation of discomfort amounting almost to *angor animi*. At the same time she noticed hoarseness of her voice and what she described as attacks of "laryngitis," by which she meant total inability to phonate, not associated with pain. All these symptoms waxed and waned. She would have attacks lasting up to three months, and intervals of comparative freedom lasting from 18 months to two years. During an attack she insisted that her neck was actually swollen. Her eyesight was not affected; she did not suffer headaches, and her general health was quite unimpaired.

There was one other curious symptom which defied explanation. She insisted that since childhood she had been unable to tell left from right, and this caused her much embarrassment at school and while she was in the Women's Auxiliary Services. During parades the platoon might be given the order to turn left or right and would duly go off in that direction, but she would invariably go off in the opposite direction. The only way she could find her bearings was by feeling her fingers and working out which was her left hand and which her right. In October, 1953, after the loss of a child, her symptoms became more noticeable, and she consulted a physician. An "x-ray of the oesophagus" was done, and this was reported to have been normal. After several visits to her doctor she was referred to a psychiatrist for treatment of the "globus hystericus." The psychiatrist made a note on his card that, in his opinion, this patient, although outwardly stoical, was a hysteric. He explained to her that she had a "wall of tears" in the chest, and that psychiatric treatment would enable her to have a good cry and rid herself of this wall. She refused treatment, and went back to her work as a doctor's receptionist. She had learnt to live with her symptoms previously, and, in the absence of medical help, would continue to do so.

She consulted me at a time when her symptoms were particularly bad. She said that she had been moderately well until a few days previously, when she had developed a cold, associated with tightness in the chest, "as if something was stuck." She had a feeling of wanting to belch, but the gas would only come up as high as the angle of Louis, and then would be propelled down and up the oesophagus, crackling and rumbling all the while. This symptom had persisted day and night for three days. She had been forced to sleep sitting straight up, because of a choking feeling on attempting to lie down, and she had avoided taking solids completely, as they appeared to stick in the region of the upper sternum. The swallowing of fluids, especially tea, gave some relief. Her voice had become more hoarse than usual, and the whole of her neck was swollen and tight. The attack, above her usual background of symptoms, had begun suddenly, accompanied at first by some nausea which soon passed off. Apart from

the removal of her gall-bladder and appendix, she had always been in good health, and there were no other symptoms.

Examination

The patient weighed 128 lb. (58 kg.), and measured 5 ft. 4 in. (162.5 cm.). There was nothing peculiar about her habitus, and she looked younger than her 54 years. She sat upright in comfort, but became extremely dyspnoeic when reclining. The base of the neck looked full and tense, and the anterior jugular veins stood out like distended cords. The fullness of the neck suggested bilateral diffuse thyroid enlargement, which was in fact present, and over the region of the thyroid gland the neck was flushed. Her bearing was that of someone being gently choked by external physical means. In fact, she described the feeling in her neck as one of being so choked. The external jugular veins were not seen to be distended. There was diffuse tenderness around the base of the neck anteriorly. Blood-pressure readings were 110/70 (right) and 120/80 (left). The rest of the physical examination was unremarkable, except for the appearance of the optic fundi. Here the veins were over full and the disk edges were blurred bilaterally. After treatment these abnormalities resolved completely. The E.C.G. was normal, and the urine contained neither albumin nor sugar. The clinical impression gained was that of a retrosternal goitre which has undergone a recent haemorrhage.

The radiological diagnosis was made by Dr. A. Berezowski. In the postero-anterior view the typical oblique radiological defect in the oesophagus was seen (Fig. 3), and there was also an indentation on its postero-lateral aspect, caused by the aberrant vessel (Fig. 4). Aortography confirmed the presence of an aberrant right subclavian artery with a slight dilatation at its origin, the remainder of the artery being of normal calibre (Fig. 6). The right and left common carotids and left subclavian artery were normal.

Oesophagoscopy was performed by Mr. A. I. Lichter. From 20 to 22 cm. from the upper incisor teeth an obliquely running indentation of the posterior oesophageal wall was apparent. The indentation ran from left to right in an

FIG. 6.—Angiogram, showing the aberrant right subclavian artery arising from the aorta, with slight dilatation at its origin.

upward direction corresponding to the appearances on the barium swallow. This area showed forceful pulsation. The remainder of the oesophagus showed no abnormality. While the oesophagoscope lay on and beyond the pulsatile swelling, the right radial pulse was completely obliterated. On withdrawing the oesophagoscope to just above the swelling, the pulse returned immediately. The left radial pulse remained constant throughout. These observations on the pulse were made by Mr. M. Arnold.

Operation (Mr. A. I. Lichter)

The left chest was entered through the bed of the resected fourth rib. The mediastinal pleura was incised and displayed the posterior mediastinal structures. The left subclavian artery and the aberrant right subclavian artery were identified. The aberrant artery arose about 0.5 cm. beyond the left subclavian from the posterior aspect of the aortic arch. The origin of the aberrant artery showed a slight aneurysmal dilatation. The artery was traced backwards and obliquely upwards towards the right. It was freed from its bed, where it lay between the posterior wall of the oesophagus and the spine. The thoracic duct was identified and retracted. The artery was freed up to the right border of the oesophagus and ligated at that level and at its origin from the arch of the aorta. The intervening portion of the artery was excised and on examination was found to show slight atheromatous changes. No fibrous bands accompanied the artery. In its final position the distal stump of the artery lay along the right border of the oesophagus. The chest was closed in layers with an underwater tube drain. The patient made a rapid recovery from the operation, with the immediate disappearance of all symptoms.

Results

At a follow-up examination on February 24, 1960, the patient insisted that immediately after coming round from the anaesthetic she was aware that all symptoms of the past 20 years had vanished. She was now able to lie flat, and, in fact, slept with no pillows. The hoarseness of the voice had disappeared. The optic fundi presented normal appearances, and there was no thyroid swelling. She was able to swallow anything and everything with ease. The size of her neck had returned to what it was 25 years ago.

Discussion

An aberrant right subclavian artery is fairly rare, but dysphagia lusoria is extremely so. It is possible that, with increasing recognition of the clinical syndrome, the two conditions will approach each other in incidence. It has been suggested that when symptoms are present they are due to the diverticulum of Kommerell; but this is unlikely, because symptoms may be episodic and also because, as the present case demonstrates, a relatively slight dilatation at the origin of the artery is compatible with symptoms severe enough to warrant a thoracotomy. Moreover, that part of the artery which causes the oesophageal obstruction is usually not dilated.

It is unlikely that increasing atherosclerosis or tortuosity of the vessel is important in symptom production, as will be seen from the age of onset in Palmer's (1955) 11 cases: three patients were 20-29 years, two 30-39, three 40-49, one was 50-59, one 60-69, and one 70+.

In symptom production we have to answer three questions.

1. Why, in the majority of cases, are there no symptoms at all? Possibly in most people the capacity of the superior mediastinum is such that the anomaly is accommodated freely.

2. Why, in some cases, are symptoms chronic and progressive, as in Bayford's original patient? Probably a combination of factors, local and general, associated with advancing age play a part—such as progressive aneurysmal dilatation of the vessel, emphysema, and kyphoscoliosis. All these would decrease the capacity of the already tightly packed mediastinum.

3. Why, as in the case described above, is this abnormality capable of producing a picture of superior mediastinal obstruction? It is clear from the clinical picture described that there was obstruction to the venous return from the head and neck, as evidenced by the swollen appearance of the neck, the flush, the enlarged thyroid, the appearance of the optic fundi, and the choking sensation. Anatomically it is unlikely that the innominate veins or superior vena cava would be compressed directly by the arteria lusoria, as they lie far forward in the mediastinum in front of the trachea. They could, however, be compressed, owing to their relatively low internal pressure, by oedema, although why any swelling should occur is not clear. Such oedema would be unlikely to affect the arteries with high internal pressures, or the trachea with its more rigid structure. The peristaltic waves of the oesophagus might, with difficulty, produce sufficient mechanical force to propel food through the area of relative obstruction—hence the lack of total obstruction and regurgitation.

So far as can be ascertained from the English-language literature, this is the first reported case of dysphagia lusoria presenting as a superior mediastinal syndrome. It is also the first time that an adult patient has been operated upon for the relief of symptoms due to an aberrant right subclavian artery.

Summary

Adult dysphagia lusoria is due to an aberrant right subclavian artery compressing the oesophagus. This disorder is probably not as rare as the paucity of case reports would suggest; methods are therefore described by which more cases of "globus hystericus" might be found to have an organic basis.

A case is described presenting as chronic superior mediastinal compression: complete disappearance of the symptoms and signs occurred after surgical resection. This is the first known adult patient to be operated upon for the condition.

My thanks are due to Dr. A. Berezowski for the x-ray films; to Mr. A. I. Lichter for his operation notes; and to Mr. A. M. Shevitz, of the Witwatersrand University Medical School, for the photographs.

REFERENCES

- Asherson, N. (1952). *J. Laryng.*, **66**, 111.
Copleman, B. (1945). *Amer. J. Roentgenol.*, **54**, 270.
Felson, B., Cohen, S., Courter, S. R., and McGuire, J. (1950). *Radiology*, **54**, 340.
Gross, R. E. (1946). *Ann. Surg.*, **124**, 532.
— and Neuhauser, E. B. D. (1951). *Pediatrics*, **7**, 69.
Neuhauser, E. B. D. (1946). *Amer. J. Roentgenol.*, **56**, 1.
Palmer, E. D. (1955). *Ann. intern. Med.*, **42**, 1173.
Raphael, R. L., Schnabel, T. G., and Leopold, S. S. (1952). *Radiology*, **58**, 89.
Stauffer, H. M., and Pote, H. H. (1946). *Amer. J. Roentgenol.*, **56**, 13.

The sixth portion of the Signet Library, Edinburgh, consisting of seventeenth, eighteenth, and nineteenth century pamphlets and a few books, was sold recently at Sotheby's. A first edition, 1811, of Sir Charles Bell's *Idea of a New Anatomy of the Brain* went for £410.