ON THE
ANATOMY AND PHYSIOLOGY
OF THE
CYSTICERCUS TENUICOLLIS.

By C. B. ROSE, F.R.C.S., F.G.S., &c., SWAFFHAM.

Received April 11th—Read May 9th, 1848.

IN the year 1833, I contributed a paper on the Vesicular Entozoa,* to the London Medical Gazette, in which, I fear, I did little more than repeat the errors of my predecessors ; still, in that imperfect summary, I *did* call the attention of zootomists to the existence of an organ, hitherto unobserved, which I had discovered in my examinations of the *Cysticercus tenuicollis*, and which is readily observed *within* the caudal vesicle, suspended from that portion where what has been called the body of the entozoon terminates and the vesicle commences. I long hoped that the above notice would have caught the eye of some comparative anatomist possessing more facilities for the investigation than myself, and have incited him to undertake a more elaborate dissection of the entozoon ; but my anticipations not having been realized, I again took up the subject, and the result of my researches I have now the honour of laying before the Fellows of this Society.

Previously to entering upon the history of my own inquiries and dissections, I will briefly trace chronologically the steps of preceding naturalists in the examination of this genus of the entozoa *Cystica*.

As early as the year 1654, Bartholin cursorily described

* London Medical Gazette, 1833, vol. xiii. p. 205.

Hydatids (believed by Rudolphi to be Cysticeri), which he considered to be animated, but did not esteem them to be worms.*

About the same period, Steno discovered bladders (*Cysticercus tenuicollis*) in the omentum of a reindeer, which he described as if pregnant, or containing one within another.† It is most probable that Steno mistook the omental cyst for the parent hydatid, and the true parasite for the offspring; the *cyst* being commonly so transparent, as to give it very much the appearance of a bladder of water, suspended by a pedicle.

Hartmann, in the year 1685, was the first to pronounce this hydatid to possess *independent* vitality, he having placed one (which he had taken from the omentum of a goat) in warm water, and observed its undulatory movements.‡ He also noticed the retracted neck and head.

Our countryman, Tyson, in 1691, made discoveries to the same extent, without being aware of what Hartmann had published. Tyson considered the caudal vesicle to be the stomach of the worm. His paper (published in the *Philosophical Transactions*) is entitled, "*Lumbricus hydropicus*; or, an Essay to prove that Hydatides, often met with in morbid animal bodies, are a species of Worms, or Imperfect Animals."§

Pallas, who wrote in 1766, was the first to enter elaborately into the natural history of vesicular worms: having minutely examined this hydatid, and imagined that its mouth resembled that of the *Tæniæ*, he named it *Tænia hydatigena*.||

Goeze, Zeder, Steinbuck, and Rudolphi, did but perpetuate the errors of their predecessors, contributing nothing to the knowledge previously acquired respecting their anatomy and generation. The recent writers on the same subject, of this country, Houston, Knox, Owen, Gulliver, and Goodsir, have

* *Entozoorum, sive Verm. Intest. Hist. Nat.*, vol. i. p. 119.

† *Idem*, loc. cit.

‡ *Idem*, p. 112.

§ *Opus cit.*, vol. i. p. 112.

|| *Elench. Zooph.*, p. 413; also *Spicileg. Zoolog.*, Fasc. 12, p. 42.

made us much better acquainted with the anatomy of this entozoon; to whose contributions I shall have occasion to refer, *en passant*.

Till the period of Harry Goodsir's researches, the *Cysticercus* had been described as made up of a head armed with a double row of hooklets, and furnished with four mouths or suckorial cups, a neck, body, and caudal vesicle; and various had been the offices assigned to these components of the parasite. Of the parts connected with the imaginary head, some authors have considered the hooklets to be for the purpose of attachment to the containing cyst, and the whole *body*, therefore, an organ of prehension; others have looked upon them as teeth; and a third party thought them subservient to the generative process. With regard to the four oscula, Steinbuck, finding that they were not patent, and consequently could not be mouths, conceived the preposterous notion (as late as 1802), that they performed a respiratory function. Rudolphi and several earlier naturalists described them as true mouths, for the purpose of sucking in nutriment, each mouth opening into a canal, which either passed separately towards the caudal vesicle, or, uniting into one common canal, terminated in the caudal vesicle. Some recent writers have given credence to the above statements of Rudolphi, and also reiterated them; but whoever reads his remarks upon this portion of the subject will be quite satisfied that he had promulgated conjectures rather than observations. The late Dr. Houston, of Dublin, in reference to the so-called mouths, says, "These depressions are called mouths, meaning thereby patulous orifices, but from my own observations I believe them to be covered over by the same kind of transparent membrane as that which completes the anterior planes, for I had abundant opportunities of remarking that the fluid of the caudal vesicle when squeezed along the neck into the head, protruded and rendered them prominent, without making its escape from apertures in their centre."* The appearance of their being covered by a

* An account of Hydatids found in the omentum of an axis-deer. Dublin Journal of Science, &c., vol. viii. p. 271.

membrane, and made to protrude by propelling the fluid of the vesicle against them from behind, is fallacious. I have repeatedly produced a similar aspect with the compressing instrument (a paper knife), when effecting the extrusion of the retracted head. It is, in fact, an eversion of the suctorial cups; and with regard to the passage of the fluid along the body, the Doctor was likewise deceived; for the fluid cannot be forced along the body, without first rupturing the lining membrane of the vesicle, as I shall show hereafter.

Dr. Roget also imbibed the erroneous idea of four mouths communicating with the caudal vesicle, which he pronounced a stomach:* but it is manifest that he did not investigate the point himself.

Even Professor Owen has in part adopted the errors of the early naturalists, as he states that, "In the larger *Cysticerci*, lateral canals may be traced from the suctorious pores extending down the body towards the terminal cyst, but they appear not to terminate in that cavity, the fluid of which is more probably the result of secretion or endosmosis."†

After this brief summary of the labours and opinions of those who have preceded me in this field of inquiry, I proceed now to lay before you the result of my own investigations and dissections.

The *Cysticercus tenuicollis* infests chiefly Ruminants, is continually met with in sheep, and occasionally in the hog; its proper *habitat* is the peritoneum, or the pleura, but it is very much more frequently seen in the former than in the latter; a serous membrane appears to afford its proper nidus, as it never invades the parenchyma of a viscus, but is entirely confined to the peritoneal or pleural covering. It is invariably found in a cyst, and in the abdomen is met with enveloped, either in the omentum, or hanging from various parts within the peritoneal bag. The formation of the cyst, and the morbid changes which it frequently undergoes, I

* Bridgewater Treatise, by Dr. Roget, vol. ii. pp. 84, 113.

† Dr. Todd's Cyclopædia of Anatomy and Physiology, vol. ii., article Entozoa, p. 131.

shall only briefly notice hereafter ; but, on those matters, I would more particularly direct attention to an elaborate paper by the late Dr. Houston,* before referred to, and in which will be found a full and accurate description of this envelope, in its normal and also in its anormal states.

The entozoon itself is a vesicular body, constructed of translucent membranes, containing a limpid fluid, and from its anterior part is produced an elongated cylindrical structure, hitherto considered the true body, or vermiculum, but which, from analogies afterwards to be offered, I am disposed to call a *proboscis*.†

This hydatid acquires a magnitude sufficient to weigh twenty-five drachms, and to possess fluid contents amounting to three ounces by measure. I have repeatedly met with them of this bulk ; to what extent they may continue to grow I have not determined : the above sized specimens were taken from a one-and-a-half year old sheep—a *sheerling*—and as their *habitat* offers no physical impediment to their progressive increase, we may fairly infer that an extension of life to the animal would be productive of a further growth of the parasite ; doubtless, also, their magnitude is a measure of their age. It is almost unnecessary to remark that, where a colony of these hydatids is established, they occur of various sizes ; I have had twenty-two specimens sent me from the same abdomen, varying from the size of a nutmeg to that of a goose's egg. The large hydatids do not appear to distend their cysts at all more than the smaller ones, the cyst unquestionably growing with its inhabitant.

A glutinous fluid is invariably found within the cyst, occupying the small space between it and the hydatid ; with the aid of a microscope it is discovered to consist of a large

* Opus cit., p. 277, et seq.

† It appears that Mr. Harry Goodsir is the first who has taken a more correct view of the conformation of this entozoon ; he considering the caudal vesicle to be its proper body, and the hitherto-called body but a pedicle. *Vide* Anatomical and Pathological Observations, No. XIII. p. 91.

proportion of oil globules, with a few globules of *pus*, mechanically incorporated with serum (?) : it contains a large quantity of albumen.

When the hydatid is first removed from the cyst, particularly if through a large opening, its form is nearly spherical ; and it exhibits at the anterior part of its long axis an oval, labiated, opaque protuberance—the retracted proboscis.

Everting the proboscis elongates the vesicle, and the hydatid becomes pyriform. When thus unfolded, the proboscis exhibits coarse transverse plicæ, and the vesicular body a delicately plicated surface, similarly arranged. The eversion is not a difficult manipulation,* if attempted upon a recent specimen ; but after some decomposition has taken place, it cannot be so readily effected, by reason of its delicate texture easily giving way.

Structure of the vesicle.—The vesicular *body* is constructed of two laminæ, or coats ; the external or fibrous, and the internal or granular : its form is spheroidal, but with its proboscis protruding it is pyriform, and it is replete with a transparent fluid, to the amount of three fluid ounces in large specimens.

The *external coat* is made up of three sets of fibres ; two of the three decussate each other at right angles, forming a network of longitudinal and transverse fibres ; the third set are seen to run in all directions, although those passing transversely are most distinct, and under a power of one-eighth, greatly resemble the fibres of the middle coat of an artery, as figured by Müller, Plate II. fig. 6 ; they undoubtedly form an elastic tissue, a kind of structure which alone accounts for the remarkable degree of elasticity possessed by the entozoon.

The Cysticerci exhibit muscular motion, and it is asserted that they protrude their retracted proboscis during the undu-

* I have found no difficulty in unfolding the retracted part, by pressing it gently with the fore-finger of my left hand, at the same time assisting its eversion with a paper-knife, used in a pressing and rather scraping manner.

latory movements excited by immersion in warm water; the latter circumstance I have never witnessed, although I have frequently immersed them, and watched their proceedings. There has not yet been detected in their structure, either the fibre of voluntary or of involuntary muscle; still, as modern anatomists consider that an animal so low in the scale of organization requires a less complicated muscular texture, we may, perhaps, assume that the *decussating* fibres possess the requisite muscular function.* The *internal* coat is composed of a form of albuminous matter having a granular character: it has incorporated with it some carbonate of lime, and probably a larger portion of the phosphate; it has also a great abundance of oil-globules interspersed throughout; it possesses no fibrous texture, and when decomposition has, to a certain degree, advanced, it separates from the external coat, and falls to the bottom of the vesicle.† The internal coat, after having formed a lining to the fibrous lamina, is reflected over an organ (to be described hereafter) hanging from the proboscis into the fluid of the vesicle, and to which it forms a covering. In passing from the parietes of the vesicle on to the above-mentioned organ, it forms a septum (fig. 1, *h*) between the cavity of the vesicle and the interior of the proboscis, and therefore precludes the escape of the fluid of the vesicle in that direction, and, consequently, the distention or protrusion of the proboscis. The *fluid* of the vesicle has been analysed by Collard de Martigny, and others; the former found it to be composed of *water* 96·5, *albumen* 2·9, and salts, for the most part *chloride of sodium*, 0·6. By Göbel it is said to contain some phosphate of lime.‡

The Proboscis.—The external membrane of the vesicular body is produced anteriorly, to the extent of from a quarter

* Professor Owen says,—“Although in every order, both of the Parenchymatous and Cavitory worms, living specimens have been observed to exhibit sufficiently conspicuous motions, yet their muscular fibre is not always distinctly eliminated in them.”

† The separation and deposit is well exhibited in preparation No. 224, of the entozoa *Cystica*, in the Museum of the Royal College of Surgeons.

‡ Simon's Chemistry (translation by Day), vol. ii. p. 484.

of an inch to nearly an inch in length, according to the age of the hydatid, and forms a tapering, cylindrical, transversely plicated appendage, terminated by a subtetragonal enlargement, armed with a double row of hooks, and having placed at each angle (?) a suckorial cup (fig. 1, *b*). The external tissue of this appendage is thickly beset with oval corpuscles, throughout its texture, from its commencement at the vesicle, to within a very short distance of its extremity; they are very closely arranged upon its surface, but evidently loosely attached, for a slight scraping readily detaches some of them.

The *corpuscles* have been described, and their size minutely determined, by Mr. Gulliver;* he has shown, also, that they contain a notable quantity of carbonate of lime in their composition; a circumstance which accounts for the opacity observed in the texture, of which they form so considerable a proportion. He thinks it probable that they are the ova of the *Cysticercus*; Klencke† also believes them to be ova; and Goodsir describes as ova, similar bodies found in the pedicles upon the vesicle of the *Cænurus cerebialis*.‡ Having been favoured with an opportunity of examining the oval corpuscles under one of our most powerful microscopes, and, at the same time, had the advantage of Mr. Busk's valuable assistance, I do not hesitate to pronounce (having his high authority for so doing) that they are not ova, but are, in reality, as far as structure goes, what Mr. Gulliver satisfactorily demonstrated them to be, "oval bodies possessing a granular texture, with a calcareous shell;" thus giving firmness to the structure—in short, forming a coat of mail to an organ whose duties, in all probability, required it. It will be seen, by the very accurate figures given in illustration of Mr. Gulliver's paper, that but few corpuscles are interspersed through the enlarged terminal portion of this organ, that part in which the suckorial cups

* *Medico-Chirurg. Trans.*, vol. vi. 2nd series, p. 4.

† Professor Siebold's Report on Helminthology, translated by Busk for the Report on Zoology, Ray Society, 1847.

‡ *Trans. of Royal Society of Edinburgh*, vol. xv. 1844, p. 561.

and coronet are situated; which is undoubtedly a necessary provision, for, had it been otherwise, the flexibility required by the function of the suctorial cups could not have existed, and their presence, consequently, would have been rendered useless.

I have once met with a congeries of the oval corpuscles, congregated into a circular patch, upon the external membrane of the vesicle, and very remote from the proboscis; to the naked eye it appeared a small white speck, about half a line in diameter, but which, by the microscope, was resolved into an assemblage of corpuscles, as in fig. 2. The circumstance can, of course, be only viewed as one of the inscrutable freaks of nature, analogous to the occurrence of maculæ studded with hairs, upon the human integument.

The hooks.—The circlet of hooks attached to this organ is composed of thirty-two: they are alternately placed in and out, forming a double row; its form and arrangement are so accurately described by Mr. Gulliver, in the paper before referred to, that I have but few observations to make respecting it. At the point of junction of the curved spine with its clavated arm, or *manubrium*, is a process described by Mr. Gulliver as “a short blunt lateral branch, or process;” after repeated examinations, particularly when the hooks were thrown back, and the circle was fully expanded, I have determined that the process is not placed laterally, but anteriorly, in a line with the manubrium and apex of the hook. The hooks are alternately long and short, the difference depending chiefly on the length of the manubrium. The former are 1-125th of an inch in length; the latter, 1-250th. The process of the short hooks I have found to be *bifid* (fig. 3, *a*), while that upon the long ones is formed as Mr. Gulliver has described.

The form, structure and arrangement of the hooks lead me to infer that they are capable of free movement around the centre to which they are attached; and we cannot but suppose that there are special muscles for the designed actions of the hooks. Under this impression, I am induced to think that the processes are the points of insertion of those muscles, although my microscope has not at present laid

open any trace of such a structure. Muscular action must be required, either for the convergence or divergence of these instruments; and, supposing divergence, or their expanded form, to be the quiescent state of the circlet of hooks, convergence—a forward movement, a kind of flexion—would be its instinctive and active movement, effected by muscular action; then, admitting elasticity as the antagonistic force to the muscular (a property greatly prevailing in all parts of the hydatid), an antagonism effectively employed in the economy of testaceous mollusks, by its expanded position would be recovered, and thus, with one muscle only to each, could the requisite alternate actions of the hooks be efficiently produced. The office of this interesting structure, this elegant coronet (for it is a very beautiful microscopic object), appears to be solely that of a mechanical irritant—an instrument wherewith to scratch the interior of the vascular cyst, for the purpose of exciting it to secrete a pabulum for the sustenance of its tenant,—an opinion I was led to adopt from having always observed, when examining the *Cænurus cerebrialis in situ*, that that portion of the *cyst* immediately in contact with its proboscis was much more vascular than any other part of it.*

It has been observed by those authors who have met with this entozoon in a state which showed that it had died previously to the death of the infested animal, that it was invariably found with its proboscis completely unfolded, even to the entire exposure of its crown of hooks. It has always occurred to me to find it so likewise, and the circumstance may, in my opinion, be thus explained. There is every reason to believe that the proboscis is retracted by a muscular power, the energy of which will be in direct ratio with the vigour of the hydatid. The noxious food furnished by the morbid action going on in a diseased cyst necessarily debilitates and ultimately destroys the muscular power of the parasite; its instinctive struggle for food—"a stimulus of necessity"—keeps the organ protruded, and in which state it

* See Todd's Cyclopædia, vol. ii. p. 131.

remains at its death. Dr. Knox, in an interesting paper on the *Cysticercus cellulosæ*,* writes:—"Whilst examining these hooks or spines under powerful glasses, I remarked that they were arranged in a double row, and, at the base of each alternate one, so far as I could observe, there was a small rounded body, resembling an ovum or egg, and their appearance has suggested to me the idea, that these rounded bodies may be the young *Cysticerci* developed at the base of each hook;" but the Doctor also says, "I throw out this opinion merely as a conjecture." I have myself, aided by a most satisfactory display of the parts, repeatedly searched in vain for these "rounded bodies at the base of each alternate hook" in the *Cysticercus tenuicollis*; and as it is probable that they would be more readily recognised in this much larger individual, I do not hesitate to conclude that no such bodies exist, and that Dr. Knox in all probability merely saw one of the oval corpuscles, so thinly implanted hereabout, in the position he describes.

The Suctorial Cups.—The nature of these concavities has given rise to much discussion. Up to a very recent period they were considered by zootomists to be patulous mouths for the absorption of nutriment; but since it has been satisfactorily demonstrated that they afford no channel to the interior of the hydatid, a suctorial function has been appropriated to them.

The cups are circular, and, when examined under the microscope, with a power of one-eighth, exhibit a sharp outline; they are abrupt hollows, by which I mean, not shallow. Their interior is perfectly smooth, and there is no opening in their fundus. With the same magnifying power, concentric fibres are distinctly seen around their margins, and I believe that I have seen transverse fibres dipping towards the interior at right angles to the circular ones. In this instance, also, can it be doubted that these fibres possess the property of muscularity, although the microscope does not detect in them the characteristics of those of a muscle

* *Lancet*, vol. ii., 1837-38, page 397.

of either animal or organic life? Fig. 2 of Mr. Gulliver's plate* does not give a correct delineation of the *cups*, as seen in a recent specimen mechanically unfolded; they are made to appear *convex*, or protruding.

In the study of the physiology of this parasite, various offices have been assigned to the appendage now occupying our attention. From its early history, up to a recent period, it has been considered the head and body of the worm, absorbing nutriment by four mouths, and attaching itself to the cyst by its crown of hooks. Houston having assured himself that the so-called mouths were not patent, looked upon it as a rudimentary part—a type of transition to an animal of a higher grade. Gulliver leans to the idea that it took a share in the generative process, for he writes:—"It is an interesting question, whether the spherules of the caudal vesicle may not be considered as nuclei, which advance to the neck or body of the worm, and there become invested with cells, and so formed into the complete oval corpuscles."† And Goodsir, taking a somewhat similar view, observes:—"The *pedicles* of these two latter animals (*Cænurus* and *Cysticercus*) are analogous to the reproductive nucleus."‡ For my own part, I can see in it but the analogue of the anterior extremity of an *Echinorhynchus*, or of a *Bortriocephalus*, and therefore have denominated the entire pedicle the proboscis.

Thus far, our attention has been occupied with the anatomy of the external parts of the entozoon. We will now proceed to the internal. Upon opening, by a crucial incision, the vesicular *body* of the hydatid,§ and carrying the antero-

* Transactions of Royal Med.-Chir. Society, vol. vi., second series, plate 1.

† Transactions of Royal Med.-Chir. Society, vol. vi., new series, p. 8.

‡ Anatom. and Pathol. Observations, page 93.

§ The readiest mode of dissecting the vesicular entozoa is the following:—Procure a *wooden plate*, or trencher, and paint it black. Upon this fix your hydatid (after having everted its proboscis), by passing a pin through the anterior portion of the proboscis. Now fill the plate with cold water, when, upon opening the vesicle, as above directed, and pinning the anterior flaps back, you will have the pendulous organ distinctly exhibited, and in this way only can you readily examine it, and manipulate with facility.

posterior incision forwards quite up to the commencement of the proboscis (previously everted), the observer will notice upon turning aside the anterior flaps, a transparent gelatinous filament floating in the fluid of the vesicle, and pendulous from that part of the interior of the vesicle where the proboscis commences (fig. 1, *f*). On carefully laying open the dense texture of the proboscis, and turning it aside, he will find that he can trace this pendulous body passing forward for three-fourths of the length of the former. We may therefore describe it as arising high up within the proboscis, and passing along it till it enters the vesicle, where it hangs floating in the liquid contents. I am inclined to think that, within the proboscis, its form is that of a hollow cylinder,* to admit of the inversion of that body; but within the vesicle, although retaining the cylindrical form, it appears to be solid, and tapers to a blunt point (fig. 4, *f*). From this point, a single filament (fig. 4, *f*), or two, or more, or sometimes a tuft (fig. 4, *g*) of slender filaments, is suspended, which filaments either float in the fluid, or are loosely attached to the parietes of the vesicle by still more slender filaments. These filaments may be detected by pulling the pendulous body with forceps, when a dragging of the parietes will be observed at the point of their attachment. The tissue of this pendulous body, arising from the interior of the proboscis, is very intimately connected with the tissue of the latter, particularly where it terminates in the vesicle; for it is impossible to separate one from the other without bringing, with the former, portions of the inner layer of the latter, having an abundance of oval corpuscles attached.

This pendulous organ is not always met with as a single body, but is more frequently than not accompanied with one, two, and even three off-sets: in fig. 5 it will be observed there are three, the principal organ hanging loose, and much

* I have passed a fine bristle into it, after the anterior fourth part of the proboscis has been cut off; but its texture is so delicate that it may have pierced its substance, and not entered a tube. In fig. 4, a bristle is exhibited, passing between it and the surrounding tissue.

longer than the others, an adjoining off-set recurved with its extremity adhering, a second off-set enveloped in the inner coat of the vesicle, and the third very short, and slightly projecting. It is not an unfrequent circumstance to meet with the principal organ, and also its off-sets, entirely embedded in the inner coat of the vesicle. The length of this organ is chiefly determined by the growth or age of the hydatid, but not invariably so; for although in a large hydatid you will never meet with a short pendulous body, still in a small hydatid you occasionally find a disproportionately long one, particularly if you include the slender filaments, or tuft of them.

The texture of this organ is only to be ascertained with the microscope; every part of it is very strong, and remarkably elastic, even to the finest filaments. It is composed of cellular tissue; of elastic tissue resembling that of the parietes of the vesicle, but of a more delicate texture; and also of a tissue, which, with a power of 200 diameters, appears in the form of longitudinal fibres, passing in parallelism. In dilatations of the organ which are not unfrequently seen, the fibres are observed to form volutions.

These fibres are moreover continued into the slender portions of this pendulous body and tufts. With the aid of the microscope, there are also observed in this organ deposits of granular matter and minute oil-globules, both of which are, I am inclined to think, confined to its investing membrane, which, as before noticed, is derived from the internal coat of the vesicle, and reflected over in the same manner as the pericardium is over the heart.

Of the office of this organ, or to what function it is subservient, I have but recently been enabled to form a rational opinion. In 1833, just after I first discovered it, possessed with the then prevalent idea that the four suctorial cups were mouths, I conjectured that this pendulous body and its appendages were portions of an alimentary canal,* cæca, for

* London Medical Gazette, 1833, vol. xiii. p. 205.

instance; but in 1845, I was so fortunate as to find, upon opening a large hydatid, a cluster of young ones sprouting from the upper part of the pendulous body, and, in addition, one about midway down its length, and also another at its extremity; these are well exhibited in fig. 6, but they are magnified about two diameters. In fig. 7 may be observed a single young hydatid, considerably developed, and at the upper part of the pendulous body also, which is seen floating from behind it.

I have before adverted to the circumstance of the organ from which the young are seen to sprout, not being in every instance found suspended in the vesicle, but lying enveloped in its internal coat. I possess a specimen with two or more young ones sprouting from filaments connected with an organ thus circumstanced. I have also met with instances where the young are being developed at a greater distance from the germinating organ than even in the last specimen; and in these examples, also, a slender filament may, with a pocket lens, be traced passing from the young hydatid towards the upper part of the pendulous organ.

What office, then, does this organ perform in the economy of the parasite? From the examples above given, of the development of young hydatids within its confines only, I cannot but infer that it is a special reproductive organ; and as in those instances where the young are apparently sprouting from remote portions of the vesicle, filaments may be traced connecting them with the principal organ, I propound, as a legitimate inference, that a special organ for reproduction is provided in the genus *Cysticercus*, and that the germinating function does not appertain generally, throughout the parietes of the vesicle, as in *Cænurus*. With this view of the function of the organ, I venture to denominate it a *Gemmarium*, and, for the convenience of demonstration, the more slender extremity and filaments, with or without tufts, I designate the *fimbria* (fig. 4, g).

On the nutrition of the entozoon.—Helminthologists having

ascertained that the *Cysticercus* did not possess *suctorial mouths*, they then considered that it must be by imbibition throughout the whole surface of its vesicular body it received its nutriment. In a preceding part of this paper I have shown that the enclosing cyst of the hydatid holds within it a fluid, in quantity more considerable than is required for mere lubrication, and in quality eminently fitted for the nutrition of so simple an animal. It is therefore highly probable, that the parasite does derive its nourishment from the fluid of the cyst, which it calls forth, or rather excites the secretion of when required, by irritating the interior of the cyst with its armed proboscis. The fluid of the cyst is composed of serum, having apparently a smaller proportion of water than that of the blood, and of oil-globules, &c. By *endosmosis* the hydatid imbibes the more attenuated particles of it, to the exclusion of the more *granular* particles of the albumen, as unfitted for the delicate structure of the entozoon, and, as it appears from Matteucci's experiments, ill-adapted for endosmosis. I have been led to this explanation of the mode of nutrition by the following facts: first, the large proportion of albumen in the secretion within the *cyst*; and second, the small trace of it in the fluid of the *vesicle* of the worm. But from whence, then, is derived the material for the *solid* structures of the entozoon? From the fluid of the *cyst*, also, in the first instance, undoubtedly; and as the fluid of the vesicle *does* contain some albumen, if we acknowledge the aid of *catalysis*, the material for the granular internal and fibrous external coats will be readily furnished. Endosmosis carries the materials, catalysis eliminates them, and exosmosis deposits them. It appears to me, indeed, that it is only by the aid of endosmose and exosmose we can explain the growth of animals so low in the scale of organization as not to possess vessels of any description.

On the *reproduction* or propagation of the cystic entozoa generally, we have still much to learn. With respect to the species particularly under our notice, my observations have

enabled me to contribute some additional information in this division of its economy. As regards its *dissemination*, we remain in utter ignorance. Whether these entozoa are generated as ova or gemmæ it matters not: incarcerated as they are, their mode of escape and dissemination is at present a sealed book.

In reference to their reproduction, Mr. Gulliver has considered his *oval corpuscles* to be ova; Mr. Busk and myself, after careful and repeated examinations, are satisfied they are not so. Mr. Goodsir speaks of ova in the pedicles of the *Cænurus cerebialis*; but no one can doubt, on perusal of his observations, that he also has mistaken the oval corpuscles for ova: as they possess similar characters to those of the *Cysticercus*, he has unquestionably fallen into the same error as Mr. Gulliver; therefore, the elaborate description of the process of their development by Mr. Goodsir* I can only contemplate as exceedingly imaginative. But M. Klencke, in this matter, has surpassed all other helminthologists in successful and felicitous experiments and observations: he has not only seen and measured (!) the ova of *Cysticerci* in the blood of animal, but, *mirabile dictu*, has propagated and disseminated the *Echinococcus* by *inoculation*! I cannot better exhibit to you the degree of estimation in which his opinions are held by his continental brethren, than by quoting the words of Professor Siebold, of Freiburg, from his "Report on Helminthology," for 1843-4:†—"It is to be hoped that Klencke will desist from these researches, which he has, in a short time, so completely exhausted, and the results of which have advanced to such an extraordinary extent our present possible knowledge, and relinquish a field, the praiseworthy and careful cultivation of which by other naturalists, though but slow, will be attended with more permanent results."

Professor Owen, and also most naturalists of the present day, consider all hydatids to be reproduced by gemmæ only.

* Anatomical and Pathological Observations, p. 90.

† Reports on Zoology, translated from the German, and published by the Ray Society. 1847.

This eminent zootomist observes, "No trace of a generative apparatus has hitherto been detected in the cystic entozoa. They would seem to be gemmiparous, and to have the reproductive power diffused over the whole cyst (*vesicular body*); at least in the *Acephalocysts*, in which the young are not developed from any special organ, or limited to any particular part of the cyst."* Such a mode of reproduction certainly obtains also in the *Cænurus*, as I am enabled to show by a pregnant specimen of *Cænurus* from the rabbit, in my possession; but, in the genus *Cysticercus*, I have shown that a special organ is appropriated to the generative function, instead of its appertaining to the vesicle generally—a step in advance in the scale of organisation. In the reproduction of the *Cysticercus*, we have, at present, been able to show only one process by which its species is propagated—that is, *gemmation*; it has, at the same time, been demonstrated, that the gemmæ are developed *within* the parent; it is, therefore, manifest, that no *dissemination* of the species can be effected by this mode of reproduction and development; even had the development taken place from the *exterior* of the entozoon, its dissemination would not have been facilitated a single step further, unless indeed, very simple ova,† or perhaps embryonic gemmæ, are extruded from some external part of the entozoon into the cavity of the cyst, and of such infinitesimal size as to allow of their being taken from thence into the general circulation of the infested ani-

* Dr. Todd's *Cyclopædia of Anatomy and Physiology*, vol. ii. p. 137.

† "No germinal vesicle has (we believe) yet been detected in the ovum of the hydra; and it seems to be questionable whether the entire ovum is not really homologous with the vesicle of Purkinje, which may itself contain a store of nutriment, adequate to the development of so low and simple a creature, without the additional supply furnished by the yelk in which this vesicle is elsewhere imbedded. It seems to be clearly established, however, that this separated body contains the essential parts of a true ovum, and is altogether distinct in its character from a *bud*."—See the article on the Development and Metamorphoses of Zoophytes, in the *British and Foreign Medico-Chirurgical Review* for January 1848, page 197.

mal, to be deposited in other parts of it, and also to be carried to its posterity.* But, I am entering the wilds of conjecture; still, as zoophytes are known to propagate by ova as well as gemmæ,† we may surely be permitted to imagine that the vesicular entozoa may do so likewise.

In the specimen, fig. 7, where the young hydatid is considerably advanced in growth, I cannot discover the proboscis. Does its formation await a further development of the entozoon? or are we to look upon this offspring as a "nurse," and admit it as an instance of "alternation of generations," according to Steenstrup? Not having seen a sufficient number of gradations of growth in the young of the *Cysticercus* to determine this point, I can only state with reference to it, that, in the development of the young of the *Cænurus* of the rabbit, the *proboscides* are visible in *all* the young (and several generations occur upon one common parent), and at an early period of their growth.

Of the exciting cause of, or the circumstances which mediate lead to, the *generation* of hydatids in animals, nothing certain is known; but it has very long been believed that an unsound or innutritious diet is productive of their appearance. Mr. Lawrence long since stated,‡ that the *Cysticercus cellulosæ* does not occur in the *wild* hog, although it is not unfrequently met with in the tame animal, constituting *measly pork*; we must, therefore, conclude that domestication (probably through its food and confinement) exerts considerable influence in their production. It is, likewise, a well-known fact, that monkeys and animals of the deer kind, when kept in menageries and consequently placed in an unnatural condition, are very obnoxious to the generation of vesicular entozoa in their viscera and cellular tissue.

* With the aid of the microscope, I have repeatedly searched for ova in the secretion of the *cyst*, but never discovered anything resembling them.

† Brit. and For. Med.-Chir. Review for January 1848, page 196, line 14, et seq.

‡ On Monstrous Productions, in the Trans. of Med.-Chir. Society, vol. v., first series.

I have myself caused the *Cysticercus pisiformis* to be developed in the omentum of rabbits, by feeding them on cabbage leaves alone; an experiment made many years since by the celebrated Jenner. A highly intelligent occupier of a warren has informed me that *bladdery* rabbits are most numerous in dry seasons, when the herbage is necessarily short and scanty; *bladdery* rabbits are those infested with a *Cænurus* in the inter-muscular cellular membrane of the trunk and extremities.

It has been stated, that in sheep fed on salt-marshes *Acephalocysts* are not generated: on this point I can only observe, that sheep are never fattened on salt-marshes; and that sheep fed as lambs on salt-marshes, and afterwards fattened, *are* infested with the *Cysticercus tenuicollis*: also from my inquiries, I am led to believe that sheep fattened on a luxuriant and sound herbage are much less frequently infested with the vesicular parasites. Dr. Budd, in his admirable work on the "Diseases of the Liver," makes the following remark:—"Hydatids are met with in all conditions of life; but seem to be more frequent among the poor than among the rich."*

Having now communicated the extent of my knowledge on the anatomy and physiology of this entozoon, I purpose to offer a few brief observations on the morbid changes which not unfrequently take place in it. It has long been remarked that this *Cysticercus* is every now and then found dead, and when so circumstanced is invariably enclosed in a thickened cyst. A somewhat recent writer upon this subject, the late Dr. Houston, elaborately and accurately described the changes occurring in the cyst, and attributed them to the previous death of its inhabitant. My own observations have led me to arrive at a different conclusion. It has occurred in my researches, that the cyst must have arrived at a considerable degree of thickness before the hydatid died, judging from the slight change in the latter. I have also taken living hydatids from thickened

* Op. cit., page 360.

cysts ; and we may easily conceive that when, from its thickened state, the cyst is no longer dilatable, its living tenant could no further grow, and it would die in consequence. What then, it may be asked, is the exciting cause of the thickening of the cyst? I reply—the constant demand upon it, of food for its inhabitant, leading to the frequently repeated irritation of its interior by the armed proboscis of the parasite. I have been partly led to adopt this opinion, from having found pus-globules in the viscid secretion of the cyst—a sure indication that something more than a healthy stimulus had been going on in that locality.

In the dissection of a cyst considerably advanced in morbid thickening, it will frequently be found more distinctly separable into two layers than in its normal condition, and it will be also seen that, while the outer layer is almost cartilaginous and consequently opaque, the inner layer remains in its originally transparent state ; so much so, that without a previous knowledge of the circumstance, the latter might very easily be mistaken for the hydatid itself: such an occurrence must have taken place during Mr. Goodsir's inquiries into the structure of the cystic entozoa, or he could not have given the following description of the cyst of the *Cysticercus*. He commences his section (VI.—Of *Cysticercus*) in these words: "*Cysticercus* is distinguished from *Cænurus* by its sac (vesicle) having only one pedicle ; it is also always contained in a cyst, which, in some cases, is formed from the compressed textures of the infested animal ; while, in others, it consists of two membranes, viz. one similar to that mentioned, and another *sui generis*, and belonging entirely to the parasite."* The idea of an inner membrane of the cyst, *sui generis*, and belonging entirely to the parasite, is so incompatible with the great characteristic of the entozoa, viz. their independent existence manifested by their state of insulation, that it is marvellous so distinguished a zootomist should have overlooked the connection by vessels existing

* Anatom. and Pathol. Observations, page 91.

between the opaque outer layer and the transparent inner layer of a thickened cyst.

Morbid action progressing in the cyst would naturally lead us to expect the occurrence of morbid changes in the hydatid, and such I have found to be the case. A mere exaltation of the irritation previously noticed would inevitably be productive of a supply of noxious food, which again must furnish a deteriorated material to the growth of the recipient, and the necessary consequence would be a departure from the healthy structure, and even disease unto death of the parasite. The degree of condensation of the cyst to a certain extent governs the deviation from health of its tenant; thus, in a slightly-thickened cyst the textures of the hydatid will be found somewhat opaque only, and the hydatid still living; in a cyst thickened to a higher degree the hydatid will be found dead, with its vesicle burst and shrivelled up. I have in my possession two specimens illustrating these abnormal states, which are faithfully delineated in the accompanying drawings. Fig. 8 exhibits a very elongated *gemmarium*, thickened and opaque from a redundant deposit of granular albuminous matter, and also an opacity of the parietes of the vesicle from the same cause. Fig. 9 shows a much greater extent of disease; the *gemmarium* is completely enveloped in albuminous matter, and from the fimbria radiate fibres of the same material to various parts of the vesicle; the latter is opaque, and dense white patches of calcareous matter are scattered about it. In this example the hydatid was found burst and collapsed. The containing cyst was as firm as cartilage, and its interior was lined with a coarse calcareous and granular matter.

In conclusion, I cannot but observe that there is still much which is interesting and instructive to investigate relating to the *Cysticercus tenuicollis*; and not the least important, I beg leave to suggest, is the generation of their ova (if required for propagation), and the mode by which they or the gemmæ are disseminated.

Let me not omit to acknowledge the valuable services of

Mr. Busk, by whose assistance I have been enabled to speak more confidently of what I have *seen* in my dissections of this entozoon.

EXPLANATION OF PLATE II.

- Fig. 1. A diagram exhibiting the anatomy of *Cysticercus tenuicollis*.—*a*, the crown of hooklets; *b*, the suckorial cups; *c*, the proboscis; *d*, the external coat; *e*, the internal coat; *f*, the gemmarium; *g*, the fimbria of the gemmarium; *h*, the septum between the cavity of the vesicle and the interior of the proboscis, formed of the internal coat.
- Fig. 2. Patch of oval corpuscles; see page 223.
- Fig. 3. Portion of the coronet of hooklets (magnified) in position.—*a*, bifid process of the short hooklets; *b c d*, three hooklets lying in their natural position when expanded; *e*, the manubrium; *f*, the claw; *g*, the conical process.
- Fig. 4. The *Cysticercus* opened longitudinally, exhibiting the *gemmarium*.—*f*, the gemmarium; *g*, the fimbria, a bristle placed under the portion of the gemmarium within the proboscis.
- Fig. 5. The anterior portion of a *Cysticercus* laid open, showing a gemmarium with two or more off-sets, viz. a long central portion, a short lateral off-set, a recurved off-set, and others less distinct.
- Fig. 6. A large *Cysticercus* opened, exhibiting a cluster (I counted nine) of young ones at the upper part of the *gemmarium*, a single one about the middle of its length, and another near its extremity.

- Fig. 7. A *Cysticercus* opened, having a young one, considerably developed, hanging from the upper part of the gemmarium, the inferior portion of the latter floating beside it.
- Fig. 8. A diseased *Cysticercus*, its coats and gemmarium thickened and opaque, particularly the gemmarium.
- Fig. 9. A dead, ruptured and shrivelled *Cysticercus*, taken from a very thickened cyst. Its coats are studded with patches of *calcareous* matter; and the gemmarium is enveloped in granular albumen; also fibres of similar matter are seen radiating in various directions from the fimbriæ to the parietes of the vesicle. See page 236.

Fig: 6.

Fig: 5.

Fig: 7.

Fig. 8.

Fig. 9.