Published in final edited form as:

Biophys Chem. 2007 December ; 131(1-3): 111–114.

Mechanism of Na⁺ binding to thrombin resolved by ultra-rapid kinetics

Stefano Gianni^{a,*}, Ylva Ivarsson^a, Alaji Bah^b, Leslie A. Bush-Pelc^b, and Enrico Di Cera^{b,*}

^a *Istituto di Biologia e Patologia Molecolari del CNR, Dipartimento di Scienze Biochimiche, Università di Roma "La Sapienza", 00185 Roma, Italy*

^b *Department of Biochemistry and Molecular Biophysics, Washington University School of Medicine, Box 8231, St. Louis, MO 63110*

Abstract

The interaction of Na⁺ and K⁺ with proteins is at the basis of numerous processes of biological importance. However, measurement of the kinetic components of the interaction has eluded experimentalists for decades because the rate constants are too fast to resolve with conventional stopped-flow methods. Using a continuous-flow apparatus with a dead time of 50 μs we have been able to resolve the kinetic rate constants and entire mechanism of Na⁺ binding to thrombin, an interaction that is at the basis of the procoagulant and prothrombotic roles of the enzyme in the blood.

Keywords

thrombin; sodium; rapid kinetics; allostery; monovalent cation activation

Enzymes activated by monovalent cations (M⁺s) [1] require Na⁺ or K⁺ for optimal activity and utilize M⁺ as a cofactor or allosteric effector [2,3]. The allosteric effect of Na⁺ on thrombin activity and specificity carries significant physiological relevance [4–6], is at the basis of the procoagulant and prothrombotic roles of the enzyme in the blood and has been studied in great detail both functionally and structurally [7,8]. However, aspects of this physiologically important interaction remain unresolved because Na⁺ binding and dissociation occur on a time scale too fast to measure with conventional stopped-flow techniques [5]. A similar limitation applies to the study of other well characterized M⁺-activated enzymes like Trp synthase [9, 10], pyruvate kinase [11,12], Hsc70 [13], β-galactosidase [14,15] and inosine monophosphate dehydrogenase [16]. Indeed, no protein system characterized to date has offered direct information on the kinetic rate constants for Na⁺ or K⁺ binding/dissociation, therefore leaving a gap in our understanding of how M⁺s in general regulate protein function [3].

To circumvent the limitations of standard stopped-flow systems, we used a continuous-flow instrument equipped with an ultra-rapid mixing device and extended our recent characterization of Na⁺ interaction with thrombin [8] into the μs time scale. To avoid autoproteolysis at the high concentrations (40 μM) needed for the measurements, we inactivated thrombin with the S195A mutation, prepared as described [6,7,17], to selectively knock out the catalytic Ser-195 with minimal structural perturbation [18,19]. As for wild-type thrombin, the fluorescence

*To whom correspondence should be addressed. Tel +39 06-4991-0548, Fax +39 06-444-0062, e-mail: Stefano.gianni@uniroma1.it; Tel +1 (314) 362-4185, Fax +1 (314) 747-5354, e-mail: enrico@wustl.edu.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

increase observed upon Na^+ binding to S195A has an initial rapid phase that cannot be resolved within the dead time (<0.5 ms) of the stopped-flow apparatus, followed by a single exponential slow phase (Figure 1) with a k_{obs} that decreases as $[\text{Na}^+]$ increases (Figure 2). The decrease of k_{obs} is conducive to the existence of a conformational change that precedes Na^+ binding [8]. Of crucial importance is to establish whether the fast phase reports just the binding/dissociation of Na^+ with thrombin or also depends on additional conformational changes. In the former case, k_{obs} should change linearly with $[\text{Na}^+]$. In the latter, a hyperbolic dependence of k_{obs} on $[\text{Na}^+]$ should be observed.

The rapid phase was resolved with the continuous-flow apparatus (Figure 1) and yielded a single exponential time course with a k_{obs} that increases linearly with $[\text{Na}^+]$ (Figure 2). The two-step mechanism of Na^+ binding to thrombin solved by combination of the stopped-slow and continuous-flow measurements is then

SCHEME 1.

This is the mechanism recently proposed by Bah et al. [8], but validated for the binding/dissociation components of Na^+ by direct resolution of the rapid phase. Thrombin exists in equilibrium between two forms, E^* and E , that interconvert with kinetic rate constants k_1 and k_{-1} . Of these forms, only E can interact with Na^+ with a rate constant k_A to populate $E:\text{Na}^+$, that may dissociate into the parent components with a rate constant k_{-A} . The fast phase is due to the E - $E:\text{Na}^+$ interconversion involving Na^+ binding/dissociation, and the slow phase is due to the E - E^* interconversion that precedes Na^+ binding. The exact analytical solution of scheme 1 calls for two eigenvalues that give the k_{obs} for the two exponential transitions reported in Figure 1. Separation of time scales for the E - E^* (slow) and E - $E:\text{Na}^+$ (fast) interconversions simplifies the eigenvalues to

$$\lambda_{\text{slow}} = k_1 + k_{-1} \frac{1}{1 + K_A [\text{Na}^+]} \quad (1)$$

$$\lambda_{\text{fast}} = k_{-A} + k_A [\text{Na}^+] \quad (2)$$

where $K_A = k_A/k_{-A}$. The best-fit parameter values derived from simultaneous analysis of both experimental data sets in Figure 2 are: $k_1 = 67 \pm 7 \text{ s}^{-1}$, $k_{-1} = 69 \pm 6 \text{ s}^{-1}$, $k_A = 56,000 \pm 200 \text{ M}^{-1}\text{s}^{-1}$, $k_{-A} = 4,800 \pm 200 \text{ s}^{-1}$. These values differ somewhat from $k_1 = 224 \pm 4 \text{ s}^{-1}$, $k_{-1} = 34 \pm 4 \text{ s}^{-1}$ and $K_A = k_A/k_{-A} = 57 \pm 4 \text{ M}^{-1}$ reported for wild-type under identical experimental conditions [8], indicating that the S195A mutant has a 5-fold reduced Na^+ affinity and is more stabilized into the E^* form compared to wild-type. These properties do not invalidate the basic mechanism of Na^+ binding uncovered for wild-type and may have actually facilitated measurement of the fast phase. Furthermore, they document an energetic perturbation caused by disruption of the linkage between the side chain of Ser-195 and the water network reaching out into the Na^+ site documented by structural studies [7].

The important conclusion emerged from measurements of the fast phase of Na^+ binding to thrombin is that its k_{obs} increases linearly with $[\text{Na}^+]$. We can therefore rule out additional conformational transitions following Na^+ binding that would have produced curvature in the dependence of k_{obs} vs $[\text{Na}^+]$. The interconversion of E and E* that precedes Na^+ binding is the only conformational transition in scheme 1.

It is of interest to comment on the magnitude of the kinetic rate constants for Na^+ binding and dissociation determined here for the first time for a protein- M^+ interaction. The value of k_A is four orders of magnitude slower than the diffusion controlled limit [20]. Although we lack data on other protein systems for comparison, we can assume that several factors contribute to slow down the kinetics of Na^+ association. Without these factors, Na^+ binding to thrombin would likely be too fast to measure even with the continuous-flow method. Na^+ binds to its site after penetrating a pore defined by the 186- and 220- loops [3,17] whose size would not admit a fully hydrated M^+ . In this regard, the interaction of Na^+ with thrombin has features in common with the permeation and selectivity of ion channels [2,3,21]. Dehydration of Na^+ required to enter the pore is expected to slow down the kinetics of association significantly [21,22]. Once inside the pore, Na^+ is re-hydrated into its coordination shell that involves four water molecules and the backbone oxygen atoms of Lys-224 and Arg-221a [7]. That in turn organizes a network of water molecules that spans the interior of the enzyme for over 15 Å up to the catalytic Ser-195 [7]. Ordering of this network of water molecules may require a longer time scale than the simple process of penetrating the pore, thereby contributing to the drastic reduction in the rate of binding k_A measured experimentally. The role of dehydration/re-hydration postulated here for the kinetics of Na^+ binding to thrombin can be tested experimentally by studying the kinetics of interaction with other M^+ s that differ in hydration energy and ionic radius. Unfortunately, measurements carried out with K^+ , that binds to thrombin with lower affinity compared to Na^+ [17], have so far failed to yield large enough spectral changes with the continuous-flow method. Combination of dehydration, penetration of the pore, and re-hydration with organization of the water network around the Na^+ site could account for the significant slow down in the kinetics of Na^+ association.

The results presented in this study offer valuable new information on the physico-chemical components linked to Na^+ binding to thrombin. Future work on this and other M^+ -activated enzymes will establish if general features indeed exist for the interaction of M^+ s with proteins.

Acknowledgements

We are indebted to Prof. Maurizio Brunori for valuable suggestions. This work was supported in part by NIH research grants HL49413, HL58141 and HL73813 (to E.D.C.). Y.I. is supported by a grant from the Wenner-Gren Foundations (Sweden).

References

1. Suelter CH. Enzymes activated by monovalent cations. *Science* 1970;168:789–795. [PubMed: 5444055]
2. Di Cera E. A structural perspective on enzymes activated by monovalent cations. *J Biol Chem* 2006;281:1305–1308. [PubMed: 16267046]
3. Page MJ, Di Cera E. Role of Na^+ and K^+ in enzyme function. *Physiol Rev* 2006;86:1049–1092. [PubMed: 17015484]
4. Di Cera E. Thrombin interactions. *Chest* 2003;124:11S–17S. [PubMed: 12970119]
5. Di Cera E, Page MJ, Bah A, Bush-Pelc LA, Garvey LC. Thrombin allostery. *Phys Chem Chem Phys* 2007;9:1292–1306.
6. Dang OD, Vindigni A, Di Cera E. An allosteric switch controls the procoagulant and anticoagulant activities of thrombin. *Proc Natl Acad Sci U S A* 1995;92:5977–5981. [PubMed: 7597064]

7. Pineda AO, Carrell CJ, Bush LA, Prasad S, Caccia S, Chen ZW, Mathews FS, Di Cera E. Molecular dissection of Na⁺ binding to thrombin. *J Biol Chem* 2004;279:31842–31853. [PubMed: 15152000]
8. Bah A, Garvey LC, Ge J, Di Cera E. Rapid kinetics of Na⁺ binding to thrombin. *J Biol Chem* 2006;281:40049–40056. [PubMed: 17074754]
9. Woehl E, Dunn MF. Mechanisms of monovalent cation action in enzyme catalysis: the first stage of the tryptophan synthase beta-reaction. *Biochemistry* 1999;38:7118–7130. [PubMed: 10353822]
10. Woehl E, Dunn MF. Mechanisms of monovalent cation action in enzyme catalysis: the tryptophan synthase alpha-, beta-, and alpha beta-reactions. *Biochemistry* 1999;38:7131–7141. [PubMed: 10353823]
11. Mesecar AD, Nowak T. Metal-ion-mediated allosteric triggering of yeast pyruvate kinase. 2. A multidimensional thermodynamic linked-function analysis. *Biochemistry* 1997;36:6803–6813. [PubMed: 9184163]
12. Mesecar AD, Nowak T. Metal-ion-mediated allosteric triggering of yeast pyruvate kinase. 1. A multidimensional kinetic linked-function analysis. *Biochemistry* 1997;36:6792–6802. [PubMed: 9184162]
13. O'Brien MC, McKay DB. How potassium affects the activity of the molecular chaperone Hsc70. I. Potassium is required for optimal ATPase activity. *J Biol Chem* 1995;270:2247–2250. [PubMed: 7836457]
14. Xu J, McRae MA, Harron S, Rob B, Huber RE. A study of the relationships of interactions between Asp-201, Na⁺ or K⁺, and galactosyl C6 hydroxyl and their effects on binding and reactivity of beta-galactosidase. *Biochem Cell Biol* 2004;82:275–284. [PubMed: 15060622]
15. Neville MC, Ling GN. Synergistic activation of beta-galactosidase by Na and Cs. *Arch Biochem Biophys* 1967;118:596–610. [PubMed: 4860414]
16. Digits JA, Hedstrom L. Kinetic mechanism of *Trichomonas foetus* inosine 5'-monophosphate dehydrogenase. *Biochemistry* 1999;38:2295–2306. [PubMed: 10029522]
17. Prasad S, Wright KJ, Roy DB, Bush LA, Cantwell AM, Di Cera E. Redesigning the monovalent cation specificity of an enzyme. *Proc Natl Acad Sci U S A* 2003;100:13785–13790. [PubMed: 14612565]
18. Krem MM, Di Cera E. Dissecting substrate recognition by thrombin using the inactive mutant S195A. *Biophys Chem* 2003;100:315–323. [PubMed: 12646374]
19. Bah A, Chen Z, Bush-Pelc LA, Mathews FS, Di Cera E. Crystal structures of murine thrombin in complex with the extracellular fragments of murine protease-activated receptors PAR3 and PAR4. *Proc Natl Acad Sci USA* 2007;104:11603–11608. [PubMed: 17606903]
20. van Holde KE. A hypothesis concerning diffusion-limited protein-ligand interactions. *Biophys Chem* 2002;101–102:249–254.
21. Roux B, MacKinnon R. The cavity and pore helices in the KcsA K⁺ channel: electrostatic stabilization of monovalent cations. *Science* 1999;285:100–102. [PubMed: 10390357]
22. Eisenman G, Dani JA. An introduction to molecular architecture and permeability of ion channels. *Annu Rev Biophys Chem* 1987;16:205–226. [PubMed: 2439095]
23. Gianni S, Walma T, Arcovito A, Calosci N, Bellelli A, Engstrom A, Travaglini-Allocatelli C, Brunori M, Jemth P, Vuister GW. Demonstration of long-range interactions in a PDZ domain by NMR, kinetics, and protein engineering. *Structure* 2006;14:1801–1809. [PubMed: 17161370]
24. Shastry MC, Roder H. Evidence for barrier-limited protein folding kinetics on the microsecond time scale. *Nat Struct Biol* 1998;5:385–392. [PubMed: 9587001]
25. Jemth P, Gianni S, Day R, Li B, Johnson CM, Daggett V, Fersht AR. Demonstration of a low-energy on-pathway intermediate in a fast-folding protein by kinetics, protein engineering, and simulation. *Proc Natl Acad Sci U S A* 2004;101:6450–6455. [PubMed: 15096617]
26. Peterman BF. Measurement of the dead time of a fluorescence stopped-flow instrument. *Anal Biochem* 1979;93:442–444. [PubMed: 464271]

Figure 1.

A,B (A) Kinetic traces of Na⁺ binding to human thrombin in the 0–250 ms time scale. Shown are the traces obtained at 50 and 100 mM Na⁺ with the stopped-flow method using an Applied Photophysics SX20 spectrometer, with an excitation of 280 nm and a cutoff filter at 305 nm [8]. Traces are averages of three determinations. Binding of Na⁺ obeys a two-step mechanism, with a fast phase completed within the dead time (<0.5 ms) of the spectrometer, followed by a single-exponential slow phase. The k_{obs} for the slow phase decreases with increasing [Na⁺] (Figure 2A). (B) Kinetic traces of Na⁺ binding to human thrombin in the 0–700 μs time scale. Shown are the traces obtained at 36 and 91 mM Na⁺ with the continuous-flow method as single determinations with an exposure time of 3 s. We built an ultra-rapid mixer [23] of similar design and methodology to that described by Shastry and Roder [24,25]. The flow cell was purchased from Hellma (Germany). The mixed sample was illuminated with an A1010B Mercury-Xenon lamp (PTI, UK) at 280 nm, using a Model 101 6 Monochromator (PTI, UK). Fluorescence was recorded with a Micromax CCD camera (Princeton Instrument, USA), with a typical exposure time of 1–3 s and employing different emission glass filters. An original, pneumatically driven, loading syringes unit was designed. Data were recorded at 3 bars, leading to a linear velocity in the flow cell of about 16 m/s. Experimental conditions were identical to those used in the stopped-flow experiments. The dead time of the instrument was determined by fluorescence using the quenching of N-acetyl-tryptophanamide (NATA) by N-bromo-succinamide (NBS) [26]. The pseudo-first-order quenching reaction of NATA by NBS was measured at NBS concentrations from 3.5 to 33 mM. The observed fluorescence traces obtained at different NBS concentrations yielded single exponential time courses extrapolating to a common point near $f_{\text{rel}} = 1$, the expected initial fluorescence of NATA in the absence of quencher. The time delay from this point to the first data point that falls on the fitted exponential provided an estimate of the dead time, 40–50 μs , of the instrument. Binding of Na⁺ in the 0–700 μs time scale obeys a single-exponential phase with a k_{obs} increasing linearly with [Na⁺] (Figure 2B). This resolves the fast phase detected with the stopped-flow method and shown in (A). Experimental conditions for the two methods are: 5 mM Tris, 0.1% PEG8000, pH 8.0 at 25 °C. The thrombin concentration was 50 nM for the stopped-flow measurements and 40 μM for the continuous-flow measurements. The [Na⁺], as indicated, was changed by keeping the ionic strength constant at 400 mM with choline chloride. Continuous lines were drawn using the expression $a + b\exp(-k_{\text{obs}}t)$ with best-fit parameter values: (A) $a=8.944\pm0.001$ V, $b=-0.10\pm0.01$ V, $k_{\text{obs}}=111\pm9$ s⁻¹ ([Na⁺]=50 mM); $a=9.427\pm0.001$ V, $b=-0.29\pm0.01$ V, $k_{\text{obs}}=96\pm6$ s⁻¹ ([Na⁺]=100 mM). (B) $a=0.417\pm0.002$, $b=-0.072\pm0.02$, $k_{\text{obs}}=7\pm2$ ms⁻¹ ([Na⁺]=36 mM); $a=0.477\pm0.002$, $b=-0.10\pm0.03$, $k_{\text{obs}}=10\pm2$ ms⁻¹ ([Na⁺]=91 mM). All data were collected at least in duplicate.

Figure 2.

A,B Values of k_{obs} vs $[\text{Na}^+]$ for the slow and fast phases of fluorescence change due to Na^+ binding to thrombin shown in Figure 1. Shown are the results pertaining to the stopped-flow (**A**) and continuous-flow (**B**) measurements. Note the different time scale between the two panels. Experimental conditions are given in the legend to Figure 1. Continuous lines were drawn according to eqs 1 and 2 in the text, with best-fit parameter values: $k_1=67\pm7\text{ s}^{-1}$, $k_{-1}=69\pm6\text{ s}^{-1}$, $k_A=56,000\pm200\text{ M}^{-1}\text{s}^{-1}$, $k_{-A}=4,800\pm200\text{ s}^{-1}$.