ON
DISLOCATIONS
OF
THE VERTEBRÆ.

BY
W. LAWRENCE, F.R.S.

SURGEON TO ST. BARTHOLOMEW'S HOSPITAL, ETC.

Read June 12, 1827.

COMPLETE dislocations of the vertebræ without fracture, whether produced by accident or disease, are so uncommon, if we except the first and second bones of the neck, that the possibility of the occurrence has been doubted, and even denied by some of the highest authorities. In the “*Traité des Maladies Chirurgicales*”, which contains the valuable results of his long experience, Baron Boyer points out the circumstances which prevent the displacement of the bodies of the vertebræ, and, without expressly denying it, appears clearly to disbelieve its possibility*. He asks the question, whether the bodies of the vertebræ can be luxated? alludes

* Tom. IV. Chap. IV. Article 4.

to the kind of accidents, under which such displacement has been considered probable, and then, having shewn the powerful obstacles to it, concludes by observing, that if we examine the cases in which dislocation has been supposed to have occurred, we shall find invariably that the posterior laminæ of the vertebræ have been broken; that in many instances they have been comminuted, and further that when the body of a vertebra has undergone any displacement, the laceration of the fibro-cartilage has been almost always accompanied by detachment of some portion of the bony substance.

Delpech represents it as a point proved by observation, that the bodies of the vertebræ cannot be dislocated*.

Sir A. Cooper† says, that “if luxation of the spine ever does happen, it is an injury which is extremely rare, as in the numerous instances which I have seen of violence done to the spine, I have never witnessed a separation of one vertebra from another through the intervertebral substance, without fracture of the articular processes; or, if those processes remain unbroken, without a fracture through the bodies of the vertebræ.”

* *Précis Élémentaire des Maladies réputées Chirurgicales.* Tom. III. p. 42.

† *Treatise on Dislocations and on Fractures of the Joints.* 1st edition, p. 539.

Other surgeons, however, have affirmed that the vertebræ may be luxated. After pointing out, in his treatise on diseases of the joints, the circumstances which render this occurrence very improbable, Rust affirms that even the lumbar and dorsal vertebræ may be dislocated, and refers to instances recorded by various writers*. He mentions the following as a case of dislocation in the cervical region of the spine, successfully replaced by himself. "The injury was produced by a severe fall on the head. The neck was completely bent to the right side, the upper extremities already paralysed, and repeated attacks of convulsions and hiccough came on. Replacement was immediately attempted, and succeeded. I made the patient sit on the ground, and had the head drawn straight upwards by a strong assistant. The patient got well under the employment of cold locally."†

Mr. Bell mentions a case of complete dislocation between the last dorsal and first lumbar vertebra, with entire division of the spinal cord. A small portion of bone was broken off. He gives two views of the parts‡.

The greater mobility of the individual bones, the comparative smallness of their bodies, and the ob-

* *Arthro-kakologie*, p. 71.

† *Medicinisch-chirurgische Zeitung*, 1813, Vol. III. p. 127.

‡ *On Injuries to the Spine and Thigh-bone*, p. 25. Plate II. fig. 2 and 3.

liquity of the articular processes, point out the cervical vertebræ as those most likely to be luxated; at the same time the form of the neck, and its connexion with the head, are favourable to the application of such violence as may cause luxation. Hence not only does dislocation of the atlas occasionally occur, but we have also instances of luxated articular processes in the case of the five inferior cervical vertebræ. Baron Boyer even considers that this may happen without external violence, and that the inferior articular process of a cervical vertebra may be carried in front of the superior articular process of the vertebra below it, by a sudden and forcible rotation of the head and neck towards the opposite side. He says that "Desault mentioned in his lectures the case of an advocate, who met with this kind of dislocation, by turning his head suddenly round to see who was coming in at a door situated behind his seat. Chopart also shewed us a young man, 24 years old, in whom a similar accident had occurred, in consequence of an extreme rotation of the head, leaving the head permanently inclined upon the left shoulder."*

Baron Boyer mentions another instance, which Desault used to relate in his lectures, of a child, eight or nine years old, who, in turning heels over head on a bed, luxated the right lower articular process of a cervical vertebra. The head was turned

* P. 114.

towards the left shoulder, and so firmly fixed that it could not be brought into its natural position, even by the employment of considerable force. When the proposal of attempting reduction was made, Desault decidedly discountenanced it*. Another case of a child terminated fatally, under the employment of efforts at reduction; and, when the body was examined, one of the inferior articular processes of a cervical vertebra was found dislocated forwards†. It does not appear that Baron Boyer had ever ascertained the existence of this dislocation by actual examination, and it is even doubtful whether he had ever seen a case at all. Further, he admits that no example can be adduced of both articular processes being dislocated at the same time.

Sir A. Cooper‡ says that he has never seen a dislocation of the cervical vertebræ: he does not deny the possibility of the occurrence, but seems hardly to believe it.

In the anatomical museum at St. Bartholomew's Hospital there are specimens of luxated cervical vertebræ. In one of these the right inferior articular process of the fifth vertebra is dislocated forwards. The portion of the vertebral column above the seat of the injury is twisted to the left, and the

* P. 117. † P. 118. ‡ On Dislocations, p. 539.

body of the fifth, having been partially displaced, projects beyond that of the sixth vertebra. This displacement could not have been effected without laceration, or considerable injury of the fibro-cartilage. The upper and anterior part of the body, both of the sixth and seventh vertebræ, has been slightly fractured on the left side, without detachment of the broken portion. It appears as if a small piece of the edge had been broken off by the twisting of the spine to the left. As the specimen, which is dried and inclosed in a bottle, has not been completely cleaned of the soft parts, the exact nature and extent of the fracture cannot be ascertained, nor can it be pronounced with perfect certainty, that there is no fracture of the processes. For the same reason we cannot now make out the precise degree and kind of injury sustained by the fibro-cartilage.

In another, the inferior articular processes of the fifth cervical vertebra are partially separated from those of the sixth, having been drawn upwards; but they have not been thrown forwards. The bodies of the same vertebræ are partially separated behind, but they preserve their natural level and relative position in front. The connexions of the two vertebræ must have been torn by a force applied behind, bending the neck powerfully forwards, and the bones must have subsequently resumed their natural position, so that there would have been no

pressure on the spinal marrow. This kind of injury has been called by Mr. Bell*, diastasis or subluxation.

A third specimen exhibits a dislocation of the sixth upon the seventh cervical vertebra. The inferior articular processes of the sixth are completely displaced forwards, and its body projects over that of the seventh. The spinous processes have been sawn out, so that it is uncertain whether they were injured or no. This may be deemed a case of complete dislocation; but there is an appearance of fracture near the basis of the articular process, the exact nature of which cannot be ascertained in the present state of the specimen. Mr. Stanley considers that this fissure was made by the saw in removing the spinous processes.

* Observations on Injuries of the Spine and Thigh-bone, p. 9, fig. III. plate III. Diastasis is called by Mr. Bell, "separation of the vertebræ by violence, where they resume their relations again", p. 9.—Speaking of subluxation, he says, "the accident happens to young people, and from the operation of a force, which, in advanced years, would fracture the bodies of the vertebræ. A weight on the head and shoulders overpowering them, and bending them double, the articulating processes of the upper lumbar vertebræ are burst from their connexions; if they again fall into their places the case is diastasis; but sometimes their edges meet, then it is subluxation; they are not restored to their natural position." P. 11.

Mr. Bell relates a case at p. 9, to exemplify diastasis occurring between the last cervical and first dorsal vertebra. The parts are represented in pl. III. fig. III.; in the explanation of the figure, he calls it a specimen of subluxation.

The following case, which was recently under my care at St. Bartholomew's Hospital, shows that complete dislocation, both of the articular processes and body, without fracture, may occur in the cervical region of the spine. I submit at the same time to the examination of the Society, the dislocated bones, and two drawings of them.

C A S E.

Charles Butcher, 22 years old, a robust young man, in perfect health, was brought to the hospital at five o'clock in the afternoon of January 8th. He was carrying a heavy barrel on the back of his head and neck, slipped in descending some steps, and fell on the buttocks, the burthen resting on the head and upper part of the neck. He was immediately deprived of sensibility in the trunk and limbs, and of all power over the voluntary muscles of those parts. When he was brought to the hospital, he was completely insensible and incapable of voluntary motion below the neck. The functions of the brain were not disturbed. Respiration was apparently performed merely by the diaphragm: its descent enlarged the chest, while the elasticity of the thoracic and abdominal parietes restoring the parts, which had been previously displaced by the diaphragm, seemed to be the only agent of expiration. The chest was motionless, except towards its inferior margin, and the abdominal muscles were

soft. The pulse was weak and slow, and the body cold; the penis was in a state of permanent erection. No irregularity in the line of the spinous processes could be discovered by accurate examination. He was placed on the back in bed, with the head carefully supported. In four or five hours the pulse had become full and strong, and the heat of the body was greater than natural; respiration was rather hurried. Venesection to sixteen ounces, a dose of Calomel and Jalap. Four ounces of urine were drawn off by the catheter.

Jan. 9th.—He dozed in the course of the night, but did not sleep. There is pain in the lower part of the neck. He can move the arms very slightly, and has a little feeling in the front and upper part of the chest. Dark offensive stools have been passed involuntarily. When the catheter was introduced in the morning, only a table-spoonful of urine came away: six ounces of high-coloured urine were drawn off in the evening. Respiration is slower, the pulse full, heat natural. Saline draught every six hours.

10th.—Feels better this morning, having slept three or four times in the night, for about an hour each time. Has experienced a tingling sensation in the hands, and is sensible to impressions on the upper part of the arms and thighs: stools have passed involuntarily; he complains of distention of the bladder; eighteen ounces of high-coloured urine

were drawn off, and deposited a dark-brown sediment. The priapism continues. Four ounces of urine drawn off in the evening.

11th.—He has not slept during the night, and is worse this morning. Respiration becomes difficult, so that speaking requires a painful effort; and the countenance expresses distress and anxiety. The tongue is light-brown, and dry in the middle. Fæces have passed involuntarily, and six ounces of urine, lighter-coloured than that of yesterday, have been drawn off. Breathing became more laborious in the evening, and the countenance extremely anxious; the pulse weak and low, surface of the body cold. He expired at one in the morning of the 12th, from increasing embarrassment in respiration.

Examination.—No displacement nor irregularity could be discovered by external examination, when the body was laid on the face. After cutting away the muscles from the back of the spine, the cartilaginous surfaces of the superior articular processes of the fifth cervical vertebra came into view: they were exposed in consequence of the inferior processes of the fourth vertebra having been completely dislocated forwards, and remaining fixed in their unnatural position. The yellow ligaments connecting the laminæ of the two vertebræ (ligamenta subflava) were torn through, and the bifid apex of the fourth spinous process lay in close con-

tact with the basis of the fifth. On the front of the column an unusual projection was observed, but the anterior longitudinal ligamentous expansion was entire. The body of the fourth was completely detached from that of the fifth vertebra, the connecting fibro-cartilage being torn through, and the body of the former projecting by its whole depth in front of the latter. In consequence of this displacement, the antero-posterior diameter of the vertebral canal is lessened by about one-third. The section of the bone was not made till some days after death, so that the recent state of the spinal marrow could not be estimated.

The nature of the accident in this case sufficiently accounts for the effect produced. The head and four upper vertebræ, were driven forwards by the heavy burthen which rested on them, while the fall on the buttocks forcibly propelled the lower part of the column in the opposite direction. As the case was simple dislocation, unattended with any other effect, except such pressure on the spinal cord as sufficed to destroy sensibility, and the power of motion below the seat of injury, it affords us the opportunity of observing how, and in what time, death occurs under such circumstances. It is caused by the paralysis of a large portion of the muscles concerned in respiration, and by the consequently imperfect mode in which the mechanical phænomena of that function are executed.

The intercostal and abdominal muscles, and nearly all those of the chest, which are capable of assisting in the motions of inspiration and expiration, were completely palsied in this case, and the diaphragm alone remained capable of action, the pressure on the spinal cord being below the origin of the phrenic nerve. Breathing was at first so well executed by this single power, that, on a cursory view of the patient, there was nothing alarming in his situation: the inadequacy of the diaphragm, however, to the sole performance of respiration soon became apparent, and the patient sunk on the fourth day under the continually increasing difficulty.

In a patient under my care at St. Bartholomew's during the last winter, with fracture and displacement of the spine at the seventh cervical vertebra, and paralysis of the parts below, death occurred on the tenth day. Sir A. Cooper observes, that "Death ensues in these cases in from three to seven days, as it (the injury) is seated in the fifth, sixth, or seventh vertebra. I have scarcely known the subject of this injury to live beyond a week, and but rarely to die on the second day, although they sometimes do if the fifth cervical vertebra has sustained the injury." *

* On Dislocations, pp. 555, 556.

Dislocation of the Atlas and Axis from Disease, with consequent Anchylosis.

Although accidental luxation of the two first cervical vertebræ is immediately fatal from pressure on the commencement of the spinal cord *,

* A case, which occurred to Mr. Cline, and is related by Sir A. Cooper, forms an apparent exception to this statement. A boy, who had met with a severe fall on the head, and had been subsequently obliged to observe great caution in all the motions of that part, steadying it and supporting it with the hands, died twelve months after the accident. Mr. Cline found "the first vertebra of the neck broken across, so that the dentiform process of the second vertebra had so far lost its support, that under the different inclinations of the head, it required great care to prevent the spinal marrow from being compressed by it, and as he could not depend upon the actions of the muscles of the neck, he therefore used his hands to support the head during different motions and positions." On Dislocations, pp. 549, 550.

If the case recorded by Ehrlich was, as he supposes it to have been, a dislocation of the atlas, it proves that such an accident is not only not necessarily fatal, but that it even admits of complete remedy. A young man of sixteen fell in carrying a sack of flour upstairs, and his head was forcibly bent forwards by the burthen. He was found senseless, with livid countenance, prominent eyes, tongue hanging out of the mouth, respiration slow and interrupted, and pulse scarcely sensible. The limbs were motionless and apparently paralysed, urine and fæces had passed involuntarily. The head was inclined to the right shoulder and had lost its firmness, so that it fell by its own weight from side to side when not supported. The articular process of the second vertebra projected on the left side. Ehrlich considered it a case of dislocation with pressure on the spinal cord, and caused extension of the head to be made, while he endeavoured to force back the atlas, and to bring the second vertebra forwards. After some at-

these bones may be slowly displaced, in consequence of disease in their articulations, without affecting the functions of that important part; and even, after disease is at an end, may become permanently fixed in their new positions by ankylosis. A remarkable specimen of this kind was given to me by Mr. Wigan, a member of this Society; and I lay it before the Society for their examination, as the nature and extent of the displacement could hardly be credited without actual observation. Mr. Wigan, who attended the patient throughout, had promised to furnish me with his notes of the case; but he could not find them before he left England; I can therefore only mention the few leading circumstances in the history, which Mr. Wigan communicated to me verbally when he brought me the parts in their recent state.

CASE.

At the age of five or seven, a child became the subject of an illness supposed to be hydrocephalus.

tempts, the replacement was effected with a snap. The head now became steady and the arms moved, but the patient remained insensible, with dilated pupils; the respiration and pulse were improved. Consciousness returned on the next day. Great swelling of the neck came on from ecchymosis and subsided in a short time. He was much recovered in eight days, and soon after got quite well. *Chirurgische Beobachtungen*, Vol. II. No. 25.

lus. After it had existed for some time, a swelling took place in the side of the neck, obviously containing fluid, and this slowly increased to a considerable magnitude, so that its contents must have amounted to several ounces. Pressure on this swelling affected the brain, producing a state of coma. It was supposed at this time that the fluid in the neck communicated with that suspected to exist in the ventricles of the brain. The child used to move the head cautiously and slowly, supporting it with the hands at the sides. After a long continuance the symptoms slowly subsided, and at last entirely disappeared, together with the swelling of the neck. At no period of the complaint was there any interruption or diminution of sensation or voluntary motion, and the recovery of health and activity was complete, the child being able to walk and run, and engage in the active sports suited to its age. There was nothing to attract particular notice in the position of the head. After some time had elapsed, disease came on in the lumbar vertebræ, attended with bending forwards of the spine and the formation of a large lumbar abscess. In consequence of this affection the child died at the age of twelve.

Examination.—The head was examined during the hottest part of last summer, and the brain had become so soft, that the changes produced in it by disease could not be ascertained. Mr. Wigan brought to me the basis of the skull, in which we

were surprised at observing a considerable bony prominence standing up in the right side and front of the foramen magnum.

The projection in question was smoothly covered by the dura mater, and it was soon apparent that it must be the dentiform process of the second vertebra. When the soft parts had been completely removed by maceration, I found an extensive displacement of the occiput, atlas, and axis, and a firm consolidation of these bones in their new relative positions by the complete bony ankylosis of several articulations. The atlas is partially dislocated towards the left, and at the same time thrown a little forwards and upwards; hence the right and posterior part of its bony ring intercepts a considerable portion of the spinal canal. The middle anterior protuberance now corresponds to the left side of the basilar process; the extremity of the left transverse process projects three quarters of an inch beyond those of the two following vertebræ, while the right transverse processes of those vertebræ project one quarter of an inch beyond the corresponding one of the atlas. A considerable part of the right side of this bone has been destroyed by absorption: that is, the surfaces by which it is articulated to the occiput and atlas, a part of the transverse process, and that groove on which the right vertebral artery rested. The axis is completely dislocated from the atlas and occiput to the right, so that its left portion in-

tercepts about one third of the spinal canal, and the dentiform process projects by its whole length, into the cavity of the skull at the anterior part of the foramen magnum, close to the right anterior condyloid foramen.

In the natural position of the parts, the apex of this process is a little below the level of the occipito-atlantal articulation ; here it is an inch above the same level, while its projection into the cavity of the skull is between five and six-eighths of an inch. The lateral displacement is no less extensive ; the measurement from the left anterior condyloid foramen to the middle of the basis of the dentiform process being seven-eighths of an inch, while the distance from the right foramen to the same point is only two-eighths. We cannot but be astonished at finding that the immediate pressure of this bony projection on the under surface of the medulla oblongata caused no paralytic affection, even when we allow for the very gradual manner in which it must have taken place. The course of the right lingual nerve and vertebral artery must have been greatly altered.

The bodies of the second and third vertebræ are displaced towards the right, so that a line drawn along their middle and continued upwards, would strike the right margin of the basilar process ; and the small tubercle on the anterior arch of the atlas corresponds to the left side of those bodies.

The right occipital condyle, the remains of the right transverse process of the atlas, the inferior articular plane of the axis, and the right side of the basis of the dentiform process are anchylosed, so as to form one solid bony mass.

The left occipital condyle is partially anchylosed with the atlas; the left articular plane of the axis, its transverse process, and that of the third vertebra, are all consolidated with the partial anchylosis just mentioned. The articular processes of the axis and the third vertebra are anchylosed, the union extending on the left side as far as the bases of the spinous processes. The bodies of the axis and third vertebra are not anchylosed.

The preternatural bony connexions just enumerated are perfect, equalling in solidity the natural bony structure, so that the limits of the anchylosed bones are confounded. The bony texture is quite natural; there is no roughness of surface, no diminution of solidity, nor any appearance of caries.

The dimensions of the vertebral canal at its commencement are greatly reduced by the extensive displacement of the atlas and axis. The antero-posterior diameter of the foramen magnum is in this case one inch and a half; the greatest measurement from side to side is one inch and a quarter. The diameter of the ring of the third vertebra,

from side to side, is seven-eighths of an inch, from front to back six-eighths. The measurement of the canal, between the displaced portions of the atlas and axis, is half an inch from before backwards, and five-eighths of an inch from side to side.

The history and examination of this case clearly show, that the opinion originally entertained of its being hydrocephalus was altogether erroneous, that the primary disease was an affection of the articulations, the fluctuating tumour in the neck being a chronic abscess consequent on that affection, while the spontaneous disappearance of the tumour is referrible to the cessation of the irritation which caused it. The swelling bore the same relation to the vertebral disease that lumbar abscess does to disease in the lumbar region of the spine.

OBSERVATIONS SUGGESTED BY THE FOREGOING CASE.

Although disease of the upper cervical vertebræ, similar to that which had occurred in the preceding case, is not uncommon, the symptoms of the affection have not been carefully attended to, nor its history clearly made out; so that it is hardly mentioned by any English writer, while by most the subject is altogether unnoticed. It has, no doubt, been overlooked, in consequence of the affected parts being out of sight and reach, and the

symptoms being therefore somewhat obscure, and liable, on superficial observation, to be confounded with those of other affections. Professor Rust, of Vienna, has given the best account of the disease. He first noticed it shortly in the *Salzburg Medico-Chirurgical Journal**, and afterwards described it at greater length in his work on diseases of the joints†, where he mentions that he had seen thirteen cases, in all of which the symptoms were nearly alike.—Pain in the neck, becoming more severe at night, or in swallowing a large mouthful, or drawing a deep breath, is the first symptom. This pain affects one side of the neck, especially when the head is moved towards the shoulder; it extends from the larynx towards the nape, and often to the scapula of the pained side. No external alteration is perceptible; but firm pressure on the region of the first and second vertebræ produces considerable pain, and thus points out the seat of disease. The difficulty of swallowing and breathing, and hoarseness increase, alternating with pain in the neck, which seems to fix about the back of the head, and becomes intolerable on moving that part. The head sinks towards one shoulder, the face being turned a little down, for

* 1813, V. III. p. 108, 109.

† *Arthrokakologie*, 1817. § 111—119.

The short but clear notice of the subject by Roche and Sanson in their "*Nouveaux Elémens de Pathologie Medico-Chirurgicale*. Tom. II. p. 255—258", seems to have been derived chiefly from the observations of Rust.

in general the articulations are affected on one side only, and that was the left in seven out of nine examinations after death. If both sides are affected the head will incline directly forwards. In this state things continue for several weeks or months; and before worse symptoms come on, there is often apparent improvement, freer motion, and more natural situation of the head. But the uneasiness in speaking and swallowing returns, the pain becomes more severe and extensive, the head falls a little backwards and sinks towards the opposite side. The patient feels as if the head were too heavy, and he carefully supports it with his hands, when he moves from the sitting to the lying position, or vice versâ. This may be considered a pathognomonic symptom of the affection. Another symptom, which at this period shows the true nature of the disease, is a peculiar expression of pain in the countenance, which, combined with the position and stiffness of the head, constitutes so characteristic an assemblage of appearances, that it is enough to have seen it once, in order to recognize it again immediately. This look of the patient, which Rust has endeavoured to represent in an engraving, consists especially in a general alteration of the features, with heavy motion of the eyes, and a dull melancholy expression of internal painful sensations. More active indications of severe suffering are observed whenever the head is moved.

In the further progress of the case, noise in the head, deafness, giddiness, cramps and convulsions, partial paralysis, particularly of the upper limbs, loss of voice, purulent expectoration, and hectic symptoms supervene. Generally, no external change is observable, either in the neck or in the nape; and Rust observed, in one case only, swelling of the affected side, which broke and left fistulous ulcers. But the slightest pressure in the region of the three upper vertebræ is acutely painful, and sometimes in the advanced period of the disease a grating of rough surfaces is distinctly perceptible when the head is turned. The patient may continue for months in this helpless and painful state, and then dies, either from exhaustion and debility, or, which is more frequent, suddenly and unexpectedly.

Such is Rust's description of this formidable disease, which we can have no hesitation in regarding as originally ulceration of the cartilages, proceeding to destruction of the ligaments, and caries of the bones, and extending in various shapes and degrees to the neighbouring important parts. I have not met with any instance illustrating the condition of the disease, while confined to the articular surfaces. Reil shortly mentions the case of a young man who was attacked with what was deemed inflammatory sore throat, the head being bent forwards and towards the right.

The sore throat was removed, but the neck became more bent. Hectic fever came on, and the patient died with paraplegia on the seventeenth day.

The occipital condyles, the bodies of all the cervical, and of the four upper dorsal vertebræ were found in a softened state, while the articular surfaces of the condyles and the two first vertebræ were partly destroyed*. Here then, in addition to the ulceration of the cartilages, there seems to have been extensive inflammation of the bodies of the vertebræ.

Mr. Bell mentions two instances in which the connexions of the occiput and atlas, and those of the atlas and axis had become so much weakened by disease, that they gave way under a movement of the head forwards, causing sudden luxation of the dentiform process, and immediate death by pressure on the spinal cord. A figure is given representing the parts in one of these cases, but the condition of the articular surfaces is not described †. The appearances of the diseased parts in a more advanced period are described by Rust. "I have always", says he, "found, in larger or smaller quantity, matter between the pharynx and vertebral column; it has sometimes been of a

* Fieber-lehre. 2d Edit. Vol. II. p. 356.

† Observations on Injuries of the Spine, &c. p. 27. pl. III. fig. 1.

good kind, sometimes sanious. The pharynx is more or less ulcerated (*excoriirt*), and the periosteum and ligaments of the two first vertebræ are destroyed. I have always found the original seat of disease in the articulations of the occiput and atlas, and of the atlas and axis; caries has attacked the occipital condyles, more frequently the left than the right, the atlas and the dentiform process. Where death has come on suddenly, the diseased dentiform process has been entirely broken off, or blood has been effused from ulceration of the vertebral artery, or pus has been found in the chest. The neighbouring important organs may be more or less seriously affected. Serous effusion is sometimes found between the dura mater and the arachnoidea, and in the ventricles. In the vicinity of the foramen magnum, where the space for lodging the spinal cord is sometimes much contracted, the dura mater is found livid, ulcerated and sloughy, and the neighbouring parts of the brain and medulla more or less altered in appearance. I have placed several preparations illustrating this affection in the Pathological Museum of the Vienna General Hospital”*.

In some cases the disease appears to be confined

* Arthro-kakologie, § 117.

In the fifth and sixth plates, figures are given of the occiput and three first vertebræ of a boy twelve years old, who died of the disease. The articular surfaces and other parts of the bones are carious.

to the articular surfaces; after destroying the cartilages and ligaments, and producing more or less extensive displacement of the occiput and two first vertebræ, it stops, and the process of reparation commences, ending in bony ankylosis, by which the displaced parts are firmly fixed in their new position. Yet, in these more favourable instances, there is great suffering during the inflammatory stage; the extension of active disturbance to the dura mater, and the neighbouring contents of the cranium and vertebral canal, may cause dangerous and alarming symptoms, while chronic abscess may be formed externally, as in the case which I have related.

The size of the foramen magnum and the dimensions of the vertebral canal in the neck, are considerably beyond what would be necessary for simply containing the spinal marrow, so that the free lateral movements of the head and atlas can be executed without any risk of pressure on that important part. Hence, spontaneous displacement can occur in these cases to a considerable extent, without impairing the functions of the spinal cord. The case, which I have related above, strikingly illustrates this circumstance; no paralytic symptoms occurred, and the spinal cord, which was still in its situation, when the bone was brought to me, was free from pressure, though the dimensions of the canal had been reduced to about one-half their natural extent. This circumstance

has not been sufficiently attended to by some of those who have described the affection. Observing the diminution of the vertebral canal, they seem to have inferred that the spinal cord must be compressed; and they accordingly mention such pressure among the ordinary effects of the disease*. In most instances, however, as there are no paralytic symptoms, we may safely conclude that there has been no pressure, while in the cases in which sudden dislocation of the dentiform process has

* Boyer expresses his surprise that patients should still live "malgré la gêne prodigieuse qu'un semblable déplacement a du exercer sur la moëlle épinière".—*Trait. des Mal. Chirurg.* Edit. 2. Tom. IV. p. 105.

Rust observes, that the pressure on the spinal cord, which the luxation of the occiput upon the atlas must cause in these cases, is not fatal, because it occurs gradually, p. 82. He mentions that a specimen exists in the anatomical cabinet at the Jardin des Plantes, where the contraction of the vertebral canal is so considerable, that a common quill cannot be introduced into the opening. *Ibid.* As he does not state that he had seen it himself, he probably speaks on the information of others, of the instance quoted at p. 415 from Buffon, in which the transverse measurement of the opening was three lines.

"En même temps que la tête se déplace, la moëlle épinière se trouve comprimée à son entrée dans le canal vertébral, et les mouvemens volontaires de tout le corps, y compris ceux de la respiration, deviennent ordinairement de plus en plus difficiles, ou même impossibles. Cependant, quand la maladie marche avec lenteur, le rétrécissement du canal, et par conséquent la compression de la moëlle vertébrale, peuvent être portés très loin, sans qu'il en résulte beaucoup de difficulté dans les mouvemens." Roche et Sanson, *Nouveaux Elémens de Pathologie Medico-Chirurgicale.* Tom. II. p. 256.

occurred, either from accident or disease, the consequent pressure on the spinal cord has been immediately fatal. Slight compression of the cord or its membranes, or the consequences of such inflammation as may be produced by that pressure or by the disease of the bone, are probably the causes of the paraplegia, or other partial paralysis, which occasionally takes place. In his first observations on the subject, Rust gives a very unfavourable opinion of the probable termination of the complaint. He says that it had been fatal in all the cases under his observation, and he makes it a question whether it admits of cure *. He mentions, however, in his treatise on diseases of the joints, that nature has effected a cure in some instances †, and he cites the case of a patient who was under his care in the Vienna General Hospital with a stiff neck and head, and obliquity of the latter; in other respects he was in good health. He found a motionless state of the three first vertebræ with exuberant bony deposition, and the history of the case left no doubt that it was a spontaneous luxation of the head.

Reil saw three instances, in which the upper vertebræ of the neck swelled suddenly with pain and fever, the head was bent backwards, degluti-

* *Medicinisch-Chirurgische Zeitung*, 1813, V. III. pp. 108, 109.

† *Arthro-Kakologie*, § 118.

tion painful and difficult. In fourteen days the swelling and fever lessened, but the head remained aside and limited in its motions, as he concluded from ankylosis*.

The cases of perfect ankylosis afford the clearest proof that the affection admits of cure, and the considerable number of such instances shews that the prognosis cannot be very unfavourable.

In the museum of St. Bartholomew's Hospital, there is a head dug out of a churchyard, with a dislocated atlas firmly ankylosed to it. The latter bone has been partially displaced towards the left, and at the same time a little twisted, so as to be placed obliquely in its new position. Its right side intercepts about one-third of the vertebral canal. Both occipital condyles are completely ankylosed to the atlas, the union extending also a little towards the front. The bony junction, which seems to be of very old date, is so perfect that the limits of the bones are not visible. The anterior arch of the atlas is situated lower than in the natural state, the space between it and the foramen magnum being at least half an inch.

Sandifort† describes six specimens, belonging

* Fieber-lehre, V. II. p. 356.

† Museum Anatomicum Academiæ Lugduno-Batavæ, Vol. I. p. 143—146, Tab. XIV.

to the anatomical collection at Leyden, of ankylosis between the occiput and atlas. In some the latter bone was partially displaced. The ankylosis embraced one condyle, or both, or even included, in addition, a part either of the anterior or posterior bony arch of the atlas, or one or both transverse processes. In two instances there was a considerable deposition of bony matter about the anchylosed parts.

Wynpersse had seen six specimens of ankylosis between one or both occipital condyles and the occiput ; and he has given an engraving of one *. Boyer met with one at La Charité†. In other cases there is not only dislocation of the occiput upon the atlas, but also of the latter upon the second vertebra, with ankylosis.

Daubenton mentions a remarkable instance, which is probably of this kind, though he does not expressly state that the occiput had been displaced. It is a specimen contained in the cabinet of the Jardin des Plantes. “ The second vertebra of the neck has been displaced, and pushed so far back, that there remains an interval of three lines only between the dentiform process and the posterior arch of the atlas ; the second vertebra is at the same time inclined towards the right. We

* *Dissertatio Medica de Ancylosi*, § 21, Tab. I, Fig. 1.

† *Traité des Mal. Chirurg.* Tom. IV. p. 105.

can easily conceive how such a dislocation might occur ; but, when the vertebral canal has been so considerably contracted, and the spinal marrow consequently so powerfully compressed, it is surprising that the patient should have lived long enough for ankylosis to be effected.”*

Sandifort has described and delineated a specimen in which the atlas and axis have been displaced, the former to the right, the latter to the left, and they have become ankylosed to each other and to the occiput. The change of situation in these bones has lessened the dimensions of the vertebral canal, which measures transversely only half an inch instead of an inch †.

The most extensive occurrence of ankylosis is however seen in another specimen, in which the occiput, all the cervical, and the two upper dorsal vertebræ are firmly consolidated, the head being strongly twisted to the right, so that the left side of the alveolar arch corresponds to the middle of the bodies of the vertebræ. The parts ankylosed in this remarkable specimen are the occipital condyles and front of the foramen magnum to the atlas ; the latter to the dentiform process, that process to the occiput ; the articular processes and the laminæ of all the enumerated vertebræ. The

* Hist. Natur. Générale et Particulière avec la Description du Cabinet du Roi, Tome III. p. 99.

† Museum Anatomicum, p. 148, Tab. XV. Fig. 4.

bodies of the vertebræ were still connected by their fibro-cartilages*. Wynpersse has an instance of luxated atlas and axis, with ankylosis of those bones to each other and to the occiput†.

There is a remarkable case in Reil's Archives, of a patient sixty years old, who had for many years laboured under gout, and whose head was so bent towards the right side, that it was necessary to place a cushion between it and the shoulder. He died of apoplexy, and was examined. The atlas was placed obliquely, and rather thrown forwards, while the axis was pushed so much backwards, that the dentiform process projected into the middle of the foramen magnum, and the vertebral canal was diminished by one half. Although there had been no paralytic affection, it is added that the spinal cord was powerfully compressed‡.

A preparation in the anatomical collection at St. Bartholomew's, exemplifies ankylosis of the occipital condyles and atlas, of the left transverse process of the atlas and occiput, and of the

* Museum Anatomicum, p. 147, Tab. XV. Fig. 1, 2, 3.

Wynpersse shortly mentions an instance in which the occiput and the six upper cervical vertebræ were ankylosed together. *De Ancylosi*, p. 21.

† *De Ancylosi*, p. 19, Tab. I. Fig. 2 and 3.

‡ *Archiv. für die Physiologie*, Vol. IV. p. 407.

articular processes of the second and third vertebræ.

Anchylosis of the dentiform process and atlas is not so frequent an occurrence as that of the occipital condyles. In the collection just mentioned there is a specimen of complete bony anchylosis, between the anterior arch of the atlas and the dentiform process. The occipital condyles are here nearly flattened, so that they probably executed some degree of lateral movement upon the atlas. A curious example of this anchylosis is described and figured by Wynpersse. The dentiform process is united to the atlas, and the articular processes and bodies of the two vertebræ are also anchylosed. A leaden bullet was lodged in the left transverse process of the atlas, which is formed into a rounded bony cyst, firmly containing the foreign body, and constituting an unnatural prominence on the front of the vertebra.

The principles of treatment applicable to these cases, in the commencement and during the progress of disease, are the same which guide our conduct, when disease is seated in other parts of the vertebral column. The immediate vicinity of so many important organs renders it a matter of great consequence to lessen the degree and extent of inflammation, and thus to limit the disorder in its early stage to the parts first attacked. The

local abstraction of blood, with the other parts of the antiphlogistic plan, and perfect rest of the affected bones should be persevered in until all symptoms of active mischief are removed. Counter irritation in its various shapes, and more particularly by the moxa, seton, or issue, will contribute essentially to assist the subsequent restorative process.

Fig. 4.

Fig. 3.

Fig. 2.

Fig. 1.