OBSERVATIONS
ON
VASCULAR APPEARANCES
OF
MUCOUS AND SEROUS MEMBRANES,
AS INDICATIVE OF INFLAMMATION.

By JOHN YELLOLY, M.D., F.R.S.,
PHYSICIAN TO HER ROYAL HIGHNESS THE DUCHESS OF GLOUCESTER, AND
LATE PHYSICIAN TO THE LONDON HOSPITAL, ETC.

READ NOVEMBER 10TH, 1835.

MANY years ago, the Society did me the honour to insert in its Transactions, a paper on the Vascular Appearances of the Human Stomach, which are often mistaken for inflammation of that organ*. In that communication, my attention was principally directed to the various states of the mucous membrane of the stomach; and though I had occasion to advert likewise to the appearances of the villous coat of the intestines, and of their outer, or serous covering, it has long been my intention to offer to the Society, a few additional remarks on the appearances of membranous surfaces generally.

* Medico-Chirurgical Transactions, Vol. IV. p. 371. Published in 1813.

In the former paper, my object was principally to shew :

1st. That appearances of vascular fulness in the villous coat of the stomach, whether florid or dark-coloured, in distinct vessels, or in extravasations of different sizes, are not to be regarded as unequivocal marks of disease ; inasmuch as they occur in every variety of degree and character, under every circumstance of previous indisposition, and in situations where the most healthy aspect of the organ is to be expected.

2d. That from the circumstance of such appearances being in many cases not preceded by symptoms of inflammation, many authors of great eminence were led to the conclusion, that gastritis might exist in the living body, without the indications of either pain or pyrexia ; that it was a very common and frequent complaint, even terminating occasionally in gangrene, though there were no indications during life of its existence : and that the absence of the usual symptoms of inflammation of the stomach, when unequivocal marks of such inflammation were supposed to be found on dissection, could not but be regarded as a very remarkable, and puzzling circumstance.

3d. That the same misconceptions attached to intestinal vascularity ; and that MORGAGNI himself, in order to get quit of the difficulty, was in some cases disposed to attribute the absence of pain, in inflammation of the bowels, to a paralytic affection, which took off the sensibility of the parts ; but that, as this was a circumstance which could but seldom occur, he

came to the general conclusion, that pain and fever were not necessary for the existence of enteritis. HALLER, too, was induced to infer, from so uniformly finding vascularity in his inspections, that inflammation of the bowels was almost constant in every kind of fever, and was frequent in every other complaint.

4th. That the supposition of a necessary connection between vascularity and inflammation, has led to erroneous conclusions in many cases of sudden death, where the effects of putrefaction, and the spontaneous changes which the loss of vitality produces on the human body, were misunderstood, and sometimes confounded with the proper and necessary operation of poisons.

5th. That some diseases of a very obscure nature, as hydrophobia, have been considered as examples of gastritis, and plans of extensive depletion adopted for their treatment, merely from the villous coat of the stomach exhibiting appearances of vascularity, though there was no palpable difference between such vascularity, and that which is ordinarily seen where there had been no affection of the stomach during life.

6th. That such vascularity is entirely venous, though sometimes florid, and sometimes of a dark red; that it depends on a power capable of being exercised on the artery itself at the close of life, which carries on the blood to the veins, after the further supply of fresh blood from the heart is stopped; and that the branched or stellated form of vessels, under which the vascularity usually appears, is capable of

being imitated, whether by injection of the veins with fine injection, or by forcing back with the finger, or the back of a scalpel, the blood from the larger branches of veins into the smaller.

7th. That vascularity soon becomes diffused redness, by transudation through the coats of the containing vessels, just as happens with the bile in the gall-bladder, or in the speedy production of a blue colour, when the mesenteric arteries are injected with a solution of prussiate of potash, and the veins with that of green sulphate of iron.

8th. That digestion of the coats of the stomach after death, is by no means a common occurrence; inasmuch as extravasation does not take place in the great end of the stomach, however thin its coats may be, when fine injection is thrown into it, either by the superior coronary, or the splenic artery, as would be the case, if there were usually an erosion of the ends of vessels produced after death, by the action of the gastric juice, according to the opinion of Mr. HUNTER.

9th. That it is very difficult to ascertain, merely from the aspect of the vessels of a dead part, that they had been affected with inflammation during life, unless for the occurrence of unequivocal results of inflammatory action.

These are the views which I felt myself justified in taking, relative to the subject of vascularity, as referable to the human stomach; and, with certain modifications, they will apply both to *mucous* and *serous* surfaces generally.—Next to the stomach, it is in

the intestinal canal and trachea that vascularity is most frequently found ; but it is in the two former, and more particularly in the stomach, that the greatest disposition to this sort of appearance takes place: for the softness and laxity of texture which distinguish the mucous surface of those parts, and which are admirably fitted for freedom of circulation, render such surfaces peculiarly permeable to vessels through their whole substance, and therefore eminently liable to vascular turgescence after death.

Serous membranes, on the other hand, are smooth, thin, and transparent, and allow vessels to be seen through them, without exhibiting them in their substance, as in mucous surfaces, unless in the inflamed state*. The separation of such membranes from the vascular surface beneath, is easily made, at least in the peritonæum and pleura: and the existence of inflam-

* Professor RUDOLPHI, of Berlin, is of opinion (*Elements of Physiology*, translated from the German by WM. DUNBAR HOW, M.D., Vol. I. p. 92) that serous membranes are to be viewed as merely affording coverings, like the epidermis, to the parts beneath, and an egress to serous fluid, just as the cuticle does to the ordinary exhalation of the skin. It does not appear to me, however, that the evidence which he brings forward in support of his views, is sufficiently decisive of the question, to invalidate the usual opinion entertained upon this subject; namely, that these structures possess minute and colourless vessels, which become apparent only under inflammation. DR. CRAIGIE makes some judicious observations on this subject in his review of KOLK's "*Observationes Anatomici-Pathologici et Practici Argumenti*," in the 45th volume of the *Edinburgh Medical and Surgical Journal*, p. 511.

mation in them is readily detected, by the effusion of coagulable lymph which takes place in or upon them, very soon after its occurrence.

On the surface of the brain and cerebellum vascular appearances often occur, which, like those already mentioned, are principally the result of venous accumulation after death.

The *spinal marrow* is not the subject of convenient observation, and is therefore seldom inspected, except with some precise and determinate view. Its usual appearances are therefore but little known; but I have made them the objects of considerable attention, and have much reason to suppose, that a state of vascularity which is in no way morbid, has been sometimes regarded as imparting a certain character of spinal inflammation to some diseases of an obscure nature, as tetanus, which did not actually belong to them.

In my former paper I thought it expedient, in order to obtain a knowledge of those appearances of healthy stomachs which were the product merely of a particular mode of death, to give an account of the examinations of five malefactors who forfeited their lives to the justice of their country; and a plate which accompanies the communication, affords, in addition to some other sketches connected with the subject, a correct delineation of such appearances in one of them. I have, since that time, examined the bodies of several other persons who were executed, with a view to observe the vessels generally, and in particular the state of those of the spinal column.

The drawing which accompanies this paper, is a good example of the appearances in the spinal column of one of these cases, and may serve as an accompaniment to that above alluded to, which it very much resembles.

From this it may be seen, that a florid vascularity, communicating a scarlet hue to the whole column, is nothing more than venous turgescence, accompanied, as in the case of the stomach depicted in my former paper, with slight extravasation.

To those who have attended much to pathological inquiries, it must be obvious, that nothing is more important for the successful prosecution of morbid anatomy, than the habit of accurately connecting the phenomena which present themselves after death, with the symptoms which occur during life; and of carefully distinguishing between the varieties of form and appearance which exist in healthy parts, and such as are the result of derangement or disease.

Very careful attention and discrimination are invariably required on this subject; and even with all the aids which the most accurate knowledge, and the most extended experience can communicate, it is not always that the best qualified pathologist feels himself justified in speaking with decision, as to the nature of certain appearances which come before him.

This is particularly the case with *inflammation*, a subject of the highest importance in pathology; for while on the one hand, it either constitutes or gives rise, in some one or other of its various seats and forms, to many of the most important diseases, it is

likewise on the other; one of the principal agents by which injuries are repaired, and maladies prevented; thus affording an interesting example, which the well directed studies of modern times have so successfully elucidated, of the provident mode in which nature employs her resources, and aims at accomplishing as many ends as possible by the same means. The subject of inflammation is very extensively interwoven with medical reasoning and practice. Where its existence is well marked, we have an important guide in our curative indications; and even its fainter appearances may exercise some portion of directing influence on the treatment of disease*.

The phenomena of redness, swelling, heat, and pain, sufficiently indicate the existence of inflammation in external parts; but in affections of internal organs, we are often led to our conclusions by imper-

* In the fifth volume of the *Mémoires de l'Académie Royale de Médecine* lately published, are some observations on the cure of wounds *without inflammation*, by DR. MACARTNEY, the distinguished professor of Anatomy and Surgery in the University of Dublin, which, if fully established, would affect much the prevailing ideas on the subject. In this communication we are informed, that DR. MACARTNEY has for more than thirty years been of opinion, that inflammation retards the healing of wounds when moderate, and entirely prevents it when very considerable; that the proper means of treatment are, to soothe a wounded part as much as possible, by repose; by the absence of every kind of irritation, and by the application of water or vapour of regulated temperature; and that, when by these means inflammation is prevented, a cure takes place without that deposition

fect analogies, and by symptoms and appearances more or less doubtful. Sensations themselves are, to a certain degree, deceptive or equivocal. They may be connected with very different states of body, and may be very dependent on the temperament of the patient by whom they are described. In internal diseases, we are by this means often left to form our conclusions, as to the existence of certain states of disease connected with the extinction of life, by the appearances offered on examination after death.

Where inflammation has existed *externally*, death makes a very important alteration of appearance. The redness goes off, except where effusion has taken place, and the remains of it are seen; while the tumour sinks, except in as far as any deposition in the cellular membrane has taken place, and may render it permanent. These particular changes are necessarily owing to the altered state of the blood-

of coagulable lymph, which is necessary to ordinary healing by the first intention. The communication is of great practical importance, but if DR. MACARTNEY'S opinions are correctly represented, (of which, from some observations made by him in the Medical Gazette, there is some reason to doubt,) I should feel much difficulty in adopting the conclusion, that irritation, under the circumstances of injury mentioned by him, could ever be so far prevented or repressed by medical treatment, as to preclude altogether the effusion of coagulable lymph from divided vessels. It is easy to conceive, that a very small portion of that deposition, may be a sufficient bond of union under the favourable position contemplated, but not that it should be entirely wanting; which would appear to me to involve the anomaly of a change of law, in circumstances very nearly approximating to each other.

vessels after death, when both the arteries and veins have parted with their contents, and left the parts to which they belonged nearly exsanguous.

But in the *internal* organs, and more particularly the abdominal viscera, various modifications take place, from the peculiar situation which these parts occupy in the animal body. Being nearly connected with a double series of venous structure, that of the liver, and that of the heart, in which a considerable portion of the blood is concentrated after death, a retarding or obstructing cause operates on the veins which belong to them, and which, added to the softness of texture to which I have already alluded, very generally produces more or less fulness of the vessels, and thus an appearance of more or less vascularity, both on the external and internal surfaces of the chylopöietic viscera.

It may be readily seen, therefore, that vascular fulness in the dead body is exceedingly delusive, and cannot usually be regarded as any indication of what was the state of vessels during life, where both arteries and veins bear their part in the circulation, and where there are none of the causes of accumulation in the lungs or heart, or in the vena cava or vena portæ, which take place at the close of life. Under this view of the subject it is likewise apparent, that though a florid state of vessels after death has often been regarded as a proof of arterial congestion, yet it merely arises from the arterial character of the blood remaining in the veins, for some time after its transmission from their continuous arterial capillaries.

Since the time when my former paper was published, many years have elapsed of a more active and general cultivation of every branch of medical science, than any similar period ever before witnessed. The consequence has been favourable; and as far as morbid anatomy is concerned, a more correct knowledge has been thus obtained, of the nature and indications of disease; and many misconceptions on the subject have been cleared up, or removed. Few of the pathologists of this country are now, therefore, disposed to consider mere turgescence of vessels as indicating inflammation; and they consequently look to some of the results of inflammatory action, for proofs of its existence in internal parts.

The enterprise and energy of continental, and particularly of French physicians and surgeons, during the same period, have not been less marked and creditable; but it is only of late years that their attention has been applied very strongly to the importance of a correct discrimination relative to the vascularity of membranes, to which I directed the attention of the professional public, through the medium of our Transactions. There are still, however, as it appears to me, opinions entertained relative to this subject, by gentlemen of much character and influence in the profession, which do not altogether accord with the present state of pathological knowledge; and on these it is proper that I should offer some observations.

The frequent appearances of vascularity which exhibit themselves in the examination of bodies, have had a powerful influence in forming the pathological opi-

nions of two distinguished physicians of the present day, TOMMASINI and BROUSSAIS. Both these gentlemen feel themselves warranted in concluding, that general fever is to be attributed to local inflammation alone; that it is never to be viewed as an idiopathic disease; and that it consists of an assemblage of symptoms, depending on the operation of such local affection on the constitution. The vascularity frequently observed in the lining membrane of the stomach and intestinal canal, BROUSSAIS regards as gastro-enteritic inflammation; and as this is generally seen on examining the bodies of persons affected with fever, he has been induced to conclude, that fever depends, in all cases, on inflammation of the gastro-enteritic mucous membrane.

TOMMASINI, on the other hand, though he considers fever, equally with BROUSSAIS, as produced by inflammation, regards its seat as in the brain. But he carries his opinions as to the extensive bearings of inflammation, in the production of disease, so far as to believe, that much the greater number of maladies affecting the human body arise from *inflammation*, either acute or chronic, apparent, deep seated, or concealed, and of various extent and relation; and that this is particularly the case with many obstinate convulsive affections; with hysteria, tic douloureux, sciatica, and the different forms of neuralgia, in many of which stimulant and sedative remedies have been given in vain*.

* TOMMASINI's Précis de la nouvelle Doctrine médicale Italienne, &c., traduit de l'Italien, par P. L. VANDER LINDEN, p. 69. Paris, 1822.

In considering the application of these doctrines in pathology, whether in their more limited or extended view, it has been a natural object of inquiry with the best observers, whether the appearances of inflammation which were represented as thus exhibiting themselves, were original, necessary, and constituent parts of the disease, or were only accessory and occasional. There is much reason for believing that the latter is the fact ; but though there is no doubt, that both in fevers and many other complaints, unquestionable marks of inflammation are occasionally found in various organs of the body, and particularly in the intestinal canal and encephalon, yet it is an important subject of consideration, in connection with the present paper, whether the indications of inflammation mentioned by those gentlemen and their supporters, as so often occurring in morbid dissections, are always to be viewed as unequivocal, or as being frequently nothing more than turgescence or congestion of vessels.

TOMMASINI informs us, that in the examination of one hundred cases of fever, the brain exhibits undoubted marks of inflammation in at least ninety of them* ; but he does not mention the appearances on which his inferences as to the existence of inflammation are founded. If those appearances were the

* "... si l'on consulte les résultats des dissections pathologiques et si l'on apprécie bien les faits que chacun de vous aura sans doute pu observer ; l'on voit que, sur cent morts de Typhus, ou de Fièvre nerveuse, quatre vingt-dix au moins nous présentent dans le cerveau les traces manifestes et les résultats d'une véritable inflammation."—TOMMASINI, p. 72.

unquestionable and necessary results of inflammation, the correct pathologist could have no difficulty in admitting his conclusions ; and it would indeed be a matter of great surprise, that there should be any hesitation in acknowledging them, with such weight of evidence in their support. But my own knowledge of the appearances of the brain in fever patients, and the descriptions given by those whose opportunities of observation on the subject have been very extensive, lead me to doubt, whether the appearances in question consist in more than turgescence of vessels, between which, and inflammation, it is of such high importance to discriminate.

In reference to this point, M. ANDRAL states his belief, that there are no characters by which it can with certainty be ascertained, whether the sanguineous injection often observed in the brain, is the result of inflammation, or cerebral congestion ; and he adds, “ that simple congestion often passes insensibly into inflammation, a circumstance which is not peculiar to the brain, for it is not always possible to distinguish positively between mere engorgement of the lungs, and recent pneumonia. In typhus fever of every form,” he continues, “ the nervous system acts a very important part ; there is disturbance in the brain, as well as in the alimentary canal ; but inflammation is no more necessary in the one, than the other.” *

* *La Lancette Française*, *Gazette des Hôpitaux civils et militaires* pour Dec. 8, 1835. *Med. Gazette* for Jan. 9, 1836.

The zeal and ability with which M. BROUSSAIS has propagated his opinions, and the seductive influence of doctrines, which promise a ready and unhesitating removal of previous difficulties, seem to have led to an extensive reception of the deductions of that eminent physician, in respect to the dependence of fever on gastro-enteritic inflammation.

In the early period of his professional labours, M. ANDRAL, to whom the world is greatly indebted for the industry and talent of his pathological researches, appears to have been under considerable influence from the opinions of M. BROUSSAIS, which greater experience, and larger opportunities of observation, have enabled him, in some degree, to modify. His favourable impressions were, indeed, from the first, blended with considerable distrust of the extent to which BROUSSAIS' opinions could fairly be admitted; but in examining, with care, the dissections of fever cases contained in the 1st volume of his *Clinique Médicale*, (which are thirty-eight in number, the whole number of cases described being 117,) it will, I think, be seen, that notwithstanding the general precision and correctness which they exhibit, some of them are described as examples of gastro-enteritic inflammation, which can scarcely be regarded but as instances of congestion*.

The brown spots too, which are mentioned as indi-

* In Case 82, Tom. I. p. 186, it is said, that the stomach itself presented a *moderate degree of inflammation*, “ *mediocre degré de phlegmasie* ; ” but the appearances, as given at p. 185,

cative of inflammation * ; and the greyish brown tint ("teinte de gris brunâtre") of gangrene, will be readily recognized as resembling much that transudation through the coats of veins, which a short time generally suffices to produce after death † ; while the follicles which not unfrequently present themselves in va-

and at p. 358, are merely those of injection of the pyloric portion ; "la portion pylorique était fortement injectée."

Case 95, p. 268, is mentioned as an example of a "Gastrite intense" attributed to the use of cinchona, and evinced by the occurrence of dryness of tongue about four days before death. But the appearances are those of injection only : "La membrane muqueuse dans la presque totalité de son étendue était rouge et molle."—p. 266.

Case 107, p. 316, is spoken of as an instance of gastro-enteritic inflammation occurring at the latter period of life ; while the appearances are only those of injection of the mucous coat, "La muqueuse elle-même était injectée mais non uniformément rouge," with the large veins below filled with black blood, "remplies de sang noir."

In Case 113, p. 338, "une longue ribbande rougeâtre formée par des vaisseaux injectés, autour des quels existait de petits points rouges," is mentioned at 339 as "une gastrite partielle peu intense."

In Case 114, p. 339, a "petite plaque rouge dans l'estomac," is viewed as having a contingent connection with gastro-enteritis, p. 344.

In Case 115, p. 344, we learn that the pulse only became frequent during the two days previous to death, notwithstanding the inflammation of the digestive canal ; but the appearances are merely those of the stomach "généralement injecté, à sa surface interne ;" and at the posterior part of the great curvature, having "plusieurs plaques d'un rouge brunâtre," &c.

* ANDRAL'S Clinique Médicale, Tom. I. Case 89, p. 223.

† Ib. Case 97, p. 277.

rious parts of the stomach, may easily be mistaken for the minute ulcerations of the size of an ordinary pin's head, which are stated by M. ANDRAL to be occasionally seen in the villous coat of that organ *. It might readily, however, be ascertained, whether, at the place where ulceration was supposed to exist, the vessels were actually divided, so as to admit the passage of injection through their open extremities, in the manner which I have mentioned, as affording a ready means of determining the existence of digestion of the stomach after death.

In order to exemplify the occurrence of such follicles, I have the honour to transmit, for the inspection of the Society, a very interesting drawing of a stomach, which SIR ASTLEY COOPER was so obliging as to get made for me many years since, and to which I alluded in my former paper; in which there were found a great number of depressions of various sizes, which a superficial examination might refer to ulceration, but which we found to be natural depressions followed by the mucous covering. The patient had no symptom of stomach-disease.—Notwithstanding, however, the occasional slight deviations from precision, which I have had occasion to notice in the *Clinique Médicale*, M. ANDRAL inculcates, in various parts of that valuable work, the importance of not mistaking congestion of vessels for the result of inflammatory action, as appears to have been so

* ANDRAL's *Clinique Médicale*, Tom. I., Case 97, p. 277.

uniformly done in BROUSSAIS' system, as well as in that of TOMMASINI.

In M. ANDRAL's later and more peculiar work*, the Précis d'Anatomie Pathologique, the distinctions between congestion and inflammation are more pointedly laid down and systematized, and are made to occupy a more prominent position than they possessed in the occasional remarks of the Clinique. In this work, he rejects the term *inflammation* as old-fashioned, vague, and confused; though I would submit that there is nothing more inconvenient than to part with an appellative which has been long accredited and understood. That its etymology may not be in exact accordance with its present signification, is necessarily the case with much of the language even of science. But we should gain little in precision by the employment of the designation which M. ANDRAL would substitute, that of *Hyperémie active ou sthenique*; nor does he appear to be at all able practically to follow up, either in his Précis d'Anatomie Pathologique, or in the Clinical Lectures which lately appeared in the Gazette des Hôpitaux, the abandonment of the term which he has deemed it proper to condemn.—Congestions, under whatever circumstances they may arise, he regards as forming other descriptions of hyperemiæ, as the *Hyperémie passive ou Asthenique*, when there is a diminished

* M. ANDRAL's Clinique Médicale, was based on a report of cases visited by M. LERMINIER, at the Clinique of La Charité.

tone in the capillary vessels; *Hyperémie mécanique*, when there is an obstacle to venous circulation; and *Hyperémie cadaverique*, which occurs after death.

In considering the last division of the hyperemiæ, the *Hyperémie cadaverique*, his deductions are very analogous to those arrived at in my former paper, both as to the cause of such congestions, (which he imitates as I have done, by pressing back the blood contained in the veins into the capillaries, by the back of a scalpel;) the circumstances under which they occur, and their liability to be mistaken for morbid appearances. He considers it as established by his observations, that the hyperemiæ which have come on during life, or after death, are not always to be distinguished from each other by their anatomical characters; and that simple inspection does not, in a great number of cases, suffice to determine the cause of sanguineous congestions found in pathological examinations *. The gastro-intestinal membranes, he continues, may be indifferently white or red, according to the circumstances under which death has taken place; but as the circumstances which produce a red colour are most frequent at the close of life, it will

* “ Telles sont les différentes classes d’Hyperémies survenues pendant la vie ou après la mort, dont une rigoureuse observation me semble démontrer l’existence; et s’il est vrai, comme je l’ai établi qu’elles ne se distinguent pas toujours les unes des autres par les caractères anatomiques bien tranchés, il devient manifeste que dans un grand nombre de cas, il ne suffira pas de la simple inspection pour décider qu’elle cause a produit les congestions sanguines trouvées sur un cadavre.” ANDRAL’s Précis, Tom. I. p. 68.

oftener happen, that congestion of vessels in the alimentary canal will be found in the dead body, than a want of colour. It does not follow, however, he adds, that a red colour is any proof that disease existed in the alimentary canal during life; inasmuch as an intestine found to be red in the dead body, was not necessarily so during life*.

M. ANDRAL attributes the position of intestinal congestion, with Messrs. TROUSSEAU and RIGOT, to the mechanical descent of the fluid blood, according to the posture which the body is made to assume after death; and in one case, where the vascularity was in the anterior part of the stomach, owing to the body lying on its back, he was disposed to refer the appearance to *gastritis*, which, with much candour, he states he should not do now†. That this cause operates, to a certain extent, in producing the phenomenon in question, particularly when the larger

* —“ La membrane muqueuse gastro-intestinale peut-être indifféremment blanche ou rouge, sans que l'une ou l'autre de ces couleurs indique nécessairement que la membrane soit dans un état morbide; elle est ou blanche ou rouge, à divers degrés, selon qu'avant la mort a existé l'une ou l'autre des conditions soit mécaniques, soit organiques, soit vitales, que nous avons cherché à faire connaître. Or, comme, parmi ces conditions celles qui en déterminent la colorisation rouge existent le plus fréquemment, il s'ensuit que sur la cadavre on doit plus souvent rencontrer l'injection du tube digestif, que son état de décoloration.” ANDRAL's Précis, Tom. II. p. 10.

† “ Je crus alors qu'il y avait *gastrite*; je ne l'affirmerais pas aujourd'hui.”

vessels are concerned, is quite certain; but, though I have remarked, that vascularity is found “ principally in the posterior part of the great end of the stomach, and in the lesser curvature,”* yet nothing is more common than to see great turgescence of vessels on the upper surface (in relation to the position of the body) both of the intestines and brain; and it occurs everywhere in the minute vessels of the inner membrane of the stomach and bowels, where an attraction is exercised between the coats and their contents, which prevents any material change of position in the latter.

M. ANDRAL is disposed to refer the transudation which takes place through the sides of vessels some time after death, to *putrefaction*†. But I see no reason to alter the opinion which I have expressed on this subject, namely that “ putrefaction will doubtless increase, but it does not seem at all *necessary* to transudation.”‡ The animal fibre has a power to resist the passage of a fluid during life; but very soon after death, and long before putrefaction can be considered as having at all taken place, the power is lost, and various well known phenomena speedily occur in every part of the body, which are dependent on that unknown and inscrutable change, which the loss of vitality has effected in the animal machine. Transudation of bile from the gall-bladder, and of

* Medico-Chirurgical Transactions, Vol. IV. p. 388.

† ANDRAL's Précis, Tom. II. p. 15.

‡ Medico-Chirurgical Transactions, Vol. IV. p. 398.

blood into the thoracic or abdominal cavities, are among the commonest examples; and the veins of the stomach and bowels will exhibit the same appearance in a day, or day and a half from death; but in a still shorter time if an inverted vascular stomach is extended upon a flat surface, and kept moist and undisturbed. I may mention too, as an additional proof of the same circumstance, the point mentioned at page 4 of this communication; and may add, that the same will happen in a tied carotid artery or intestine, containing either of the fluids employed, and made to dip into a vessel containing a portion of the other*.

It would lead me much beyond the limits which I have prescribed for my present communication, to enter into a particular examination of the opinions which M. ANDRAL entertains, as to the existence of organic affections of the stomach generally, and their connexion with complaints of the chest and encephalon. It appears to me, however, that his deductions as to the extent to which inflammation of that organ prevails, whether acute or chronic, original or accessory, are not warranted, either by the symptoms which he narrates, or the phenomena described by him as appearing after death. That the stomach should exhibit functional disease more frequently than any other organ of the body, is not at all surprising, when we consider, not only its remarkable sympathy with every part of the animal œconomy, and its liabi-

* Medico-Chirurgical Transactions, Vol. IV. p. 398.

lity, therefore, to be more or less affected in all kinds of complaints, but its unceasing employment, and its perpetual exposure to the somewhat unreasonable duties, which the habits of society are so frequently apt to impose. That it should, however, be subject to the large amount of organic affection mentioned by M. ANDRAL, is not only, as it appears to me, an error in pathology, but would seem to indicate that this viscus, so essential to the support of life, and so remarkable for its power of accommodation to every variety of food, is devoid of much of the conservative faculty with which the animal body is so eminently endowed, and is continually liable, to a much greater extent than most other organs, to have its peculiar and important functions injuriously and permanently disturbed.

M. OTTO appears to have attended much to the appearances of vascularity after death, and is of opinion that the "reddening of particular viscera, in persons who have died of adynamic fevers, which BROUSSAIS and TOMMASINI have considered as a sign of inflammation," as "not in the least so, but the consequences of mere irritation, congestion, obstruction, &c.)*" At the same time he mentions, as of great importance, "*occult* or *hidden* inflammations, so called, which are seated especially in the viscera of the three great cavities of the body, and during life often exhibit no distinct or determinate symptoms."

* OTTO's compendium of human and comparative Pathological Anatomy, translated by SOUTH, p. 47.

Whether such cases were connected with ulceration, or the deposit of some unequivocal product of inflammation, we are not informed ; but if such were not the case, it would be difficult to regard them in any other light, than as mere examples of vascular turgescence.

MM. TROUSSEAU and RIGOT, and M. BILLARD have turned their particular attention to the nature of the vascularity which is found after death. The former, in a memoir published in the *Archives générales de Médecine* *, give many interesting details on the subject ; and shew, by careful investigation, not only that the redness so often observed in the mucous membrane of the lungs and alimentary canal, comes on subsequently to death, and is independent of disease ; but that this is likewise the case with the redness seen in the inner membrane of the blood vessels and heart.

M. BILLARD has, with much zeal and industry, prosecuted a long and laborious inquiry into the gastro-intestinal mucous membrane in its healthy and diseased states, and during every period of life †.

He admits the difficulty of discriminating accurately between inflammation and passive congestion of the alimentary tube ; and describes, very correctly, and at great length, the various appearances which

* For October and November, 1826.

† De la Membrane Muqueuse gastro-intestinale dans l'état sain et dans l'état inflammatoire ; ou Recherches d'Anatomie Pathologique sur les divers aspects sain et morbides qui peuvent presenter l'estomac et les intestins. Par C. BILLARD.

this passive state of congestion assumes ; but at the same time he considers the redness which many anatomical writers describe as being ordinarily found in the villous coat of the stomach and bowels, as only existing when death may have occurred during digestion*. It may be remarked, however, that though the stomach, as a living secreting organ, becomes redder during digestion, by having an augmented distribution of blood through both the arteries and veins of its secreting surface, yet this is no criterion of the state of things after death, when its arteries become entirely empty, and its veins often nearly so.

In considering the circumstances occasioning what he terms passive congestion, M. BILLARD does not seem to recollect that such circumstances occur under almost every kind of death ; and hence he reprehends, and, as it appears to me, somewhat unduly, the opinions of many of the best anatomists, and most correct pathologists of his country, such as HABICOT, SABATIER, PORTAL, and BOYER, who have all of them offered a very judicious and appropriate caution, against mistaking a congested state of vessels for an inflamed one †.

* “ Je n’ai trouvé cette coloration rouge, ou plutôt rose, de l’estomac, ou du duodénum, que sur des cadavres d’individus *morts pendant la digestion*.”—BILLARD, p. 95.

† Of SABATIER, M. BILLARD says, “ Sabatier indique a tort la couleur rouge pourpre, comme étant naturelle à la membrane muqueuse de l’estomac. Il partage, en cela, l’erreur du célèbre chirurgien,” (Habicot,) “ dont il emprunte l’opinion.” But Sabatier only says, “ Le grand nombre des vaisseaux qui s’y distribuent lui *donnent souvent* une couleur pourpre obscur,”

In adverting to the present state of pathological opinions relating to vascularity, it is particularly satisfactory to me to find, that there is so much accordance between my own conclusions, and those which have been arrived at by gentlemen, who have availed themselves, so sedulously, of the ample field of pathological investigation which they possess in a neighbouring kingdom. It would have saved them, however, much labour, and perhaps, to a certain degree have modified some of their deductions, had they been so far acquainted with our Transactions, as to have known what had been done in this country on the same subject. Life is, indeed, too short, and the difficulties which attend the prosecution of philosophical inquiries too considerable, to afford that there should be no division of labour in science; and that men of talents and energy should occupy themselves in investigations which have already been prosecuted in other quarters, without being aware of what has already been done on the same subject. Science

which is the fact. I have quoted in my former paper, the original and discriminating remark of HABICOT on this subject, which M. BILLARD seems to have entirely misconceived. The passage is as follows: “ La partie interieure d’iceluy vëtricule estât nettoyée demōstrerez, tant la tunique interne qui est velontée, et comme l’abondissemēt qui se faict des vaisseaux provenās de la vene gastre epiploïque en la partie gibbe dudit vëtricule, le rend de couleur noirastre, qui a faict croire à beaucoup d’inexpert, en la medicine et chirurgie, aux maladies violentes y avoir en poison ou venosité en leur mort.” HABICOT NICOLAS, *Semaine ou Prâctique Anatomique*, p. 48, leçon 3, Paris 1610.

is common property, of which every one may avail himself at his pleasure ; but the talents of able men belong also in a certain degree to the public ; and it is a species of duty for such as are qualified, by their abilities and industry, to enlarge the boundaries of science, to employ themselves in the most provident manner possible for the common benefit*.

The study of the phenomena, and various results of inflammation, has been very successfully prosecuted by modern pathologists ; but still it has not been sufficiently ascertained, on what circumstances the peculiar affection depends, which constitutes unequivocal inflammatory action. The general suffusion of a blush, the increased colour produced by exercise, or by rage, and the flush of hectic fever, all of them resemble much the first appearance of erysipelas ; but

* Considering the almost national character of the Medico-Chirurgical Transactions, it is very remarkable how little they seem to be known among the best French medical writers, even in reference to some of our most distinguished names. A curious exemplification of this occurs in CRUVEILHIER, who informs us, in the 3d livraison of his *Anatomie Pathologique*, (pl. 6, p. 10) that he lately made a very convincing experiment (“une expérience bien convaincante”), to prove, that disease on one side of the spinal marrow, produces its effect on the same, and not the opposite side of the body. But the point had been examined many years previously by SIR ASTLEY COOPER, at my request, in consequence of the contradictory evidence on the subject ; and a similar conclusion, confirming a valuable experiment of GALEN, arrived at by direct experiment on a dog, of which an account was published by me in the first volume of the *Medico-Chirurgical Transactions*, p. 197.

no length of continuance, nor frequency of recurrence of these phenomena, will give them at all the character of inflammation ; while the slightest efflorescence may, in erysipelas, go on progressively, to the exhibition of all the local, as well as the general characteristics of that disease.

The local congestions which are apt to occur in various parts of the body, from slight causes in certain habits, neither constitute inflammation, nor give origin to it. They may produce effusion in the head, and in this way occasion the most serious apoplectic, or paralytic affections ; or they may occasion discharge of blood from the lungs, or Schneiderian membrane ; but they do not give rise either to phrenitis, or pneumonia ; nor is that state of vessels which in the stomach occasions hematemesis, and in the large intestines hæmorrhoidal discharge, productive of gastritis, or of inflammation of the bowels.

Mr. HUNTER saw and entered into these difficulties ; for in speaking of the vascularity of membranes, he remarks, that there is no difference between the blush which belongs to “briskness in the circulation in the part at the time,” and “the very first step in inflammation” ; that both are connected with a certain “action of the vessels” ; and that when this blush is “an effect of an inflammatory cause, *it is then only that the inflammation has not yet produced any change in the natural structure of the parts, but which it will soon do.* What the action is,” he continues, “or in what it differs from

the common action of the vessels, is not easily ascertained, since we are more able to judge of the effects than the immediate cause.”*

Not much has yet been done to remove the obscurity of the subject ; but the important influence which modern physiology has referred to the nervous system, in secretion, has been considered, and with much appearance of truth, as applicable to the production of that peculiar change in the action of vessels, on which depend the more characteristic phenomena of inflammation; and, in particular, the property acquired by inflamed vessels of throwing out coagulable lymph.

The existence of pain in an inflamed part, has, from the earliest periods, been regarded as one of the characteristics of inflammation ; and this symptom not unnaturally, therefore, refers itself, and the phenomena connected with it, to some affection of the sensorium. But degrees of pain are very various, according to the character of the part affected ; and even the total absence of sensibility is, in some cases of anæsthesia, found to be not incompatible with the existence of the other phenomena of inflammation. The circumstances which produce pain are likewise very obscure. Distension will not alone give rise to it ; for in all the states of congestion to which I have alluded, pain is neither a frequent, nor a necessary attendant. Under a great variety of circumstances, likewise, pain can be present without inflammation,

* HUNTER'S Treatise on the Blood, Inflammation, and Gun-shot Wounds, p. 281.

of which headache, gastrodynia, and the peculiar state of bowel which exists in colic are well marked examples, to say nothing of those diseases which are denominated more particularly neuralgic. For distension, therefore, to produce pain, it must exist in a part otherwise disposed to it; and of the peculiar nature of this disposition we are at present entirely ignorant.

In mentioning the existence of doubts as to the amount of our acquaintance with particular subjects, to which my inquiries into the vascular appearances of membranes have so much led me, the task, I am aware, is somewhat ungracious; for it is much more agreeable to refer to the possession of knowledge, than to hesitate in admitting its extent or application. But though there may be many, as DECAN-DOLLE well observes, "*qui aiment à embrasser des hypothèses hardies, au risque même des erreurs,*" rather than submit to the feelings of suspense and indecision which arise from the difficulties which there often are in forming correct conclusions, yet in no branch of science is it more important, and perhaps less easy, than in Pathology, to restrain or to temper the love of speculation, and to guard against its powerful influence.