OBSERVATIONS
ON
FIBRO-CALCAREOUS TUMOURS
AND
POLYPI OF THE UTERUS.

BY ROBERT LEE, M.D., F.R.S.

PHYSICIAN TO THE BRITISH LYING-IN HOSPITAL, AND SAINT
MARYLEBONE INFIRMARY; LECTURER ON MIDWIFERY AT
ST. GEORGE'S HOSPITAL.

READ MAY 27TH, 1834.

THE most important diseases of the human uterus accompanied with sensible alteration of structure, may be divided into three classes.

I. Those which are produced by inflammation of one or more of the textures which enter into the composition of the uterus.

II. Those which arise from the formation of tumours in the parietes of the organ, or from enlargement of the glands situated in its orifice, which have no tendency to degenerate into a malignant form, and do not contaminate the surrounding structures.

III. Those which result from a specific and malignant action of the uterus, by which its dif-

ferent textures and the adjacent viscera become disorganized.

The phenomena and the treatment of the diseases comprehended in the first of these classes, I have endeavoured to describe in the fifteenth and sixteenth volumes of the Medico-Chirurgical Transactions, and I now propose to make some observations on the pathology of fibro-calcareous tumours and polypi, the most important organic affections included in the second class of diseases of the uterus.

The fibrous tumour, or fleshy tubercle of the uterus, as it was termed by Dr. William Hunter, is sometimes met with not larger than a pea, in other cases it grows as large as a walnut, and occasionally is equal in size to a cricket-ball, or even the gravid uterus in the ninth month. It is generally of a globular form, or kidney-shaped, and when cut into, presents a laminated or radiated semi-cartilaginous structure, the fibres being often disposed in a concentric manner. At other times, this tumour has a granular appearance, or seems to consist of a congeries of smaller tumours, each having a thin capsule of cellular membrane. Most frequently, it has a yellowish white colour, but several specimens of the disease have been of an ash-grey colour, or approaching to a dark slate. When large, the tumour is often unequal on its surface, being lobulated or divided by deep fissures, and arteries and veins of

considerable magnitude can be traced into its substance. Cavities containing a bloody or dark-coloured gelatinous fluid are sometimes formed in the central part of the tumour, probably by a process of softening which its substance undergoes. In a specimen of large fibrous tumour imbedded in the walls of the uterus, which was removed from the body of a woman who died in the Saint Marylebone Infirmary, there is a considerable cavity which contains a coagulum of blood.

In other cases, the tumour does not manifest a disposition to become softer as it enlarges, but its density gradually increases, until the whole or the greater part of the mass has become cartilaginous, or like intervertebral substance, without vessels containing red blood; or calcareous depositions are gradually formed in the substance of the tumour, until it is either partially or completely converted into a concretion composed of carbonate and phosphate of lime. Most frequently the calcareous depositions are first formed in the central and most dense parts of the tumours, but this is not invariably the case, and in a few rare instances the deposit has taken place around the circumference of the tumour, and has enclosed it, as the shell encloses the kernel of a nut. If injection be thrown into the vessels of the uterus, it does not penetrate the substance of the tumour when in this dense state. Fibro-calcareous tumours of the uterus are generally soft and porous, like pumice-stone; but instances have occurred in which they were so hard,

that they admitted of being polished, like ivory or marble. Two specimens of this description are in the museum of St. Thomas's Hospital. Portions of these tumours were analyzed by Dr. Bostock, ten years ago, and were found to consist chiefly of phosphate of lime.

Andral, on the authority of Brugnatelli, states that carbonate and phosphate of lime, with animal matter, enter into the composition of these bodies. Breschet states, that one of the uterine calculi examined by Brugnatelli was a shapeless mass, with a white unequal surface; it emitted a peculiar odour; and was insipid, and insoluble in water. Being broken with a hammer, the surprise was extreme when a portion of the tibia of a chicken was discovered in the centre. The whole white mass forming the calculus was phosphate of lime. The second calculus, when divided into two equal parts, presented on the surface a great number of crystals of ammoniaco-magnesian phosphate; the centre was composed of phosphate of lime. There are several circumstances mentioned in the account of these concretions, which might lead us to suspect that they were urinary and not uterine calculi which Brugnatelli analyzed*.

The body of a woman advanced in years was examined by my colleague Dr. Hope, at the St. Mary-

* Dictionnaire de Medicine, Tom. IV. Art. Calcul.

le-bone Infirmary, on the 17th of August, 1832. The uterus was larger and much heavier than natural, and under the peritoneal coat of its fundus were several small fibrous tumours, with calcareous deposits. On dividing the parietes, a soft yellowish-coloured calcareous tumour, the size of a goose's egg, was found situated under the lining membrane, at the posterior part of the uterus, and distending it like an ovum at the end of the second month of gestation. There are portions of this tumour in which the fibrous structure still remains distinct. The parietes of the fundus and body are thinner than natural; the os and cervix uteri are healthy, but the latter is greatly elongated. The existence of an organic disease of the uterus was not ascertained before death*.

Mr. Henry Johnson shewed me a large fibro-calcareous tumour imbedded in the walls of the fundus uteri, which he had removed from the body of an aged woman who died in St. George's Hospital, in whom, during life, the existence of an organic affection of the uterus had not been suspected. Towards the circumference of the tumour, the fibrous structure was distinct, but the central part consisted of a hard, yellow-coloured concretion of carbonate and phosphate of lime. The parietes of the uterus surrounding the tumour were hypertrophied.

* Cases of elongated cervix uteri have been recorded by Morgagni, Levret, Lallemand, Desormeaux, Cruveilhier, Boivin, and others.

In Mr. Howship's collection, there is a uterus which weighs several pounds, from the presence of a number of dense fibro-calcareous tumours in its parietes. One of these, which hangs by a slender peduncle from the fundus uteri, is only partially converted into calcareous matter, the remaining portion exhibits the usual compact structure of the fibrous tumour of the uterus. Mr. Howship presented a portion of one of these concretions to Dr. Bostock for analysis. When sawn across, the tumour appeared harder than bone, and might have been polished like ivory.

On the 21st of January, 1834, a woman, æt. 64, died of apoplexy in the St. Mary-le-bone Infirmary. Mr. Blenkins examined the body. The fundus, body, and cervix uteri were all reduced to a very small size. Adhering to the fundus, and covered only by peritoneum, were two large fibrous tumours. In the most dense portions of one of these were several yellow-coloured calcareous deposits. Arteries and veins of considerable magnitude were seen ramifying under the peritoneum of the larger tumour, but there was no appearance of a blood-vessel in the central part*.

* Atrophy of the uterus is a rare disease. Mr. Griffiths, of Pimlico, has a remarkable specimen of it in his collection. The muscular coat of the uterus has been entirely removed. There is a fibrous tumour under the peritoneum of the fundus. When the cavity is distended with air, the uterus is diaphanous, like a urinary bladder. The os uteri is healthy. Walter described this disease in 1786, and termed it a membranous uterus.

In the cases which I have now related of calcareous tumours of the uterus, they were accompanied with little or no pain, and the existence of disease was not ascertained before death. In the following example of calcareous concretion, malignant ulceration of the body of the uterus was also present, and the patient sunk after long protracted suffering.

CASE.

In the month of September, 1832, I was requested by Sir Gilbert Blane to see Mrs. B., aged sixty-two, who for many years had suffered from constant sense of weight and uneasiness in the back, loins, and hypogastrium, with almost constant purulent and sanguineous discharge from the vagina. She had been married for many years, but had never become pregnant, and from the age of forty-five, when she ceased to menstruate, she had suffered several severe attacks of uterine hemorrhage. On examination, the hollow of the sacrum was found occupied by a large and hard tumour connected with the posterior part of the uterus. The os uteri had undergone little change, but the peculiar foetor of the discharge, and the constitutional symptoms, led me to suspect the existence of a malignant disease of the body of the uterus. In the course of a few months, after suffering excruciating pain in the region of the uterus, difficulty in passing the urine, with a profuse discharge of thin, offensive fluid from the vagina, several portions of small irregular-shaped concretions escaped from the vagina,

with a temporary relief of the most distressing symptoms.

During the remainder of 1832, Mrs. B. continued to suffer severely from the same symptoms, and she uniformly experienced relief after a calcareous concretion had passed from the vagina, which happened four or five times during that period. In the month of November, 1833, a few days after travelling a distance of eighty miles from the country, she was attacked with rigor, vomiting, exquisite tenderness over the lower part of the abdomen, and other symptoms of peritonitis, and died in forty-eight hours.

I inspected the body the following day with Dr. Webster.

The usual effects of severe peritonitis were seen on laying open the abdomen. The fundus and body of the uterus were extensively disorganized by malignant ulceration; to the posterior part of the body of the uterus was adherent a large fibro-calcareous tumour, which filled up the hollow of the sacrum, and displaced the rectum. The ulceration had extended through the parietes of the uterus to the tumour.

One of the concretions which had been passed during life was analyzed by Dr. Turner, and found to consist entirely of carbonate of lime and animal matter. Dr. Bostock analyzed another concretion passed

by the same patient at a later period, and a portion of the tumour removed with the uterus after death. An interesting account of these and other specimens of uterine calculi, has been laid before the Society by this distinguished chemist.

A case of malignant ulceration of the uterus with a calcareous tumour, in some respects analogous to the preceding case, has been recorded by M. Louis in the second volume of the Memoirs of the Royal Academy of Surgery. With the symptoms and consequences produced by these concretions of the uterus, M. Louis was well acquainted, but it does not appear from any observation contained in his memoir, that he possessed a knowledge of the manner in which these bodies are formed, and the state of chemical science at the time did not enable him to know their composition. The term calculous concretion of the uterus, employed by M. Louis, proves that he knew them to be different from bone.

Schenkius has collected together, from the works of Hippocrates, Vallesius, Salius, Marcellus Donatus, &c., the histories of many wonderful cases of stones discharged from the uterus during life, or found after death.

Michel Morus gives the history of a woman, upwards of forty years of age, who died of a pleurisy, and had suffered for a long time severe pains of the

hypogastrium, for which all remedies had failed to procure relief. On examination, a hardness was felt in the uterus. There escaped from the vagina an acrid discharge, like the washings of putrid flesh. Thirty-two stones were found in the uterus, the smallest of which was the size of an almond. Different folds of the uterus retained them, and some of them were in the Fallopian tubes. He believed these concretions to be of the same nature as bezoards, and he affirms that he saved the lives of several persons by their use. The stones found by Michel Morus in the Fallopian tubes and folds of the uterus, were probably phlebolites or vein-stones, and not fibro-calcareous concretions of the uterus.

With the origin of calcareous concretions of the uterus, pathologists do not appear to have become acquainted till a comparatively recent period. Walter has given representations of these bodies in his *Annotiones Academicæ*, published in 1786, and he states that calculi and polypi are sometimes simultaneously present in the uterus and vagina. It does not appear, however, from this observation, that he was aware of the intimate relation which exists between them, and from an examination of some of the preparations in the Hunterian Museum at Glasgow, I am disposed to believe that Dr. William Hunter was acquainted with the different situations which fibrous tumours occupy in the uterus, and with the various changes which they undergo in the progress of their development.

From an examination of a single specimen of the disease, Dr. Baillie was led to suspect, in 1787, that calcareous concretions of the uterus commence as fibrous tumours. "In the cavity of the uterus", he observes, "a bony mass is sometimes found. When this is the case, I suspect that the hard fleshy tubercle within the cavity of the uterus, such as I have already described, has been converted into bone. This at least had taken place in the only instance which I have known of this disease, for a great part of the tubercle still remained unchanged, and I think it very probable that such a change most frequently happens where the bony tumours are found."

Dr. Baillie refers to Lieutaud for proof of the fact that stones have been found in the cavity of the uterus. "These are described by authors", he adds, "as varying in their appearance, some being of a dark, others of a light colour. They are silent about their nature, and I can say nothing of it from my own knowledge, as it has never occurred to me to see an instance of this disease. Such concretions are probably formed from matter thrown out by the small arteries which open on the internal surface of the uterus, and are in some degree analogous to the concretions formed in some glands of the body."*

Bayle, Bichat, Roux, Breschet, Andral, and other

* The Works of M. Baillie, M.D., by James Wardrop. 1825. Vol. II. p. 331.

writers on the pathology of the uterus, have been fully aware of the fact that fibrous tumours occasionally become calcareous, or as they have inaccurately been termed, bony. Whether all the concretions reported to have been found in the cavity of the uterus, and imbedded in its walls, are formed by deposits in the substance of fibrous tumours, and whether the substance of the uterus itself is ever converted into bone, as several authors affirm to have been the case, it is impossible, in the present imperfect state of our knowledge, satisfactorily to determine. One of the most recent systematic writers on pathology, observes, "that the history of the mode of development of this deposition is not exactly known, and that it is not quite certain whether the ossification originates invariably in the mucous chorion. This, indeed, appears to have taken place in the instance mentioned by Walter, and in such cases of uterine ossification as that recorded by Dr. Caldwell."*

Breschet believes that calculous concretions of the uterus may come from the Fallopian tubes, and that these canals are sometimes obstructed by calculi.

Bayle has described the fibrous tumour of the uterus as fleshy at its commencement, and of a red colour, like muscular fibre, then as becoming carti-

* Craigie's Elements of General and Pathological Anatomy. Edin. 1828, 8vo. p. 732.

laginous, and in the last stage osseous. This may be the case with a few examples of the disease, but I am disposed to think that it is not generally so, and that the greater number of these tumours never exhibit a muscular or fleshy appearance at any period of their existence, but have a fibrous structure equally distinct, when not larger than a pea, and when exceeding in magnitude the head of the human adult.

Sometimes we find only one tumour present in the walls of the uterus, at other times several are met with of different sizes, and not unfrequently they are combined with cysts and tumours of the ovaria. They have no disposition to ulcerate, nor to assume a malignant character, though they are not unfrequently observed in individuals who have cancerous affections of the uterus, bladder, mammæ, liver, and other organs. They have never been observed before the age of puberty, and M. Bayle affirms, that they are most frequently met with in the bodies of those women in whom the physical signs of virginity are present, and that in twenty out of 100 women, taken indiscriminately, after the middle period of life, the fibrous tumour is found imbedded in the walls of the uterus. Of twenty uteri examined by Portal, thirteen contained fibrous tumours, and Dupuytren affirms, that there are few women of a certain age who are without tumours of this description about the uterus. From the observations I have made, I am persuaded that M. Bayle's estimate is correct.

Fibrous tumours are developed either in the cellular membrane under the peritoneal coat of the uterus, or between the layers of its muscular or middle coat, or immediately between its middle and mucous coats. When situated between the peritoneum and muscular coat, they give rise to no irritation, hemorrhage, or derangement, either in the uterine functions, or general health, and their existence even can only be guessed at during life. But when they attain a large size, and occupy a great part of the abdominal cavity, they produce all the injurious consequences of enlarged ovaria, from which, indeed, during life, they are distinguished with difficulty, and death takes place usually from interrupted circulation and long continued pressure on the bladder and other contiguous viscera. Retroversion of the uterus and retention of urine have taken place in the latter stages of the disease.

When situated under the peritoneum of the uterus, fibrous tumours do not prevent impregnation, because they do not interrupt the communication between the vagina and ovaria, but when adherent to the posterior part of the body or neck of the uterus, they sometimes produce fatal consequences both to the mother and child, by impeding its progress through the pelvis. M. Chaussier presented to the School of Medicine, the uterus of a woman who died in labour at the Maternité, in which there was a fibro-cartilaginous tumour as large as the fist, imbedded in the walls of the neck of the uterus. This tumour had formed such an obstacle to labour, that

the head of the child was crushed to pieces in its passage through the pelvis.

In the museum of the London University there is a fibrous tumour as large and nearly as hard as a cricket-ball, which was removed from the body of a woman who had died undelivered. The tumour was situated under the peritoneum, at the posterior and inferior part of the uterus.

Dr. Merriman has cited a case from Van Doeveren, where the expulsion of the child was prevented by the presence of a large polypus of the vagina. He twisted its pedicle and tore away the tumour with his hands, after which a dead child was expelled. Dr. Merriman has also related the history of another case, in which a fibrous tumour of considerable size was connected with the os uteri of a pregnant woman. A ligature was applied around the peduncle, and in a few days the tumour fell off. The general health of the patient improved after the operation, she went to the full time, but the child was still-born.

Dr. Gooch relates a case which occurred in the practice of Mr. Borrett, of Yarmouth, in 1799, and terminated fatally soon after delivery. At the commencement of labour a tumour was discovered in the vagina. After the rupture of the membranes, as the child did not advance, it was delivered by turning, and was born alive. The placenta was expelled spontaneously, but some hours after, a soft round

tumour was found pressing on the os externum. Violent expulsive pain continued for many hours, and twenty-four hours after delivery, a large fleshy tumour, like an inverted uterus, had been forced out of the vagina. She continued to suffer during the whole of this day, and died in the evening. The body was examined the following day. The uterus was contracted, but its mouth was dragged down as low as the external orifice by a tumour, which grew from it by a broad base. It was attached to the posterior part of the mouth of the womb, and some way up the neck was of a livid colour, and weighed three pounds fifteen ounces. The patient had borne her last child before easily and naturally, but some time before her present pregnancy, she looked as large as if seven months with child.

M. Deneux relates the history of a case of fibrous tumour of the uterus expelled into the vagina after an abortion at the fourth month. "The lady, æt. 30, after her second child, observed the abdomen larger than natural, menstruation became irregular, and she had occasional attacks of menorrhagia. She again became pregnant, and miscarried at the fourth month. The after-birth was expelled with difficulty, and the uterus remained larger than usual. Fever followed, with pain of abdomen. After some days a soft fleshy body was perceived at the vulva, which was supposed to be the placenta. The febrile symptoms continued; and this body, which was discovered to be a fibro-cartilaginous tumour, was re-

moved by a ligature applied around its neck, but the patient died. On examining the tumour, which was the size of the fist, its form was found to be irregular, and it was composed of two distinct parts: 1st. An exterior portion in a putrid gangrenous state: 2dly. A central portion, white, fibrous, lamellar, presenting an appearance of little cells, and hard and resisting when cut with the knife. Uterine and abdominal inflammation followed. The tumour had sprung from the inner part of the anterior wall of the uterus. The ligature had been applied to the proper tissue of the uterus. A smooth cavity was found in the anterior wall of the uterus, which was lined with a fine membrane, a portion of which was enclosed in the ligature. Uterus healthy in other respects."*

One of the most remarkable cases of expulsion of a fibrous tumour from the uterus, observed by M. Cruveilhier, occurred in a young woman who was attacked, nineteen days after a difficult labour, with pains exactly like those of labour, and which led him to believe that superfoetation had happened. After suffering so severely for three days that her life was despaired of, the patient passed three flattened bodies of considerable consistence, which were readily recognized to be altered fibrous bodies of the uterus.

When fibrous tumours are formed between the

* *Repert. General d'Anatomie, &c. Tom. VII., 1829.*

† *Anatomie Pathologique, Livraison XIII.*

muscular strata of the uterus, and they attain a large size, its fundus, body, and orifice usually become hypertrophied as during pregnancy, and greatly altered in shape. If situated mid-way between the peritoneal and mucous membranes, they press equally in all directions, as they slowly enlarge, and cause the uterus to project both on the external and internal surfaces. When a thin layer of muscular fibres is interposed between the tumour and peritoneum, the projection is observed only on the corresponding peritoneal surface of the uterus, and the cavity of the organ remains unchanged.

When fibrous tumours are imbedded in the proper tissue of the uterus, women are frequently barren, or if they become pregnant, abortion takes place in consequence of the uterus being incapable of undergoing the necessary development in the latter months of gestation. When the ovum is not prematurely expelled, death may take place in such cases from uterine hemorrhage, soon after delivery. M. Chaussier saw a woman die from flooding, soon after giving birth to a full grown child, and there was a large fibrous tumour in the posterior walls of the uterus. This tumour was not situated so as to present an obstacle to the passage of the child through the pelvis, but soon after delivery it was perceived that the uterus had not the power of contraction. Profuse hemorrhage took place from that part of the uterus in which the tumour was lodged, the flow of blood could not be arrested, and the patient died.

A woman, æt. 42, was delivered by embryotomy of a still-born hydrocephalic child. The liquor amnii amounted to sixteen pints. Profuse uterine hemorrhage followed the extraction of the placenta, and on the third day after delivery, death took place from inflammation of the peritoneal and muscular coats of the uterus. I examined the body, and found a hard fibrous tumour, the size of a hen's egg, imbedded in the muscular coat, where the placenta had adhered to the uterus.

Dr. Outrepont delivered a woman, who died soon after from uterine hemorrhage. Three fibro-cartilaginous tumours were found on dissection in the body of the uterus, the largest of which measured ten inches in the long diameter, and five in the other. They had prevented the development of the fundus uteri in the last months of pregnancy, and the child had only room in the uterus by the excessive dilatation and extreme thinning of the cervix uteri*.

A woman, 40 years of age, was delivered of twins, and died in three days of uterine inflammation. The body was examined by Dr. Chowne, and a large fibrous tumour enclosed in a shell of calcareous matter was found imbedded in the posterior walls of the uterus.

There are no symptoms by which we can positively

* Archives de Medicine, 1830.

determine, during life, the presence of fibrous tumours situated between the muscular strata of the uterus: they may, however, be suspected to exist in those individuals who, being advanced beyond the middle period of life, suffer habitually from leucorrhæal discharge, who menstruate profusely, and have frequent attacks of menorrhagia, with sense of weight and irritation in the region of the uterus and adjacent organs. No alteration of structure can be discovered in the cervical portion of the uterus; but when an examination is made, the uterus is felt larger and heavier than natural. The os uteri is neither irregular, indurated, nor painful on pressure, as it is found to be when affected with malignant disease.

But the fibrous tumour is sometimes developed between the mucous or lining membrane of the uterus and the muscular coat; and as it enlarges, it gradually distends the cavity like an ovum, and pushes before it, through the orifice, that portion of the lining membrane by which it is covered, in a manner somewhat analogous to what takes place in hernia, when the peritoneum is pressed forward by the intestine through the inguinal and crural canals. By the constant and powerful action of the uterus, the tumour is gradually forced into the vagina, where, after the lapse of a longer or shorter period, it undergoes various changes of structure in its covering membrane, peduncle, and central portion. The mucous covering of the tumour sometimes presents no sensible alteration; but more frequently it

becomes highly vascular, thickened and inflamed, or it ulcerates and sloughs, and thus gives rise to a fetid, sanious discharge from the vagina, and to all the other symptoms of malignant disease. In a few rare instances, the tumour has formed adhesions with the vagina.

M. Dupuytren was of opinion that uterine polypi, if abandoned to themselves, ultimately become disorganized by cancer. So long, he observes, as they have a red or white discharge, there is no fœtor, and they are throughout of equal density. If the speculum be introduced, a smooth rose-coloured body is observed ; but if there be a sanious discharge, there is then great fœtor, and when examined we find a soft fungous tumour extending over a surface which bears a relation to the period which has elapsed since the supervention of the symptoms in question. It is also at this time that the constitution begins to suffer in a severe degree, that the skin becomes of a pale yellow, that fever sets in, that emaciation advances, and that the appetite and sleep are lost. There appears to be a decided coincidence, continues M. Dupuytren, between the appearance of gangrene as marked by the fetid, sanious discharge, and the commencement of the cancerous degeneration. This change takes place first in the inferior part of the tumour, which is exposed to the contact of the air ; the pedicle is the part last affected. The accuracy of these statements respecting the cancerous degeneration of uterine polypi, has not been confirmed

by the following or any of the other examples of the disease which have come under my observation.

C A S E.

A woman, forty-four years of age, died in the Saint Marylebone Infirmary, with the usual symptoms of malignant disease of the os uteri. She suffered much for many months before death, from pain in the hypogastrium, and had a profuse sanguineous and purulent discharge from the vagina. She had a sallow complexion, and was much emaciated. I am indebted to my friend and colleague Dr. Sims for the history of the case and the preparation of the parts.

Under the peritoneum of the fundus uteri was a fibrous tumour with a narrow neck, the size of a large walnut. From the inner surface of the fundus uteri there hung by a soft slender root a tumour of a pyriform shape, the greater part of which had passed through the os uteri and filled the upper part of the vagina. The tumour was covered by a thick membrane continued from the lining membrane of the uterus. The membrane covering the stalk was perfectly smooth, but that portion which covered the most depending part of the tumour was soft and partially destroyed by ulceration and sloughing. The central part of the tumour had a dense fibro-cartilaginous structure. Another fibrous tumour, an inch in diameter, was imbedded in the muscular tis-

sue of the uterus near the root of the tumour, filling the vagina, which it strongly compressed. When the uterus was laid open, its cavity contained coagulated blood.

In the following interesting case, for the details of which I am indebted to my colleague Mr. Perry, the fibrous tumour was covered by a capsule, which consisted, not only of the lining membrane of the uterus, but of a layer of muscular tissue. The body was examined after death by Dr. Sims and Mr. Hutchinson, and when they laid open the tumour from the root to the apex, they were both convinced that the substance of the uterus was continued into the tumour and formed its peduncle. The preparation of the uterus with the tumour attached to its cervix, was presented to me by Mr. Perry, the day after the examination of the body took place.

C A S E.

A woman, forty-seven years of age, supposed to be labouring under ascites, and much exhausted by hemorrhage from the uterus, was brought into the Saint Marylebone Infirmary on the 26th November, 1833. A few days after her admission it was ascertained by Dr. Sims and Mr. Perry, under whose care she was placed, that there was a globular-shaped tumour, larger than the foetal head at the end of the ninth month, hanging out of the vagina. The tumour

resembled at first a prolapsed uterus, but when the finger was passed into the vagina, it was found to be connected with the anterior lip of the os uteri by a short root of considerable thickness. The surface of the tumour was of a dark livid colour, and had a sloughing gangrenous appearance in different parts. The woman was so enfeebled in body and mind that she could not communicate a distinct account of her complaints, and the precise period when the tumour appeared externally could not be ascertained.

She stated that she had suffered for several years from profuse discharges of blood from the vagina, and that at different periods a tumour had protruded, which she had always succeeded in returning within the parts by pressure. Dr. Sims and Mr. Perry being of opinion that the removal of the tumour by the ligature afforded her the only chance of relief, Mr. Perry immediately performed the operation with the double canula. She suffered little pain after the ligature was tightened. Twenty-five minims of laudanum were given, and she appeared to be going on well till the evening, when she began to sink, and died in less than twenty-four hours from the time when the ligature was applied.

Dissection. A large cyst, containing several pints of fluid, was found adhering to the left ovarium. There were several small cysts in the right ovarium. The uterus and vagina were healthy. To the anterior part of the cervix a large hard tumour, flattened

on the anterior and posterior surface, was found attached by a thick short peduncle, in which was a slight depression from the ligature. The tumour was invested by a membrane, which was continued from the lining membrane of the uterus. A yellowish-coloured exudation of lymph, which readily peeled off in flakes, partially coated the surface of the tumour, and when pressure was made, blood oozed out from numerous small openings. The root of the tumour was half an inch in length and one inch in diameter, extremely dense, and of a red fleshy appearance, like the muscular coat of the gravid uterus. Numerous large blood-vessels, resembling the sinuses of the gravid uterus, filled with coagula, were seen in the peduncle and in a considerable part of the substance of the tumour. The tumour when first laid open had a dark livid colour like venous blood. Its structure was not uniform. In the most depending part of the tumour was a mass which had the appearance of a common fibrous tumour of the uterus. The root and a great portion of the tumour surrounding this firm nodule had a different structure; they resembled the muscular coat of the uterus, and to all appearance were formed by a continuation of this tissue. Numerous large vessels, resembling the sinuses of the gravid uterus, also traversed this portion of the tumour as well as its root*.

* At a meeting of this Society, held soon after the occurrence of the preceding case, I gave an account of the symptoms before death and the appearances on dissection. Dr. Sims then stated that the neck of the tumour was formed of a portion of

In the 13th Fasciculus of M. Cruveilhier's Pathological Anatomy, there is a representation of a uterus, in the anterior wall of which has been developed a fibrous tumour whose capsule is formed of the lining membrane and a layer of muscular tissue. The cavity of the organ was completely distended by the tumour, which was traversed by large veins, some of which were filled with coagulated blood. In different parts of the tumour there were also small cavities filled with serum, and several great uterine sinuses

the proper or muscular tissue of the uterus, and that the death of the patient was occasioned by the ligature or operation involving a portion of the sensitive tissue of the uterus, which, from its known impatience of injury, occasioned the speedy death of the patient. Dr. Sims also stated that this case, and two or three others which he had seen, explained the original seat of these tumours, which have fleshy necks, and their progressive development. That they were first formed in the muscular tissue of the uterus in the following manner :—1st. If in the centre of the wall of the uterus, they would grow there, and sometimes attain the size of an orange, completely embedded in the muscular tissue. 2dly. If nearer the abdominal surface of the uterus, they would expand into the cavity of the abdomen, and form large tumours, covered by the peritoneum. 3dly. If formed near the cavity of the uterus, they would project into the cavity and protrude before them the muscular tissue, and that the neck of the tumour would be formed by a portion of the substance of the uterus, as before stated. Dr. Sims further observed, that the part of the polypus remote from the wall of the uterus would press upon its covering in the greatest degree, and would occasion its thinning and absorption, and thus account for its not being detected in this part. This view of the formation of these tumours he thought would also explain the cause of the want of success in this operation.

opened upon its surface at the apex, from which the blood had flowed which destroyed the patient. " Cette tumeur", observes M. Cruveilhier, " était ramollie ; les petites masses dont l'agglomération constitue les tumeurs fibreuses, étaient disjointes, et la sérosité remplissait leurs intervalles. La mollesse de la tumeur rendait son enucleation difficile : cependant on saisisait aisément la ligne de démarcation qui séparait le tissu utérin du tissu de la tumeur." " On conçoit," he further remarks, " que la distension de l'espèce de coque qui recouvre le corps fibreux proéminent dans la cavité utérine doive amener quelquefois l'inflammation, l'usure de cette coque, et l'expulsion définitive de la tumeur. Il existe un assez grand nombre d'exemples des ces expulsions spontanées, qui sont toujours accompagnées d'accidents très graves." " Les efforts d'expulsion peuvent avoir pour résultat le déchirement de la couche qui recouvre les tumeurs : et si de gros vaisseaux se trouvent compris dans l'épaisseur de cette couche, une hémorrhagie mortelle peut en être la suite. Ces hémorrhagies peuvent se renouveler aussi souvent qu'il se font les efforts d'expulsion, efforts qui, comme toutes les fonctions utérines, sont soumises à la loi de périodicité." P. 18.

Boivin and Duges also entertain the opinion that uterine polypi are sometimes covered with fleshy fibres, which are continuous with the muscular coat of the uterus. " Le Docteur Breschet assure qu'il a toujours vu les polypes revêtus d'une membrane mince unié, luisante ; dans d'autres cas bien distincte, charnu et d'autant plus mince qu'on se rapprochait davantage

du pedicule si la tumeur était volumineuse, d'autant plus epaisse si la grosseur était mediocre, mais toujours evidemment continue avec les fibres charnus de l'organe même dans lequel le polype a pris naissance, elle était bien manifestement due à la couche interieure de ces fibres respoussée en dedans et entraînée à la surface d'un corps fibreux, dont le siège primitif avait été l'epaisseur même des parois du viscere." *

In the following remarkable case, for the history of which I am indebted to Dr. Merriman and Mr. Cocke of Cleveland Street, the capsule, formed of the lining membrane and a layer of muscular tissue of the uterus which covered the fibrous tumour, had entirely disappeared, not only at the apex, but in the middle of the tumour on one side.

C A S E.

On the 18th November, 1833, Mr. Cocke was called to a patient about the middle period of life, who was in labour with her fifth child. There was considerable flooding, and he could feel a spongy mass adhering to the posterior part of the cervix of the uterus, which he suspected to be the placenta. No part of the child could be felt. After waiting for some time, as the discharge of blood continued, and the uterine contractions, though powerful, had little effect in advancing the child, Mr. Cocke passed his hand into the uterus, and immediately coming in

* *Traité Pratique des Maladies de l'Uterus*, Tom. I., p. 338.

contact with the arm of the infant, he brought down the lower extremities into the vagina and delivered the child. Considerable difficulty was experienced in extracting the head. The placenta was soon expelled, and the quantity of blood subsequently discharged was moderate. The pains continued severe throughout the night, and the following morning a tumour, as large as a child's head, was felt within the uterus, adhering to the posterior and inferior part. Dr. Merriman, on being consulted, was satisfied that the pains were produced by the presence of a large tumour within the substance of the uterus. She died three days after delivery, and the body was inspected by Mr. Cocke.

Seven months after, I had an opportunity of examining the tumour, which I found embedded in the walls of the uterus at the posterior and inferior part, projecting into the cavity, and almost completely filling it up. The root and body of the tumour were covered by the lining membrane and a stratum of muscular fibres of the uterus. At the most depending part there was a circular opening in the capsule, about an inch and a half in diameter, with thin smooth edges, through which aperture a portion of the fibrous tumour projected. On the right side two considerable openings had likewise been formed in the capsule of the tumour. The peduncle of the tumour consisted of the natural muscular tissue of the uterus. Large veins were visible not only in the root, but in the expansion of the mus-

cular fibres over the body of the tumour. The appearances are seen in the preparation of the parts and in the accompanying drawing *.

When a fibrous tumour is formed between the muscular strata, and consequently is covered both by the lining membrane of the uterus and a layer of muscular fibres, the peduncle is proportionably thick and short. A longer continuance of uterine action is also required to force a tumour formed in this situation into the vagina, and the patient not unfrequently dies from irritation and loss of blood before it has been expelled from the cavity of the uterus. The dissections which I have made induce me to believe that it is not on the situation or primitive state of the polypus, as Herbiniaux and Dupuytren have supposed, that the consistence and form of the peduncle depend, but on the quantity of muscular fibres carried before the tumour; and that in those cases where the root of a uterine polypus is thick and short, it will be found to be composed not only of mucous membrane but of muscular coat of the uterus. This account of the formation of uterine polypi will satisfactorily explain why it is unnecessary, as many have supposed, to pass the ligature for the removal of polypi close to the uterus, and it also explains a circumstance pointed out by Clement and Puzos, that the root of the polypus which remains never grows again after the general mass of the tumour has been removed.

* See Plate II.

Fibrous tumours are found attached either to the fundus, body, cervix, or os uteri. Inversion of the unimpregnated uterus is sometimes produced when a large fibrous tumour is developed in the walls of the fundus, and passes through the orifice into the vagina. A case occurred to Dr. William Hunter, in which the patient died by including a portion of the inverted uterus in the ligature.

Dr. Denman saw a young lady who had suffered long from frequent uterine hemorrhages, together with most violent pains recurring in the manner of those of labour. High up in the vagina he discovered a polypus, round which a ligature was with difficulty passed. When he began to tighten the ligature she complained of very severe pain and presently vomited. It was immediately slackened, but on every future attempt to draw it tighter the same symptoms were instantly produced. After many trials he was obliged to desist, leaving the ligature loose round the polypus, *merely to keep up in the mind of the patient some hope of benefit.* The health of this patient was very bad when Dr. Denman first saw her, and in about six weeks from the time of the operation she died. Leave being given to open the body, the uterus was found inverted and the ligature to have passed over the inverted part, which occasioned all the symptoms before mentioned. This polypus, Dr. Denman observes, could not have weighed more than an ounce, and had a very short, if it could be said to have any stem; so that the

uterus could not in this case have been inverted mechanically, but by its own vehement action excited to expel the polypus, which, like any other extraneous and offending body, was a perpetual cause of irritation.

In the Museum of the London University there is a specimen of inverted unimpregnated uterus. A large fibrous tumour, with a thick neck, is seen hanging from its fundus. Mr. Alexander Shaw has informed me, that this was removed from the body of a woman who lay a long time in the cancer-ward of the Middlesex Hospital. She was a woman who had borne several children, and the opinion of those who saw her was, that her uterus had been inverted after delivery.

It occasionally happens, when a fibrous tumour is large and formed under the lining membrane of the cervix of the uterus, that it is suddenly expelled from the vagina by vomiting, or any violent effort, and produces appearances externally which strikingly resemble those observed in cases of chronic inversion of the uterus. The membrane which covers the inverted uterus and the fibrous tumour being the same, and liable to similar changes of structure, without an acquaintance with the previous history of the patient, and a close examination of the symptoms, the diseases might readily be confounded.

Though the facts which have now been stated

clearly demonstrate that the greater number of uterine polypi are fibrous tumours which have been formed under the lining membrane and a stratum of muscular tissue, we are not entitled to conclude, as some have done, that these are the only tumours which make their way from the cavity of the uterus into the vagina, and which are not of a malignant nature. There is a tumour of the fundus or body of the uterus, which grows occasionally from its mucous membrane, or is formed by a morbid change of the mucous membrane itself, which does not acquire a large size, but which seems to be analogous to the common polypus tumour which is formed in the cavities of the nose. It has a broad base and flattened form, and in some cases is largely supplied with blood-vessels. Only two specimens of this disease have been observed by me, and they are now placed on the table of the Society.

There is still another tumour formed under the lining membrane of the uterus, whose structure is peculiar, and differs from any of the preceding. It consists of a congeries of small vesicles or cysts, filled with a clear or yellowish coloured ropy fluid, which cysts are embedded in a soft fibrous substance formed under the lining membrane of the uterus. Five examples of this disease have come under my observation, and in all the tumour was situated under the lining membrane of the fundus, which was very thin and highly vascular. Two of these tumours were adherent to the uterus by a

broad base. One resembled a dried fig, the other was larger than a hen's egg, and distended the cavity of the uterus, the parietes of which were healthy. The appearances presented by these tumours immediately after their removal, have been faithfully represented in the accompanying drawings.

Boivin and Duges have probably described the same disease under the term cellular excrescence of the uterus. "In a considerable number of cases," they observe, "this variety of polypus sprung from the os tincæ, and one whose presence had not been indicated during life had the shape and size of the kernel of a plum-stone; it was soft, of a red-brown colour, streaked with small vessels, and readily separated from the surface, to which it adhered by a slender peduncle. In the same body were observed three similar tumours, one of which was attached to the fundus, the other to the cervix. The exterior of these tumours was continuous with the tissue of the uterus, which tissue formed their covering membrane. Internally, this polypus was also continuous with the substance of the uterus, and was only an extension of this substance in a more cellular and filamentous form." P. 269.

A fourth variety of tumour of the uterus to which the term polypus has also been applied by writers, is produced by a morbid enlargement of the glandulæ or ovula Nabothi. One of these bodies is sometimes

converted into a cyst, as large as a walnut or even a hen's egg, and hangs by a slender peduncle from the cervix or lips of the os uteri. It is smooth and vascular, and contains in some instances a curdly matter, or yellow-coloured viscid fluid. The tumour produces great irritation, and gives rise to copious sanguineous and mucous discharges from the vagina. In a uterus presented to me by the late Mr. John Wood, there are several enlarged glands hanging from the cervix, by long, slender, and flattened stems. One of these glands, the size of a walnut, was tense and smooth, and when cut open was found to contain a yellow curdly matter. I have since met with several other examples of this affection. The appearances in these cases are represented in the accompanying drawings *. Though unacquainted with the nature of the glandular tumour of the os uteri, Herbiniaux has given a description of the appearances it most frequently presents. "There is another species of polypus," he observes, "extremely soft, of which M. Levret has not made mention : it is a little excrescence of the same form as the preceding, but which is always very small ; it arises from a segment of the orifice of the uterus, and either remains within the orifice or hangs a few lines out of it. Often it is not larger than a pea,—sometimes it is the size of the finger, but its stem is usually very large, considering the small size of the tumour."†

* Plate III. Fig. 1, 2, 3.

† *Traité sur divers Accouchemens Laborieux, &c.* Bruxelles, 1782.

Portal states that excrescences analogous to those in the nose and in the mammæ sometimes arise from great congestions of the follicles and lacunæ of the cervix uteri *.

Dr. Gooch has also described polypous tumours of the os and cervix uteri, originating in an enlargement of the mucous glands and follicles of the cervix uteri; but he appears to have been unacquainted with the differences which exist between the structure of the glandular and fibrous polypus of the uterus. "A polypus is sometimes so small," he remarks at p. 287, "that it seems incredible it should occasion the frequent hemorrhagies which attend it. Yet the hemorrhage ceases on the removal of the polypus. I have felt them, as small as a filbert without its shell, growing to the neck or lip of the uterus. They were so small, that on being touched they slipped into the orifice of the uterus, and there remained concealed till the finger was withdrawn and the patient stood up, when they dropped again into the vagina. I saw an elderly woman with a polypus of this size: the day was fixed for its removal, but before it arrived she was using a lotion with a long pewter syringe; it fell away."

Andral and Boivin have described the same disease in the following passages, though the latter appears to have confounded it with the fibrous tumour of the

* Cours d'Anatomie Medicale, Tom. V. p. 523.

uterus. "Au lieu de ces corps fibreux on rencontre quelquefois dans l'épaisseur des parois de la matrice des kystes sereux, de grandeur variable, dont il serait fort difficile d'assigner l'origine. Ces kystes s'observent surtout vers le col de l'organe. Il est des cas où ce col est rempli d'une infinité de ces kystes, qui sont tout très petits, d'égale diamètre et implanté par centaine dans le tissu du col. Quelques uns font une légère saillie du dessous de la muqueuse." *

"Dans l'épaisseur du museau de tanche, nous avons trouvé, en grand nombre, de petits corps blancs, durs comme de cartilage adhérent intimement au tissu environnant et moins gros qu'une lentille; tandis que sur divers autres points de la matrice on en voit égaler le volume d'un œuf, celui du poing et même celui de la tête d'un homme." †

The foregoing observations prove that there are at least four distinct varieties of tumours of the uterus, none of which are malignant in their nature, to which the term polypus has been applied: 1st, the fibrous; 2dly, the follicular or glandular; 3dly, the cystic or vesicular; and 4thly, the mucous tumour of the uterus. To these ought perhaps to be added, that variety of tumour of the uterus which consists of erectile tissue, or of cells and dilated arteries and veins.

* Andral Anatomie Pathologique, Tom. III. p. 691.

† Boivin et Dugès Traité Pratique, p. 314.

Before the middle of the eighteenth century, few facts of any importance had been ascertained respecting the origin and structure of polypus of the uterus. The older writers included under the term polypus all the different tumours of the uterus which have now been described, the greater number of the organic affections of the os and cervix uteri of a malignant nature, and also fleshy moles or ova in a diseased condition. The confusion and obscurity in which the pathology of uterine polypi have been so long involved, may be chiefly attributed to the circumstance that few opportunities have been enjoyed of investigating their structure before it has been destroyed by inflammation, or sloughing produced by natural or artificial causes.

With respect to the treatment of the various tumours which have now been described, I have few observations to offer. Iodine, mercury, and all other remedies have little effect either in arresting their growth or promoting their absorption. Women who have fibrous tumours formed in the walls of the uterus, should avoid mechanical pressure of the hypogastrium, violent bodily exertion, and every other cause which may excite inflammation or a determination of blood to the organs within the pelvis. Where congestion has taken place, it should be removed by local blood-letting, mild cathartics, and anodynes. Profuse uterine hemorrhage should be controlled by rest in the recumbent posture, cold applications to the

hypogastrium, and the internal use of the acetate of lead.

When any of these tumours pass through the os uteri into the vagina, they may be removed by the ligature or by the knife. If the root is soft and slender, the tumour may easily be twisted off by the forceps. In the course of the last twenty years, Dupuytren states, that he has removed two hundred uterine polypi by excision. Hemorrhage has only occurred twice in all the cases, and in both instances it was permanently arrested by the tampon. In eight or ten cases, after the application of the ligature, death took place from the absorption of pus into the system.

Where the root of the tumour is large and vascular, I am of opinion that a ligature should previously be passed around it, at as great a distance from the os uteri as is compatible with the removal of the disease.

Since this paper was read, Dr. Ferguson, Professor of Midwifery, in King's College, London, has communicated to me the following interesting cases and observations.

I.—“A midwife finding that labour did not advance sent for a practitioner, who examined and ascertained, as he thought, that the child's head was jammed in the pelvis, and required for its extraction the aid of the forceps. These he applied, but ineffectually. He left the patient, imagining that as he had succeeded in bringing down the head somewhat lower, the uterine

pains would expel it. And, indeed, some hours afterwards the scalp gave way, the brain was squeezed out, and the putrid full grown child was extracted by the midwife. Two days after this, I was called in to see this person, who had passed a polyp larger than the doubled fist, and who now was dying of peritonitis. She sunk in a few hours, and we found on examination that the polyp had been attached above the cervix uteri; that where the peduncle had adhered was a hole which penetrated into the cavity of the peritoneum, and that probably the inflammation of that membrane had been caused no less by the escape of the discharge through this orifice than by the injuries received during labour.

“It was clear the forceps had been applied to the polyp and not to the head of the child, and that the traction had probably torn the root of the peduncle, which subsequently ulcerated the uterus.”

II.—“A woman who had borne many children, and who had menorrhagia habitually, became pregnant, and the midwife sent for me, not knowing the presentation. I found what first appeared to me to be the scrotum, but a more careful examination proved that the head was presenting though it was still very high. I traced the tumour into the os uteri, near which it hung by a slender stalk. It was soft and compressible, though as large as a hen’s egg. The head descended easily, after I squeezed the tumour against the side of the pelvis. The child was born alive. The placenta came away naturally though tardily, but the most frightful hemorrhage I ever witnessed ensued. She was rescued, however, but with great difficulty. I saw her twelve months after this. The tumour was still there, retaining its usual characters of size and consistence. She would not permit me to remove it.

“I relate this case, as it has always suggested to me the connexion of polyp and hemorrhage; and, secondly, because I think it to be a polyp of the nature of those you describe under the name of disorganized Nabothian glands.”