CASES
OF
CARCINOMA OF THE THYROID
GLAND.

By CÆSAR HAWKINS, Esq.,

SURGEON TO ST. GEORGE'S HOSPITAL.

READ NOVEMBER 28TH, 1843.

CARCINOMATOUS diseases of the thyroid gland are evidently not common from the tenor of the remarks met with in different modern writers. "Malignant diseases," says Mr. King,* "sometimes encroach upon and *very rarely* develop themselves in this body." It is remarked by Dr. Walshe, in his excellent account of cancer in the Cyclopædia of Surgery, "unless as a complication of pre-existing encysted disease, or other form of bronchocele, cancer of the thyroid gland is *singularly rare*." Scarpa entertained the idea that malignant disease of this organ only took place as the consequence of preceding alteration of the parotid gland or other adjacent part.

* Guy's Hospital Reports, vol. i. 432.

I am inclined to believe that the thyroid gland may be more often the seat of primary scirrhus than is usually supposed ; and that I have seen several cases which were in all probability examples of this disease. The following case for instance was, I think, a scirrhus tumour of this body.

Benjamin Young, æt. 61, was admitted into St. George's Hospital under my care, June 28th, 1837, and the note-book observes, that " he had a very large bronchocele, occupying the whole of the anterior part of the neck, extending laterally about an inch and a half beyond each mastoid muscle, and vertically from the os hyoides to the sternum. He first perceived it about six years ago, and his health has generally been good. He has a trifling cough, but the tumour considerably affects his breathing. The tumour is particularly hard and fixed in situation, and is not tender." The effects of the tumour varied from time to time as in all bronchoceles ; and a few days after admission he complained much of weight and pain in the chest, with difficulty in breathing and lying down, and occasionally of noise and throbbing in the neck ; which symptoms were on this and subsequent occasions relieved by a blister to the chest. These chest symptoms, however, were probably in part owing to valvular obstruction of the aorta, aggravated by irregularity of increase in the tumour. He took iodide of potassium internally without advantage ; and the local employment of mercurial ointment and iodide of potassium was tried, but made the tumour more

painful. He continued in the hospital till the end of August, at which time the tumour was larger and harder, and he could scarcely lie down at any time from its pressure on the larynx; his countenance had become very anxious, and he suffered a good deal from pain in the tumour, extending upwards to the temples: and the general irritation was shown by an increase of pulse from 65 to 100. Of his subsequent history I am ignorant, as he left the hospital to go into the country.

In this case, and in others which I believe to have been of the same nature, the age of the patient has been from forty-five to sixty-five when the disease appeared; at which period of life ordinary bronchocele very seldom takes place, and the tumours most likely to present themselves are the serous tumours developed in or near the thyroid gland, or the cartilaginous and osseous degenerations sometimes observed in elderly persons, and lately described by the term *euchondroma*. From the former the distinction is sufficiently obvious, as the existence of fluid is easily ascertained by the finger or the grooved needle. From the latter, the tumour may be more difficult to distinguish, as the hardness is nearly equal in the two cases, but the scirrhus tumour seems to me to be more uniform in its texture, while the other is considerably harder at one part than another, and the surface is smoother and more regular in the scirrhus tumour than in the cartilaginous and bony tissue of the other disease. The scirrhus tumour has appeared also

to be more completely fixed to the larynx, and to the muscles around it, than in any other tumour of this gland, especially in the latter stages, when the cellular tissue is probably beginning to be converted into the same disease. All tumours of the gland necessarily move with the larynx in speaking and swallowing, but in the scirrhus tumour the motion of the larynx is itself interfered with, and therefore it not only rises less freely, but the respiration and deglutition of the patient are more affected than with most other tumours, even when of much larger size. From this cause perhaps in part, and in part from the essential nature of the disease, the patient has an expression of greater suffering and distress in his countenance, than with tumours of other kinds, and although usually of good health when the tumour first appears, he acquires, in the latter stage of the complaint, the usual sallow complexion of one labouring under malignant disease.

But it will naturally be said that these remarks must be only conjectural, unless the final termination of the tumour was known. In the following case, the appearance and symptoms of the disease, on the patient first coming under my care, were as nearly as possible the same as in the case already narrated, and in four others, in which I felt equally convinced of the nature of the tumour ; but I was fortunate enough, from the peculiar manner in which the disease spread, to have an earlier opportunity than I otherwise might have had, of verifying the

diagnosis which I had formed of the tumour on his first admission, while the patient still had the appearance of perfect health. I have thought therefore that the following history might be deserving of the attention of the Society.

Thomas Holder, æt. 50, was admitted into St. George's Hospital May 17th, 1843, having the appearance of perfect health, with a considerable enlargement of the whole thyroid gland, but particularly of the right lobe, which projected upwards more than the left; the tumour was uniformly smooth on the surface, and very firm and solid; it was completely fixed to the larynx, and sufficiently free from attachment to other parts to move with all the motions of the larynx and œsophagus; the skin was unattached and unaltered in colour, and the superficial veins were large. He breathed with some noise, and had a slight cough, but could respire naturally when told to do so. Deglutition was performed with somewhat more difficulty than respiration, although the larynx and trachea were thrown very much to the left side of the neck, nearly two inches perhaps out of the central line. The tumour was free from pain and tenderness. The patient was deaf and dumb, so that a full history could not be obtained at first; but it was learned that the first appearance of the tumour was only about five weeks before his admission.

The nature of the tumour appeared to me evidently to be carcinomatous, and probably scirrhus,

and that it did not admit of surgical relief by operation, in consequence of its intimate connection with the larynx. I determined however to admit him, and observe the effect of iodide of potassium, which was employed both internally and locally; but it was not continued for more than a month, and was intermitted twice in that time, on account of attacks of slight fever, with great pain in the head, which were supposed at first to be owing to the medicine, but were doubtless attributable to the impeded circulation through the brain, as the same attacks occurred after it was discontinued.

The patient at first thought himself better, and the tumour smaller, but the difference was very trifling, and at the end of a month its size had evidently increased. On the 13th of July it was perhaps half as large again as on his admission, and some pain was complained of in the upper part of the tumour. He was observed also to be getting thinner about the middle of June, and had become considerably emaciated before his death, which happened on the 23rd of July.

The difficulty of swallowing was considerable the whole time, but occasionally increased spasmodically, and was then relieved by a small blister to the outside of the larynx; but for the last two or three weeks the dysphagia was constant, and prevented any solid food from being taken, and even liquids were swallowed with difficulty, so that he frequently was obliged to leave the table from threatened suf-

focation in the act of eating, and from vomiting, which sometimes occurred regularly at a certain period after eating, and at other times took place violently during his meal.

Besides the vomiting, he suffered much from pain in the epigastrium and hypochondria, and had tenderness over the stomach, while the respiration seemed little interfered with, so that he could inspire quite deeply without cough or pain, and the sounds were natural on percussion, or when examined by the stethoscope.

About the middle of July, the pain being then much increased, he began to vomit some coagula of blood, but this again lessened while he was taking some lead and opium. For a few hours before his death he had much difficulty of breathing, and became of a livid colour, from this cause.

The following account of the *post mortem* examination is taken from my clinical book.

Upon the anterior surface of the windpipe was a large tumour, which extended from the thyroid cartilage to the sternum. Laterally it projected beyond the sterno-mastoid muscles, the fibres of which, as well as those of the sterno-hyoid, omo-hyoid, and sterno-thyroid muscles of both sides, were expanded over, and partly imbedded in the tumour. The right internal jugular vein, common carotid artery, and pneumogastric nerve, were separated from each other by the pressure of the morbid growth. The vein was closely adherent to the tumour, and its coats, in one place, had been absorbed, and a soft

part of the tumour projected into its interior, and a large clot of blood was firmly attached at this part. The artery was deeply imbedded in the tumour, and the pneumogastric nerve much flattened, and its fibrils separated so as to present a plexiform appearance. (Pl. I. fig. 1.)

The thyroid gland had nearly disappeared, the only part which was left being a small portion of the left lobe intimately joined to the tumour, so as to show that they were originally portions of the same body, and this portion that remained was perfectly natural, and there was a complete line of demarcation between the two structures. The anterior surface of the windpipe was healthy.

Posteriorly the morbid growth had extended to the pharynx and œsophagus, and to the cellular tissue connecting them with the larynx and trachea. The posterior part of the œsophagus was healthy, but the anterior part presented a large, irregular, ulcerated mass, (see Pl. I. fig. 2.*) extending from the aryæno-epiglottic ligaments to the first three or four rings of the trachea, and projecting into the interior of the pharynx and œsophagus, its surface being of a dark green colour, and covered with shreds and portions of sloughs, which were very foetid. The larynx and trachea had been quite pushed over to the left side, forming a curved line, and the right aryæno-epiglottic ligament was much thickened and altered in texture, and immediately

* The preparation was also exhibited to the Society.

below the cricoid cartilage a large ulcerated opening led into the trachea.

Externally, the tumour presented an irregular lobulated appearance, the greater part being situated on the right side; internally, it presented the structure of genuine scirrhus; it was remarkably firm, in some places of a light yellow colour, and in others of a pale red tinge; the variety of scirrhus which it most resembled being the solanoid form; in fact, the section was very like that of a red potatoe.

There were many small encephaloid tubercles at the base of both lungs, and in the cellular tissue under the costal pleura.

The brain was wet, and the bloody puncta large, and the veins and sinuses were gorged with dark-coloured blood.

The stomach was perfectly healthy, and rather small; the liver healthy, with a small serous cyst on the surface of the right lobe; the other viscera were healthy.

The symptoms enumerated, as caused by this tumour, included tenderness and pain in the epigastrium, vomiting after meals, and latterly hæmatemesis, which naturally led my attention to the stomach. During the first few weeks, careful examination detecting no hardness or swelling, and the symptoms not being constantly present, I did not think there was cancer of that organ; but during the last few days, the repeated vomiting of blood, and the great pain and tenderness of the epigastrium,

and in that region only, made me express an opinion that there was most probably cancer of the stomach. The difficulty of swallowing, and the vomiting of blood were, however, satisfactorily accounted for by the ulceration and sloughing of the œsophagus and pharynx ; but whence arose the marked pain and tenderness exactly over the stomach ; which were more complained of by the patient than any other symptom, and existed even at the times when there was a temporary cessation of nausea ?

Looking to the highly expanded state of the fibres of the right pneumogastric nerve, is it not very probable that the symptoms in question depended upon this circumstance, the pain being referred (as with pressure on the spinal nerves) to the part where the nerves are finally distributed below the seat of pressure and irritation, which would in this case be chiefly the pyloric end of the stomach ?

I think I never saw the larynx so much turned out of its natural course by any tumour, as in this case, and dissection showed us also a cancerous degeneration of the side of the rima glottidis, and an ulcerated opening into the trachea ; but yet the difficulty of respiration was never urgent till just before his death, and cough was very little complained of throughout the whole illness. He could always expand the chest freely, and without pain, and the stethoscopic signs showed only bronchitic effusion at the times that the obstruction about the glottis was more marked than usual. In fact, where there has been no pleuritic effusion, and no scirrhus alteration of the

pleura, the little distress occasioned by encephaloid tubercles scattered through the parenchyma of the lungs is often very remarkable.

In this case the scirrhus degeneration occupied the whole of the thyroid gland, except one small portion of the end of the left lobe, which seemed perfectly healthy. In a case published by Velpeau, and consisting probably of encephaloid rather than of scirrhus texture, "The thyroid gland contained," it is said, "about a dozen cancers," all of which were easily separated from the substance of the gland, which was itself perfectly healthy. It is said by Dr. Walshe, who quotes this case, that cancer of the thyroid gland is generally of the scirrhus kind. It is remarkable, however, that in a modern work, which contains more drawings of the various diseases of the thyroid gland than any other work, ("Atlas der Pathologischen Anatomie," by Dr. F. Albers, of Bonn,) while there are several examples of medullary or hæmatoid disease, there is not one, the texture of which looks like scirrhus. There is one called "Carcinoma Glandulæ Thyroidiæ;" it appears however not to be malignant at all, but to consist of the cellular variety of bronchocele, at any rate if malignant, it was not scirrhus. Of the instances of "Fungus Hæmatodes Glandulæ Thyroidiæ," described by this author, one resembles the case I have related, in this respect, that it had increased on each side of the windpipe, so as to adhere apparently to the œsophagus, but the interior of the tube remained quite healthy, and there is a large sloughy

mass in the front of the tumour, instead of the sloughing of the posterior part of the tumour into the œsophagus and pharynx, shown in my own case.

The only case which I have myself seen examined, of fungous or medullary disease of the thyroid gland, was one in which one lobe of this body was diseased, and I place the preparation on the table of the Society to mark the contrast between this variety, and the scirrhus kind of tumour before described. In this case, the patient, a woman, 55 years of age, was admitted into St. George's Hospital, under the late Mr. Rose, on the 9th of November 1827, for a tumour of the thyroid gland, but there was also a tumour on the head, which had been punctured by a surgeon three weeks before her admission, and from which glairy serum, mixed with blood, escaped on the introduction of the probe. To this she paid little attention, and said, when questioned about it, "Oh, Sir, that is only a little tumour." It was the cause of her death, however, about sixty hours after her admission, being fungus hæmatodes of the parietal bone. The preparations of both parts are preserved in our Museum, and the texture of the tumour of the bone was nearly the same as that of the thyroid gland, soft, elastic, dark-coloured, and with a mixture of cells of serum, or bloody fluid, and no part of it is possessed of the dense, firm, and white texture of the scirrhus tissue seen in the other case.

Whichever variety of carcinomatous disease may be the most frequent, it does not appear, from examination of pathological collections, that either of

them is common ; and wherever preserved, the preparations of the two varieties appear to show the same marked contrast, as in the specimens laid before the Society. There are, for example, two preparations of malignant disease in the Museum of the College of Surgeons, one of which came from Mr. Howship's collection, and shows a medullary tumour of large size, entirely surrounding the larynx, soft, spongy, and full of medullary substance. The other is an apparently scirrhus tumour of firmer structure, closely adherent to the larynx and trachea, for several inches, and ulcerating into the tube, which is, in part, converted into scirrhus texture.

APPENDIX TO THE FOREGOING PAPER.

Case of Schirrhous of the Thyroid Gland. By
R. WILSON BROWN, ESQ., *Surgeon to the Bath*
United Hospital. Communicated by SIR B. C.
BRODIE, BART. *Read May 28th, 1844.*

MR. G——, aged 60, of active habits, and of previous good health, began, in December 1842, to suffer from uneasiness about the larynx, accompanied by slight cough and hoarseness, but without much general illness.

A little before Christmas 1842, having caught cold, all the symptoms were aggravated, the cough

became violent, sometimes sharp and stridulous, at others spasmodic, or like the bleating of a goat; the voice feeble, and little above a whisper.

He gained scarcely any relief from treatment, such as from leeching, the use of mercury, blisters, &c.

A hard swelling presented itself in the situation of the left lobe of the thyroid gland. The swelling was not prominent nor well-defined externally, but appeared to extend internally, and press upon the œsophagus, causing great uneasiness, and great difficulty of deglutition. Mr. G. remarked that his throat seemed to be bound, as it were, with an iron hoop. The integuments in the neighbourhood became thickly studded with hard tubercles of a cancerous character.

Internally the disease continued to make progress, the difficulty of deglutition increased, the cough became more violent, convulsive, and almost incessant, with copious muco-purulent expectoration, streaked with scarlet blood. A careful examination of the chest furnished no evidence of any disease of the lungs.

Emaciation gradually took place to a very considerable degree, and eventually death, in June 1843, after much suffering.

Examination of the body thirty-eight hours after death, on June 24th, 1843.—The emaciation was very considerable. Numerous hard tubercles were seated in the skin, covering the throat and part of the chest and abdomen. The seat of the principal disease was in the left lobe of the thyroid gland,

which was somewhat enlarged, and converted into a mass of carcinomatous structure, white, hard as cartilage, and with some gritty particles dispersed through it. The chain of lymphatic glands in the neighbourhood, on both sides, had undergone a similar change, and by these the œsophagus was compressed and reduced in diameter, and the larynx so firmly fixed in its situation, as to prevent any change of its position.

There was not any ulceration of the inner surface of the œsophagus, nor any communication between it and the trachea. There was some thickening of the epiglottis, but not any disease of the trachea.

The lungs had scattered through their substance, and over their surface, several small white tubercles of cartilaginous hardness. The mucous lining of the air-tubes was more vascular than natural, and they contained some frothy mucus. The lungs were otherwise healthy.

A small quantity of serum was effused on each side of the chest, and in the pericardium.

The liver contained many tubercles of the same character as those found in the lungs; they were small, none larger than a filbert, and not softened.

The stomach was much contracted. Numerous glands about its lesser curvature, in the lesser and also the greater omentum, were converted into a carcinomatous structure of the kind already described.

The kidneys and other abdominal organs were healthy.

The cutaneous tubercles on various parts of the

surface of the chest and abdomen were examined, and found to be composed of the same firm, white, carcinomatous structure as that in the thyroid gland and its vicinity.