could breathe spontaneously and all CSF studies became normal in May 2004.

In April 2004, an abdominal computed tomography had shown a 5 cm tumour in the right ovary, which was considered a benign cyst unrelated to the neurological disorder. In June 2004, the patient developed progressive constipation and a bulging appearance of the lower abdomen. Follow-up abdominal computed tomography and MRI showed an enlarged ovarian tumour, with a transverse diameter of 10 cm. On 28 June, resection of the tumour showed an immature teratoma that contained hair follicles, cartilage tissue, glandular structures and cerebral cortex-like tissue with normal appearing neurones. No inflammatory infiltrates were evident in the tumour.

Although her Wechsler Adult Intelligence Scale—Revised score was 84, she recovered and exhibited no limitations in activity of daily living in July 2004. After she was discharged from our hospital in July, she received ambulatory neurocognitive rehabilitation. She refused follow-up Wechsler Adult Intelligence Scale—Revised, but otherwise the cognitive functions and electroencephalogram appeared normal. She returned to her job as a medical resident in April 2005.

Analysis of the patient's serum and CSF showed antibodies, colocalised with EFA6A, which predominantly reacted with the neuropil of the hippocampus and cell membrane of rat hippocampal live neurones (fig 1D).

Discussion

We consider that this patient had definite paraneoplastic encephalitis, with predominant involvement of the limbic system. Accordingly, she developed subacute onset of short-term memory loss, seizures, psychiatric symptoms, CSF pleocytosis, MRI abnormalities in the limbic system and antineuronal antibodies.³ Central nervous system tissue in the teratoma might be a trigger of the immune reaction. Central hypoventilation, skew deviation and anisocoria were observed during the most critical period. These symptoms suggest the involvement of her brain stem.

Previous reports of paraneoplastic encephalitis and ovarian teratoma showed MRI abnormalities in the frontal cortex, cerebellum and brain stem, but no case exhibited the characteristic medial temporal abnormalities observed in our patient.^{2,4} This finding might have resulted from hippocampal inflammation related to the immune response predominantly reacting with hippocampal neurones.

Most patients with PLE and ovarian teratoma improved with resection of the teratomas.^{2,4} We discovered the tumour in our patient 2 weeks after presentation of the encephalitis, but the benign appearance of the tumour and her poor physical status did not prompt for tumour resection. Instead, we started treatment with corticosteroids, plasmapheresis and intravenous immunoglobulin, and she began to improve before the tumour resection. This finding suggests that immunotherapy may provide the improvement needed to undergo the procedure for patients whose poor clinical condition prevents surgery. Furthermore, our patient began to recover faster (4 weeks after admission) than any other reported cases (7–16 weeks).² We presumed that this faster improvement resulted from the combination of immunotherapies. Immunocytochemistry with rat hippocampal live neurones showed the presence of anti-

bodies to antigens present in the neuronal cell membranes and processes and colocalised with EFA6A, as previously reported.² The surface localisation of the autoantigen might be one reason for the effectiveness of these immunotherapies.³

PLE with ovarian teratoma has a better prognosis than that associated with other tumours.² Prompt detection of antibodies that colocalise with EFA6A is useful in predicting a clinical response to immunotherapy and tumour resection and a favourable outcome despite the severity of the disorder.

Acknowledgements

We thank Dr Keiko Tanaka, Department of Neurology, Brain Research Institute, Niigata University, and Dr Yukitoshi Takahashi, Department of Pediatrics, National Epilepsy Center, Shizuoka Institute of Epilepsy and Neurological Disorders, for measurement of the anti-neuronal antibodies.

H Shimazaki, Y Ando, I Nakano

Division of Neurology, Department of Internal Medicine, Jichi Medical University, Tochigi, Japan

J Dalmau

Division of Neuro-Oncology, Department of Neurology, University of Pennsylvania, Pennsylvania, USA

Correspondence to: Dr H Shimazaki, Division of Neurology, Department of Internal Medicine, Jichi Medical University, Tochigi 329-0498, Japan; hshimaza@jichi.ac.jp

Informed consent was obtained from the patient for publication of the features of the case.

doi: 10.1136/jnnp.2006.104513

Funding: This work was supported by a Jichi Medical University Young Investigator Award.

Competing interests: None.

References

- 1 Gultekin SH, Rosenfeld MR, Voltz R, *et al*. Paraneoplastic limbic encephalitis: neurological symptoms, immunological findings and tumour association in 50 patients. *Brain* 2000; **123**:1481–94.
- 2 Vitaliani R, Mason W, Ances B, *et al*. Paraneoplastic encephalitis, psychiatric symptoms, and hypoventilation in ovarian teratoma. *Ann Neurol* 2005; **58**:594–604.
- 3 Graus F, Delattre JY, Antoine JC, *et al*. Recommended diagnostic criteria for paraneoplastic neurological syndromes. *J Neurol Neurosurg Psychiatry* 2004; **75**:1135–40.
- 4 Koide R, Shimizu T, Koike K, *et al*. EFA6A-like antibodies in paraneoplastic encephalitis associated with immature ovarian teratoma: a case report. *J Neuro-Oncol*. In press.
- 5 Darnell RB, Posner JB. A new cause of limbic encephalopathy. *Brain* 2005; **128**:1745–6.

Devic's syndrome-like phenotype associated with thymoma and anti-CV2/CRMP5 antibodies

The case of a patient who presented with a necrotic myelopathy and bilateral optic neuritis in association with a thymoma and circulating anti-CV2/CRMP5 antibodies is reported. This case shows that in some rare instances, a clinical presentation suggestive of a neuromyelitis optica can be of paraneoplastic origin.

A 45-year-old woman with a history of Hashimoto thyroiditis presented with a 4-month history of asthenia and a weight loss of 10 kg. A computed tomography of the chest showed an anterior mediastinal mass suspi-

cious for a thymoma. The mediastinal mass was completely removed by surgery. Histological examination showed a B2-type thymoma with pleural, pericardial and left phrenic local extension. There was no evidence of mediastinal adenopathy or metastasis on computed tomography of the abdomen and pelvis. Treatment with radiation therapy was planned, but 1 month later the patient developed difficulties in walking for over 2 weeks, paraesthesia of the four limbs and bladder dysfunction.

Neurological examination showed a left spastic motor paresis, brisk reflexes and a left Babinski response; proprioceptive sensation was predominantly affected on the left limbs, whereas pain and thermal sensation were affected on the right limbs, suggesting a left cervical Brown-Sequard syndrome. Visual acuity was initially normal. There was no sign of polyneuropathy, and electromyography was normal. The patient did not have fever, and had no signs of systemic disease and no sicca syndrome on general examination. Magnetic resonance imaging (MRI) of the spine showed an enlargement of the cervical cord consecutive to an extensive cervicodorsal (C1 to D7) intramedullary lesion with focal heterogeneous gadolinium enhancement (fig 1A,B). MRI of the brain was normal. The cerebrospinal fluid (CSF) had only an increased protein concentration of 82 g/dl; there was no intrathecal synthesis of IgG, and isoelectric focusing was negative. Polymerase chain reaction of herpes simplex virus (HSV)1 and HSV2 was negative in the CSF on two occasions. The following microbiological tests were also negative; enterovirus, varicella zoster virus, cytomegalovirus, Epstein-Barr virus, Lyme disease, syphilis, HIV and *Mycoplasma pneumoniae*. Anti-double-stranded DNA antibodies and anti-Sjogren's syndrome A and B antibodies were negative.

As an intramedullary metastasis of the thymoma was suspected, a biopsy of the lesion was performed at level C7. On histological examination, the lesions were found to be localised in both white and grey matter. These lesions consisted of a reactive gliosis, with foci of oedema and necrosis with numerous macrophages and some perivascular lymphocytes (fig 1C,D). Bodian luxol coloration showed demyelination. There were no features of vasculitis, nor of viral inclusion or tumour infiltration. This was consistent with a necrotic myelopathy. Serum screening for neuromyelitis optica (NMO) IgG antibodies was negative.¹ Serum screening for onconeural antibodies was negative for anti-Hu, anti-Ri, anti-Yo, and anti-amphiphysin antibodies, but was strongly positive for anti-CV2/CRMP5 antibodies. Immediately after the biopsy, the patient became quadriplegic; this deterioration was probably related to the biopsy. She developed an intestinal subocclusion complicated with aspiration pneumonia. She was treated with high-dose methylprednisolone, but her condition did not improve and she developed a respiratory insufficiency that necessitated artificial ventilation in an intensive care unit. Ten plasma exchanges were also ineffective. At 4 months after the onset of myelopathy, the patient presented a bilateral painless visual loss. Funduscopic examination was normal, and evoked visual potentials showed a bilateral optic neuropathy. The patient finally died of septicæmia 5 months after the onset of myelopathy. No necropsy was performed.

Figure 1 T2-weighted sequence of the spinal cord magnetic resonance imaging showing an extensive cervical intramedullary hypersignal (A). T1 sequence after gadolinium infusion showing focal heterogeneous gadolinium enhancement (B). (C) Haematoxylin, ploxine and saffron staining of the biopsy showing reactive gliosis, oedema, necrosis (*) and microglial infiltration. (D) CD68 immunostaining showing microglial infiltration (brown colouration).

Discussion

Histological evaluation in our patient showed a necrotic myelopathy. It seems unlikely that occlusive vascular disease was implicated because the illness progressed over several weeks. Pathologically, there was no vascular occlusion, and the distribution of the lesions did not correspond to the territory of supply of any of the cord's vessels. There were neither clinical nor biological arguments for an infectious, postinfectious or vasculitic myelitis. In particular, HSV2, which has been reported in association with acute necrotic myelopathy in patients with cancer, was negative in the CSF (polymerase chain reaction). The patient did not receive radiotherapy, thus excluding a radiation myelopathy.

At 4 months after the onset of myelopathy, the patient presented a bilateral optic neuritis

suggesting Devic's syndrome. As in our patient, the myelopathy in Devic's syndrome is usually necrotic. However, the subacute clinical onset, the context of a recently diagnosed malignant thymoma and the presence of anti-CV2/CRMP5 antibodies distinguish our case from that of patients with "classic" Devic's syndrome. These features rather suggest that in our patient this Devic's syndrome-like phenotype was paraneoplastic.² Furthermore, even if it does not exclude Devic's syndrome, we did not detect anti-NMO antibodies.¹ The clinical presentation and histological examination of the myelopathy could have been consistent with a paraneoplastic necrotising myelopathy, but a bilateral optic neuritis has never been described in this clinical entity.³ A case of Devic's syndrome occurring after surgical resection of a thymoma was recently described

by Antoine *et al*.⁴ However, this patient was different because he had myasthenia gravis, developed necrotising myositis in addition to neuromyelitis optica, and had antibodies reacting with the central nervous system and thymic epithelial cells in the serum, but no anti-CV2/CRMP5 antibodies. In fact, the most likely hypothesis in our case is that the Devic's syndrome-like phenotype was related to the presence of anti-CV2/CRMP5 antibodies. Cross *et al*⁵ recently reported three patients who had a myelopathy with optic neuritis, anti-CV2/CRMP5 antibodies and a cancer. Associated cancer was a thyroid papillary carcinoma, a small cell lung cancer and a renal cell cancer. Similar to our case, two patients had an extensive myelopathy and in one of them there was a gadolinium enhancement of the entire thoracic cord suggesting necrotic myelopathy. An autopsy was performed in a third patient (patient 15) with a less extensive myelopathy (spinal MRI abnormalities were limited to a patchy midthoracic T2 hypersignal). Spinal cord pathology showed microglial infiltration, important T cell infiltration but no necrosis. In contrast, the histological evaluation in our patient showed an important microglial infiltration with foci of oedema and necrosis. It shows that in patients with a Devic's syndrome-like phenotype and anti-CV2/CRMP5 antibodies, the myelopathy can be necrotic as in Devic's syndrome. Together with Cross *et al*'s⁵ article, our report also suggests that the presence of anti-CV2/CRMP5 antibodies should be carefully studied in cases of myelopathy of unknown origin.

Acknowledgements

We thank V Rogemond and G Cavillon for their technical assistance.

François Ducray

InsERM U 842, Université Claude Bernard Lyon 1, Hospices Civils de Lyon, Lyon, France

Richard Roos-Weil, Pierre Yves Garcia

Department of Neurology, Centre Hospitalier de Compiègne, Compiègne, France

Juliana Slesari, Olivier Heinzlef

Service de Neurologie, Hôpital de Poissy Saint-Germain, Poissy, France

Denis Chatelain, Philippe Toussaint

Department of Neurosurgery, Centre Hospitalier-Universitaire, Amiens, France

Etienne Roulet

Service de Neurologie, Hôpital Tenon, Paris, France

Jérôme Honnorat

InsERM U 842, Université Claude Bernard Lyon 1, Hospices Civils de Lyon, Lyon, France

Correspondence to: Professeur J Honnorat, Neurologie B, Hôpital Neurologique Pierre Wertheimer, 59 Boulevard Pinel, 69677 Bron cedex, France; jerome.honnorat@chu-lyon.fr

doi: 10.1136/jnnp.2006.097972

Competing interests: None.

References

- 1 Lennon VA, Wingerchuk DM, Kryzer TJ, *et al*. A serum autoantibody marker of neuromyelitis optica: distinction from multiple sclerosis. *Lancet* 2004;**364**:2106–12.
- 2 Graus F, Delattre JY, Antoine JC, *et al*. Recommended diagnostic criteria for paraneoplastic neurological syndromes. *J Neurol Neurosurg Psychiatry* 2004;**75**:1135–40.

- 3 **Ojeda VJ**. Necrotizing myelopathy associated with malignancy. A clinicopathologic study of two cases and literature review. *Cancer* 1984;**53**:1115–23.
- 4 **Antoine JC**, Camdessanché JP, Absi L, *et al*. Devic disease and thymoma with anti-central nervous system and antithymus antibodies. *Neurology* 2004;**62**:978–80.
- 5 **Cross SA**, Salomao DR, Parisi JE, *et al*. Paraneoplastic auto-immune optic neuritis with retinitis defined by CRMP-5-IgG. *Ann Neurol* 2003;**54**:38–50.

HFE H63D polymorphism is increased in patients with amyotrophic lateral sclerosis of Italian origin

A role for metal-mediated oxidative stress in the pathogenesis of amyotrophic lateral sclerosis (ALS) was proposed in 1994 in the first studies of familial ALS mutant superoxide dismutase 1, and interference with iron homeostasis is now postulated.¹ The HFE gene on chromosome 6 is a mean corpuscular haemoglobin class I-like molecule related to iron regulation. Mutations in the coding region cause hereditary haemochromatosis, a common autosomal recessive disorder of iron metabolism that leads to iron overload in adulthood. Recent reports on HFE mutations in ALS showed contradictory results. Two studies described a higher prevalence of the HFE mutations in ALS than in the control group, and one study did not find any difference between the patients with ALS and the control group.^{2–4} We analysed a series of Italian patients with ALS to investigate whether mutations in the HFE gene could represent a risk factor for ALS.

A total of 149 sporadic Italian patients with ALS (mean (standard deviation (SD)) age 59.4 (9.7) years) according to El Escorial criteria for clinically definite or probable ALS were consecutively recruited to this study. Control samples were obtained from 168 healthy people, matched by age (difference of 5 years), sex and ethnic origin (Italian region of birth) to patients with ALS. Patients and controls were informed about the objectives of the study, and informed consent was obtained. The study was approved by the institutional ethics committee. Blood samples were collected, and DNA was purified with a 6100 Nucleic acid Prep Station (ABI PRISM™). Rapid detection of H63D, C282Y and S65C, the three most common mutations in the HFE gene, was performed using the pyrosequencing technique (Pyrosequencing AB, Uppsala, Sweden). This assay is based on a duplex polymerase chain reaction (PCR) in which exons 2 and 4 are amplified together. The exon 2 PCR product is

the exon 4 PCR product for the C282Y mutation. The mutation analysis was subsequently carried out in a triplex assay by means of three pyrosequencing primers in one well. Forward PCR primers for each reaction were modified with biotin at the 5' terminus to capture single-stranded templates from PCR products for pyrosequencing. The following PCR primers were used: for exon 2, ex2-F 5'-ggc tac gtg gat gac cag c-3' and ex2-R 5'-gag ttc ggg gct cca cac-3'; for exon 4 ex4-F 5'-cct ggg gaa gag cag aga t-3' and ex4-R 5'-cag atc aca atg agg ggc tg-3'. The primers used for sequencing were: 5'-gct gtt cgt gtt cta tg-3' for exon 2 and 5'-ggg gaa gag cag aga t-3' for exon 4. The PCRs were performed for 45 cycles, with initial denaturation at 95°C for 10 min, followed by 95°C for 30 s, annealing for 30 s and extension at 72°C for 30 s. The final extension was at 72°C for 5 min. Bound biotinylated single-stranded DNA was prepared following the protocols provided by the manufacturer (PSQ96 sample preparation kit; Pyrosequencing AB). SNP/AQ analysis was performed automatically on a PSQ™96 MA system using enzymes and reagents from an SNP Reagent kit (Pyrosequencing AB).

The group with ALS included 65 women and 84 men (mean (SD) age 61.1 (11.1) years), and the control group included 66 women and 102 men (mean (SD) age 60.7 (9.2) years). Table 1 shows the findings for the three SNPs H63D, C282Y and S65C. Analysis of HFE mutations showed a higher frequency of the mutated allele H63D in patients with ALS than in controls (28.8% v 14.8%; $p = 0.004$). The odds ratio conferred by the presence of the H63D allele was 2.25 (95% confidence interval 1.30 to 3.93). An increased frequency was also found in patients with ALS when all three mutations were combined (33.3% v 17.3%; $p = 0.002$). No significant differences were found between patients with the H63D allele and patients with wild-type HFE gene considering age of onset (63.4 (SD 9.3) v 60.2 (SD 11.9)), sex (22 men and 21 women v 62 men and 44 women), type of onset (33 spinal and 10 bulbar v 80 spinal and 26 bulbar) and disease duration (median survival time, 783 v 993 days).

Our data support the hypothesis that the change in iron metabolism may confer susceptibility to neurodegenerative diseases such as ALS. Our results are consistent with those found in the USA,² and in Ireland and Britain.⁴ Interestingly, the second study reported an odds ratio of 1.85 (95% confidence interval 1.35 to 2.54) for the presence of the heterozygous H63D polymorphism, a value similar to that found in our population. In Europe, the C282Y allele has a north to west frequency-decreasing gradient, with higher frequencies reported in Ireland (28.4%) and Britain (17.4%) and lower

frequencies in Italy (3.2%) and Greece (2.6%). Conversely, the H63D allele has a higher frequency in southern Europe (Spain, 32.3%) and a lower frequency in the Celtic populations (5%).⁵ These marked ethnic differences may explain the negative findings of one study on patients with ALS in Texas, USA,³ in which no matching for ethnic origin was performed.

The possible role of the H63D polymorphism in ALS is unclear. In a human neuronal cell line transfected with genes carrying the HFE mutations, the H63D polymorphism induced a decreased expression of SOD1, α -tubulin and β -actin;² these events can cause a disruption of axonal transport, a factor implicated in ALS pathogenesis. Alternatively, the H63D polymorphism may determine a subclinical increase in intracellular iron, possibly related to neurone oxidative damage. Further studies on the analysis of iron metabolism in patients with ALS are needed to elucidate the role of the H63D polymorphism as a genetic risk factor for sporadic ALS. An alternative possibility could be a linkage disequilibrium of ALS with an unknown gene located near the HFE locus.

Acknowledgements

Funding from Compagnia San Paolo, Torino, and Progetti Finalizzati, Regione Piemonte, 2004–2005 is acknowledged. The Associazione Piemontese per la SLA (APASLA) is gratefully acknowledged for its support.

G Restagno, F Lombardo

Molecular Genetics Unit, Department of Clinical Pathology, Children's Hospital, Torino, Italy

P Ghiglione, A Calvo

Department of Neuroscience, University of Torino, Torino, Italy

E Cocco, I Sbaiz

Molecular Genetics Unit, Department of Clinical Pathology, Children's Hospital, Torino, Italy

R Mutani, A Chiò

Department of Neuroscience, University of Torino, Torino, Italy

Correspondence to: Professor A Chiò, Department of Neuroscience, University of Turin, via Cerasco 15, Torino 10126, Italy; adriano.chio@unito.it

doi: 10.1136/jnnp.2006.092338

Competing interests: None declared.

References

- 1 **Carri MT**, Ferri A, Cozzolino M, *et al*. Neurodegeneration in amyotrophic lateral sclerosis: the role of oxidative stress and altered homeostasis of metals. *Brain Res Bull* 2003;**61**:365–74.
- 2 **Wang X-S**, Lee S, Simmons Z, *et al*. Increased incidence of the HFE mutation in amyotrophic lateral sclerosis and related cellular consequences. *J Neurol Sci* 2004;**227**:27–33.
- 3 **Yen AA**, Simpson EP, Henkel JS, *et al*. HFE mutations are not strongly associated with sporadic ALS. *Neurology* 2004;**62**:1611–12.
- 4 **Goodall EF**, Greenway MJ, van Marion I, *et al*. Association of the H63D polymorphism in the haemochromatosis gene with sporadic ALS. *Neurology* 2005;**65**:934–7.
- 5 **Le Gac G**, Ferec C. The molecular genetics of haemochromatosis. *Eur J Hum Genet* 2005;**13**:1172–85.

Causes of death in multiple system atrophy

Multiple system atrophy (MSA) is a heterogeneous neurodegenerative disorder, with a clinical presentation combining extrapyramidal,

Table 1 Frequency of HFE polymorphisms in patients with amyotrophic lateral sclerosis and controls

Polymorphisms	Patients with ALS		Controls	
	No	%	No	%
H63D/wild type	41	27.5	25	14.8
H63D/H63D	2	1.3	–	–
C282Y/wild type	5	3.3	3	1.8
S65C/wild type	2	1.3	1	0.6
Wild/wild type	99	66.4	139	82.7
Total	149	100	168	100

ALS, amyotrophic lateral sclerosis.