Table 1 Disease characteristics of the six patients

	Case 1	Case 2	Case 3	Case 4	Case 5	Case 6
Sex/age (years)	M/58	F/50	M/29	F/44	M/23	F/58
Age at onset (years)	54	24	21	39	21	8
Family history	—	AD	AD	—	—	—
Presentation	CA	Dystonia	Dystonia	CA	Dystonia	CA/dystonia
Dystonia						
Cranial	+	—	—	+	—	—
Cervical	+	+	+	+	+	+
Axial	—	+	—	—	—	—
Brachial	—	+	—	+/-	—	+
Crural	—	+	—	—	—	—
Cerebellar signs						
Eyes	+	+	+	+	—	+
Speech	+	+	+/-	+	+	+
Gait	+	+	—	+	+/-	+
Arms	+	+	+	+	+	+
Legs	+	+	—	+	+	+
Ataxia disease stage	1	1	1	1	1	1
MRI of the brain						
Cerebellar atrophy	Mild/global	Severe/global (fig 1)	Severe/global	Normal	Moderate/global	Moderate/global
Brainstem	Mild atrophy	Normal	Normal	Normal	Normal	Normal
Basal ganglia	Normal	Normal	Normal	Normal	Normal	Normal
White matter	Normal	Normal	Normal	Normal	Normal	Normal

AD, autosomal dominant; CA, cerebellar ataxia; F, female; M, male; +, positive or present; —, negative or absent; +/- subtle or equivocal.

Ataxia disease stages includes scores 0 (no ataxia), 1 (ataxia, but walking independently), 2 (permanently dependent on walking aids), 3 (permanently dependent on wheelchair) and 4 (death).³

penetrance.⁴ Patients with sporadic late-onset disease, such as case 1, might represent variants of idiopathic late-onset cerebellar ataxias or late-onset primary dystonias.

In conclusion, we present six new patients with a combination of slowly progressive cerebellar ataxia and dystonia, which clinically seems to represent a separate entity, not explained by the known ataxia genes. The wide range in age at onset and variable family history suggest a heterogeneous background, probably including yet unidentified genetic conditions. However, a single causative gene mutation with variable penetrance and expression cannot be excluded.

Acknowledgements

BPCvdW was supported by grants from the Niels Stensen Foundation and the Dr Jan Meerwaldt Foundation and SAS was supported by the Brain Research Trust. We also thank the NHNN diagnostic DNA laboratory. PG and NWW acknowledge the support of EUROSCA.

Barth P C van de Warrenburg

Sobell Department of Motor Neuroscience and Movement Disorders, Institute of Neurology, Queen Square, London, UK; Department of Neurology, Radboud University Nijmegen Medical Centre, Nijmegen, The Netherlands

Paola Giunti

Department of Molecular Neuroscience, Institute of Neurology, Queen Square, London, UK

Susanne A Schneider, Niall P Quinn

Sobell Department of Motor Neuroscience and Movement Disorders, Institute of Neurology, Queen Square, London, UK

Nicholas W Wood

Department of Molecular Neuroscience, Institute of Neurology, Queen Square, London, UK

Kailash P Bhatia

Sobell Department of Motor Neuroscience and Movement Disorders, Institute of Neurology, Queen Square, London, UK

Correspondence to: Professor K P Bhatia, Institute of Neurology, Sobell Department of Motor Neuroscience and Movement Disorders, 7 Queen Square, London WC1N 3BG, UK; k.bhatia@ion.ucl.ac.uk

Informed consent was obtained for publication of the patients' details in this report.

doi: 10.1136/jnnp.2006.105841

Competing interests: None declared.

Supplementary video is available at <http://jnnp.bmj.com/supplemental>

References

- 1 van de Warrenburg BP, Verbeek DS, Piersma SJ, *et al.* Identification of a novel SCA14 mutation in a Dutch autosomal dominant cerebellar ataxia family. *Neurology* 2003;61:1760–5.
- 2 Kuoppamäki M, Giunti P, Quinn N, *et al.* Slowly progressive cerebellar ataxia and cervical dystonia: clinical presentation of a new form of spinocerebellar ataxia? *Mov Disord* 2003;18:200–6.
- 3 Klockgether T, Ludtke R, Kramer B, *et al.* The natural history of degenerative ataxia: a retrospective study in 466 patients. *Brain* 1998;121(Pt 4):589–600.
- 4 Bressman SB, de LD, Brin MF, *et al.* Idiopathic dystonia among Ashkenazi Jews: evidence for autosomal dominant inheritance. *Ann Neurol* 1989;26:612–20.

Paraneoplastic cerebellar degeneration and anti-Yo antibodies in a man with prostatic adenocarcinoma

Paraneoplastic cerebellar degeneration (PCD) is a heterogeneous group of disorders characterised by the subacute onset of a cerebellar syndrome in patients with cancer. Although PCD may occur in patients with a variety of cancers with or without autoantibodies, it is most commonly associated with ovarian or

breast tumours in women seropositive for anti-Yo antibodies. Anti-Yo antibodies bind to protein antigens of 34 and 62 kDa, respectively, found exclusively in the cytoplasm of Purkinje cells. Morphological studies have revealed loss of Purkinje cells accompanied by relatively few or absent inflammatory changes.¹ We present a case of a man with a subacute cerebellar syndrome in whom further diagnostic workup disclosed prostatic adenocarcinoma (PA) and anti-Yo antibodies. Furthermore, the PA showed expression of Yo antigen. Therefore, this case represents simultaneously the first case of PCD associated with PA as well as the fifth case of anti-Yo positive PCD in a male patient.

Case report

A 79-year-old, previously healthy man presented with a 6 month history of disequilibrium that had aggravated in the past 10 days. Neurological examination revealed a cerebellar syndrome with unsteadiness of stance and gait, scanning dysarthria, nystagmus and saccadic dysmetria, and dysidiadochokinesis. Both MRI and CT of the brain were unremarkable. Apart from oligoclonal bands, examination of CSF was normal. Routine laboratory analysis revealed elevated serum levels of prostate specific antigen (174 µg/l; normal <4.2 µg/l). In core biopsies of the prostatic gland, the diagnosis of adenocarcinoma was made. Further laboratory studies detected anti-Yo-antibodies both in the patient's serum and in CSF (fig 1A) by western blot analysis using recombinant Yo transferred to nitrocellulose and blotted against the patient's serum (standard dilution 1:100) and CSF (standard dilution 1:1). Anti-Hu and anti-Ri antibodies were not detected. The clinical status of the patient deteriorated rapidly over the following days and he died a week after admission of acute heart failure. Autopsy revealed prostatic neoplasm, metastasised to the regional lymph nodes and the osseous system. No concurrent secondary tumour was identified after

Figure 1 (A) Immunoblot analysis of recombinant Yo antigen in serum and CSF. Recombinant Yo antigen was transferred to nitrocellulose membrane after sodium dodecyl sulphate-polyacrylamide gel electrophoresis and labelled with the patient's serum (lane 1) and CSF (lane 2). Bound antibodies were visualised by alkaline phosphatase conjugated anti-mouse/human IgG. The bar to the left refers to the migration position of recombinant Yo antigen. (B) Histological section from the patient's cerebellum showing a significant reduction of Purkinje cells (arrows indicating two remaining). Immunohistochemical staining with antibody against neurofilament (clone FNP7, DA2, RMD020.11; Zymed, San Francisco, California, USA); original magnification $\times 10$. (C) Histological section of normal human control cerebellum incubated with biotin labelled anti-Yo and visualised using alkaline streptavidin-biotin complex/alkaline phosphatase/diaminobenzidine showing strong labelling of Purkinje cells (arrows); original magnification $40\times$. (D) Histological section of the patient's prostatic adenocarcinoma incubated with labelled anti-Yo, as described, showing intense labelling of tumour cells; original magnification $40\times$. (E) Histological section of control prostatic adenocarcinoma incubated with labelled anti-Yo, as described, showing no labelling of tumour cells; original magnification $40\times$.

thorough investigation, including extensive histological sampling. Histological examination showed conventional prostatic adenocarcinoma with only focal expression of prostate specific antigen and zones with large numbers ($>25\%$) of single or clustered neuroendocrine cells. A final diagnosis of prostatic adenocarcinoma with focal neuroendocrine differentiation (pT4, pN1, pM1) was made.

Neuropathological examination with sampling of various regions of the brain, including the cerebellum, hippocampus and isocortex, brainstem and central grey matter, was performed after fixation of the brain in buffered formalin for 3 weeks. Histological and immunohistochemical (GFAP, CD45, CD45R0, CD79a, CD68) studies revealed marked reduction of Purkinje cells, especially at the tip of the folia in the upper vermis of the cerebellum with discrete gliosis of the molecular layer (fig 1B). No inflammatory changes were identified.

To prove the causal relationship between the PA and PCD via anti-Yo-antibodies, we tried to demonstrate expression of Yo antigen in the patient's tumour*. Therefore, IgG from serum with known high anti-Yo concentrations was purified by binding to protein A-sepharose

* Unfortunately, at the time of the preparation of the manuscript, fresh tissue was no longer available, so western blotting could not be performed.

followed by elution with sodium citrate (pH 2.5). The resulting fractions of IgG were biotinylated using a biotin labelling kit (No 1418165; Roche, Mannheim, Germany) following the manufacturer's protocol. To show the specificity of the labelled antibodies, $5\mu\text{m}$ thick sections of autaptic, formalin fixed, paraffin embedded normal human cerebellum were dewaxed and boiled in a microwave for 20 min in 10 mM sodium citrate buffer for antigen retrieval. The slides were incubated with the biotin labelled antibodies for 2 h at room temperature followed by incubation with alkaline streptavidin-biotin complex/alkaline phosphatase following the manufacturer's protocol (No 492; Dako, Glostrup, Denmark). Bound antibodies were visualised with 3, 3'-diaminobenzidine. Thus specific and strong immunolabelling of the cytoplasm of Purkinje cells in normal human cerebellum was seen (fig 1C). Consequently, expression of Yo was sought for in the patient's PA accordingly. As a negative control, PA tissue from a patient without neurological symptoms and without anti-Yo antibodies was used. Again, we showed strong and specific immunolabelling in the tumour cells of our patient (fig 1D) whereas the tumour of the negative control yielded no labelling (fig 1E).

Discussion

Anti-Yo positive PCD is usually confined to female patients with gynaecological tumours.

Only very occasionally has anti-Yo positive PCD been reported in male patients: two cases of adenocarcinoma of the oesophagus,^{2,3} and single cases of gastric adenocarcinoma,⁴ parotid adenocarcinoma⁵ and adenocarcinoma of unknown origin.⁶ In contrast, paraneoplastic neurological syndromes associated with PA are a rarity, and PCD in patients with PA has so far never been reported convincingly.

McLoughlin *et al* published a case of a 67-year-old male with "poorly differentiated" carcinoma of the prostate and a subacute cerebellar syndrome, diagnosed by the authors as "paraneoplastic".⁷ However, we believe that it is more likely that this simultaneous presentation was in fact a mere coincidence as (1) the cerebellar syndrome developed after the diagnosis of prostatic carcinoma was made, (2) CSF analysis was unremarkable, (3) no autoantibodies were detected in the patient's serum or CSF and (4) morphological studies were not performed and the patient has not been followed for a long period of time. Furthermore, the "poorly differentiated" carcinoma could well have been a small cell cancer of the prostate and not PA.

Stern and Hulette⁸ reported on a patient with subacute memory loss and ataxia, with mild pleocytosis and elevated protein levels in CSF. Neuropathological examination showed neuronal loss, neuronophagia, "microgliosis" and reactive astrocytosis together with mild perivascular inflammation in parts of the limbic system, consistent with paraneoplastic limbic encephalitis. In the cerebellum, patchy and moderate loss of Purkinje cells, "microgliosis" of the molecular layer and clusters of histiocytes along the Purkinje cell layer were noted, consistent with PCD. However, this patient had small cell cancer of the prostate instead of PA and, furthermore, laboratory analysis failed to detect autoantibodies in the patient's serum or CSF.

As PCD is categorised as a classic paraneoplastic syndrome and further diagnostic workup in our patient disclosed cancer as well as the presence of onconeural antibodies and even expression of Yo antigen by the tumour, this case could reasonably be regarded as fulfilling the criteria for definite paraneoplastic neurological syndromes, as proposed by the members of the Paraneoplastic Neurological Syndrome Euronetwork.⁹ The present case is unusual not only because it represents the first case of PCD associated with PA, but simultaneously the fifth case of anti-Yo positive PCD in a male patient. Hence it underlines the importance of both screening the "paraneoplastic panel" and carefully searching for occult cancer in patients—male or female—presenting with subacute cerebellar syndromes.

Acknowledgments

We thank Ms Ulrike Rumpf for excellent technical assistance.

J Matschke

Institute of Neuropathology, University Medical Centre, Hamburg, Germany

A Kromminga

Institute for Immunology, Clinical Pathology, Molecular Medicine, Hamburg, Germany

A Erbersdobler

Institute of Pathology, University Medical Centre, Hamburg, Germany

K Lamszus

Institute of Neuropathology, University Medical Centre,
Hamburg, Germany

S Anders

Institute of Pathology, University Medical Centre,
Hamburg, Germany

E Köföncü

Neurological Clinic, University Medical Centre,
Hamburg, Germany

Correspondence to: Dr J Matschke, Institute of
Neuropathology, University Medical Centre Hamburg-
Eppendorf, Martinistrasse 52, D-20246 Hamburg,
Germany; matschke@uke.uni-hamburg.de

doi: 10.1136/jnnp.2006.112961

Published Online First 22 December 2006

Competing interests: None.

References

- 1 Shams'ili S, Grefkens J, de Leeuw B, *et al*. Paraneoplastic cerebellar degeneration associated with antineuronal antibodies: analysis of 50 patients. *Brain* 2003;126:1409–18.
- 2 Sutton LJ, Fursdon-Davis CJ, Esiri MM, *et al*. Anti-Yo antibodies and cerebellar degeneration in a man with adenocarcinoma of the esophagus. *Ann Neurol* 2001;49:253–7.
- 3 Xia K, Saltzman JR, Carr-Locke DL. Anti-Yo antibody-mediated paraneoplastic cerebellar degeneration in a man with esophageal adenocarcinoma. *Med Gen Med* 2003;5:18.
- 4 Meglic B, Graus F, Grad A. Anti-Yo-associated paraneoplastic cerebellar degeneration in a man with gastric adenocarcinoma. *J Neurol Sci* 2001;195:135–8.
- 5 Felician O, Renard JL, Vega F, *et al*. Paraneoplastic cerebellar degeneration with anti-Yo antibody in a man. *Neurology* 1995;45:1226–7.
- 6 Krakauer J, Balmaceda C, Gluck JT, *et al*. Anti-Yo-associated paraneoplastic cerebellar degeneration in a man with adenocarcinoma of unknown origin. *Neurology* 1996;46:1486–7.
- 7 McLoughlin J, Gingell JC, Harper G, *et al*. Cerebellar manifestations of prostatic carcinoma. *Postgrad Med J* 1992;68:584–6.
- 8 Stern RC, Hulette CM. Paraneoplastic limbic encephalitis associated with small cell carcinoma of the prostate. *Mod Pathol* 1999;12:814–18.
- 9 Graus F, Delattre JY, Antoine JC, *et al*. Recommended diagnostic criteria for paraneoplastic neurological syndromes. *J Neurol Neurosurg Psychiatry* 2004;75:1135–40.

Dementia and leukoencephalopathy due to lymphomatosis cerebri

Lymphomatosis cerebri (LC) is a rare variant of primary central nervous system lymphoma (PCNSL). Clinically, the disease typically presents with a rapidly progressive dementia and unsteadiness of gait. Its presentation on cerebral MRI, which is characterised by diffuse leukoencephalopathy without contrast enhancement, often causes diagnostic confusion¹ with suspected diagnoses ranging from Binswanger's disease to leukoencephalopathy or encephalomyelitis.

Here we report a patient with subacute dementia and diffuse bilateral white matter changes in the cerebral hemispheres and additional involvement of the brainstem, basal ganglia and thalamus on MRI. Initially, she was considered to suffer from an autoimmune encephalitis, transiently responded to

immunosuppression but then developed multiple solid appearing cerebral lymphomas.

Case report

This 65-year-old woman had a history of insidious personality changes and forgetfulness of several months. Shortly before admission she developed double vision. Neurological examination revealed bilateral sixth nerve palsy. Neuropsychological examination showed mild dementia. CSF examination showed an increased protein level (931 mg/L; normal 150–400) and cell count (18/μl; normal <4). On cytology, many activated lymphocytes but no malignant cells were found. There was IgM synthesis and positive oligoclonal bands in

the CSF. Serum levels of β2-microglobulin were normal (2.0 mg/l) and CSF levels were increased to 3.37 mg/l (normal <3). Antinuclear antibody titres (1:320; normal <1:80) and circulating immune complexes containing IgM (210 μg/ml; normal 15–114) were elevated. Extensive microbiological serology, including testing for HIV, was unremarkable. PCR for *Tropheryma whippelii* was negative, as was PCR for *Mycobacterium tuberculosis*. No virus or bacteria, including mycobacteria, were cultured from the CSF. Cerebral MRI showed diffuse symmetrical pathological signal hyperintensities in the lateral aspects of both putamina and both caudate nuclei and the bilateral frontal white matter, with relative sparing of U-fibres (fig 1A, B). None of the

Figure 1 (A–C) Cerebral MRI at admission. (A, B) Fluid attenuated inversion recovery (FLAIR) weighted images showing pathological signal hyperintensities in the deep white matter of both hemispheres and the basal ganglia. (C) T1 weighted image with gadolinium (GD) application at the corresponding plane to B showing absence of contrast enhancement in the parenchyma with only an enhanced vessel. (D–F) Cerebral MRI 18 days later revealed a prominent increase in signal hyperintensities in FLAIR weighted images (D, E) now also involving the thalamus but still lacking contrast enhancement after GD (F). (G, H) Statistical parametric mapping analysis of cerebral fluorodeoxy-glucose-positron emission tomography (FDG-PET) at the planes corresponding to D and E showing increased FDG uptake (red) in the deep cerebral white matter. (I) Cerebral T1 weighted MRI 4 months later showing multiple nodular contrast enhancing lesions.