Table Representative nerve conduction studies of patients 1–3

	Patient 1			Patient 2			Patient 3		
	Latency (ms)	Amplitude (mV)	C vel (m/s)	Latency	Amplitude	Velocity	Latency	Amplitude	Velocity
Motor nerve conduction studies									
Peroneal nerve ankle-EDB		Absent		7	1.6		7.1	0.7	
Peroneal nerve knee-EDB					0.5	27		0.7	19
Tibial nerve ankle-AH		Absent		7.3	2.2		6	8.2	
Tibial nerve knee-AH					1.9	25		4.9	21
Median nerve wrist-APB	6.4	8.4		5.7	5.6		5.2	12.4	
Median nerve elbow-APB		4.6	21		3.7	32		5.6	25
Sensory NCS									
Sural nerve		Absent			Absent			Absent	
Median nerve wrist-index		Absent		5.2	13			Absent	
Median nerve elbow-index				10.4	3	42			

AH, abductor hallucis; APB, abductor pollicis brevis; C vel, conduction velocity; EDB, extensor digitorum brevis; NCS, nerve conduction studies.

NCS studies showed relatively preserved motor amplitudes with reduced conduction velocities (table). Sensory responses were absent in both the upper and lower limbs. His symptoms progressed over 5 years and he developed an external ophthalmoplegia, bilateral ptosis, proximal limb weakness and mild limb ataxia. These new neurological signs prompted a needle muscle biopsy, which showed numerous COX negative ragged-red fibres on the combined cytochrome c oxidase succinate dehydrogenase stain, confirming the diagnosis of a mitochondrial myopathy. Unfortunately, DNA available was insufficient to perform a Southern blot to confirm an mtDNA genetic abnormality.

Discussion

Mitochondrial diseases can affect multiple organs, but have a predilection for organs with high-energy requirements such as the muscles or brain.¹ Thus they may present in a wide variety of ways, including myopathy, external ophthalmoplegia, diabetes mellitus, short stature or hearing loss. Although peripheral neuropathy is often a component of mitochondrial disease,^{2,3} it is rarely the presenting symptom. Previous reports have not emphasised the possible confusion between mitochondrial disease and CMT disease when peripheral neuropathy is the dominant feature.

Peripheral neuropathy in mitochondrial disease may occur in up to 50% of affected patients² but is often subclinical.³ In most studies, the axonal form predominates, although demyelinating forms have also been reported.^{3,4} The symptoms of MNGIE syndrome have also been reported to mimic those of CIDP.⁴ Over 30% of the patients seen in our neurogenetics clinic with genetically or biopsy-proved mitochondrial myopathy have a peripheral neuropathy on NCS, but <5% of these are demyelinating. The three patients discussed here are the only patients to initially present with the primary problem of a neuropathy, with distal weakness, pes cavus and clawed toes. The severity of sensorimotor peripheral neuropathy with slowed conduction velocities led clinicians to consider the possibility of CMT. On electrophysiological grounds alone, it would be difficult to distinguish CMT from mitochondrial disease, as both of these conditions have axonal and demyelinating forms. Clinical features distinguishing our patients from patients with CMT included external

ophthalmoplegia and bilateral partial ptosis. In the two patients with MNGIE syndrome, severe gastrointestinal involvement prompted further investigation. Although not specific for mitochondrial disease, these clinical features are exceedingly rare manifestations of CMT disease, and we note that recently there has been a case report of a Taiwanese family with probable CMT and ptosis.⁵ These ophthalmological and gastrointestinal clinical features were not initially present in our patients and thus it is important to follow up patients with undiagnosed peripheral neuropathies and negative genetic studies for CMT, particularly if the conduction velocities are slowed. On development of these features, a measurement of thymidine phosphorylase activity in the buffy coat and plasma thymidine levels for MNGIE syndrome, genetic studies and muscle biopsies confirmed the diagnosis of mitochondrial disease.

In conclusion, young patients with CMT-like peripheral neuropathies and negative CMT genetic studies should undergo close clinical follow-up. The development of any atypical clinical features, such as ptosis, external ophthalmoplegia or prominent gastrointestinal symptoms, should prompt consideration of the diagnosis of a mitochondrial cytopathy. In these cases, a muscle biopsy or relevant genetic studies should be carried out.

M Needham

Department of Neurology, Royal North Shore Hospital and University of Sydney, Sydney, New South Wales, Australia

J Duley

Chemical Pathology, Mater Health Services, Brisbane, Australia

S Hammond

Department of Neurology, Orange Base Hospital, Sydney, New South Wales, Australia

G.K. Herkes

Department of Neurology, Royal North Shore Hospital and University of Sydney, New South Wales, Australia

M Hirano

Department of Neurology, College of Physicians and Surgeons, Columbia University, New York City, New York, USA

C M Sue

Department of Neurology, Royal North Shore Hospital and University of Sydney, Sydney, New South Wales, Australia

Correspondence to: C M Sue, Department of Neurogenetics, Clinic 4, Royal North Shore Hospital, Reserve Rd, St Leonards, Sydney, New South Wales 2065, Australia; csue@med.usyd.edu.au

doi: 10.1136/jnnp.2005.078857

Funding: Supported by the Australian National Health & Medical Research Council, Canberra, Australia (Grant No 163404).

Competing interests: None.

Informed consent was obtained for publication of the patients' details described in this report.

References

- 1 DiMauro S, Bonilla E, Zeviani M, *et al.* Mitochondrial myopathies. *Ann Neurol* 1985;17:521–38.
- 2 Yiannikas C, McLeod JG, Pollard JD, *et al.* Peripheral neuropathy associated with mitochondrial myopathy. *Ann Neurol* 1986;20:249–57.
- 3 Mizusawa H, Ohkoshi N, Watanabe M, *et al.* Peripheral neuropathy of mitochondrial myopathies. *Rev Neurol* 1991;147:501–7.
- 4 Bedlack RS, Yu T, Hammans S, *et al.* MNGIE neuropathy: five cases mimicking chronic inflammatory demyelinating polyneuropathy. *Muscle Nerve* 2004;29:364–8.
- 5 Wu T, Wang HL, Chu CC, *et al.* Clinical and electrophysiological studies of a family with probable x-linked dominant Charcot-Marie-Tooth neuropathy and ptosis. *Chang Gung Med J* 2004;27:489–500.

Isolated shoulder palsy due to cortical infarction: localisation and electrophysiological correlates of recovery

Isolated pure motor involvement of the shoulder and arm muscles is extremely rare after stroke, and up to now has been documented by magnetic resonance imaging (MRI) in only one patient.¹

The corticospinal system is known to exert a greater influence over distal than proximal upper limb muscles, and the mechanisms that induce better recovery from stroke of proximal muscles are debated. The contribution of ipsilateral corticospinal fibres from the unaffected hemisphere, or of corticoreticulospinal projections, has been hypothesised.^{2,3}

A 65-year-old right-handed man awoke unable to abduct the right arm. Examination showed a strength of 0/5 (Medical Research

Council scale) in the right deltoid, 2/5 in the biceps and 4/5 in the triceps. Strength in the remaining upper and lower limb muscles as well as the sensory examination were completely normal. Tendon jerks were normal except in the right biceps, where they were reduced. Electromyography showed no voluntary activity in the right deltoid muscle, and a reduced recruitment pattern with motor units of low firing frequency in the biceps brachii. A computed tomography scan showed a small hypodense lesion in the left precentral gyrus. Two months later, the patient was able to abduct the arm against gravity and the strength in the biceps was 4/5. Biceps tendon jerk was hyperactive. At 7 months, the patient was able to hold the right arm abducted against resistance (4+/5).

Standard and functional magnetic resonance imaging (fMRI) was performed 2 months after stroke. fMRI was carried out using the blood oxygenation level-dependent contrast technique. To explore the cortical activity of the primary and secondary motor cortices, a self-paced finger-tapping task at a frequency of about 1 Hz was used. The experimental paradigm was a block design alternating a task condition (right or left finger tapping) with a control state having the same duration (30 s; 3 s for each fMRI volume) and consisting of complete motor and mental relaxation. For each run, 135 volumes were acquired. Raw data were analysed by means of the Brain Voyager 4.9 software (Brain Innovation, Maastricht, The Netherlands) and statistical analysis was performed using the general

linear model. Statistical maps were thresholded at $p < 0.001$ at the voxel level and a cluster size of at least four voxels was required. Functional volumes were co-registered with a high-resolution structural T2-weighted sequence where the ischaemic lesion was detectable. MRI showed a 3 mm cortical-subcortical ischaemic lesion in the superior genu of the left precentral gyrus medially to the knob-like structure corresponding to the motor area of the hand activated by the right finger-tapping task (fig 1A).

Magnetic stimulation was performed two and seven months after stroke using a high-power Magstim 200 magnetic stimulator (Magstim Co., Whitland, Dyfed, UK) connected to a figure-of-eight coil. Motor-evoked potentials (MEPs) were recorded bilaterally

Figure 1 (A) Axial T2-weighted image with superimposed functional magnetic resonance imaging activations obtained during a right finger-tapping task. Standard T2-weighted sequence shows an ischaemic lesion (arrow) in the superior genu of the left precentral gyrus medially to the structure activated by the finger tapping. An activated region was also present in the mesial surface of the left frontal lobe corresponding to the supplementary motor area. Image is shown using the right-left radiological convention. (B) Cortical representation maps of the deltoid, biceps and first dorsal interosseus (FDI) muscles 2 months after stroke. The stimulus intensity was set at 200% of resting motor threshold, defined as the minimum stimulus intensity that produced a liminal motor-evoked potential (MEP) (about 50 μ V in 50% of 10 trials) at rest in the FDI muscle. At this intensity, a MEP of about 3 mV was evoked in the relaxed FDI and of about 0.5 mV in the proximal muscles when stimulating the area around the optimal scalp position for each muscle. The responses evoked by three stimuli at each scalp site were averaged. The area of each map was taken as the number of scalp points at which a response was evoked, and the volume of each map was taken as the sum of the averaged MEPs recorded at all scalp sites at which a response was evoked. Each point of the scalp surface is identified by two coordinates on an x-y plane; the x axis lies on the frontal plane and the y axis lies on the longitudinal plane. The white cross indicates Cz. Positive and negative points are located on the right and left hemisphere, respectively, on the x-axis, and anterior and posterior to the line of Cz on the y axis. Cortical maps represent the motor-evoked potentials after stimulation of each point of the scalp recorded on the contralateral muscle. The amplitude of the response on each point is represented on the z axis. There are no MEPs in the right deltoid.

and simultaneously from the deltoid, biceps brachii and FDI muscles. To investigate the presence of contralateral and ipsilateral responses, we stimulated the optimal scalp position for each of the three muscles studied in the affected and unaffected hemisphere using a stimulus intensity of 100% of maximum stimulator output during tonic activation at about 20% of maximum voluntary contraction. To increase the possibility of detecting MEPs in paretic muscles, cortical stimulation was also performed with a non-focal circular coil centred over the vertex; moreover, the scalp was mapped systematically using a figure-of-eight coil and a standard TMS protocol evaluating the cortical representation of each tested muscle together with the deltoid muscle.⁴ Cortical stimulation with a non-focal circular coil centred over the vertex was performed at an intensity corresponding to the maximum stimulator output, with a clockwise inducing current flow as viewed from above for the right motor cortex and vice versa for the left motor cortex, both at rest and during voluntary contraction of the tested muscles at approximately 20% of the maximum.

Two months after stroke, maximal TMS of the contralateral and ipsilateral motor cortex during voluntary contraction did not evoke a MEP in the right deltoid either with a focal or a non-focal coil. TMS mapping showed no representation of the right deltoid in the left motor cortex and adjacent areas (fig 1B). A large-amplitude MEP was recorded at rest in the left deltoid after stimulation of the opposite motor cortex (fig 1B); this might be due to an increase in the excitability of the right, intact motor cortex consequent to an imbalance in interhemispheric inhibition.⁵ Seven months after stroke, when the patient had almost completely recovered, maximal TMS of the contralateral and ipsilateral motor cortex still did not evoke a MEP in the right deltoid either with a focal or a non-focal coil.

Recently, fMRI studies allowed us to correlate the somatotopy of the primary motor cortex to anatomical landmarks. A study indicated that the segment of the precentral gyrus containing the hand function is a knob-like structure corresponding to the middle knee of the central sulcus shaped like an inverted omega.⁶ It has been difficult to obtain a "pure" fMRI cortical activation of the shoulder muscles because the movement also propagates to the distal muscles during a shoulder motor task. Crafton *et al*⁷ demonstrated that 63% of areas activated during shoulder movement were also activated during hand motor tasks, but they were able to individuate a centre of activation for the shoulder with Talairach's coordinates 23, -29 and 55. These coordinates corresponded with the area where the ischaemic lesion was localised in the patient we report. This area was medial to the hand cortical activation area obtained by a finger-tapping task as expected from the classical human primary motor cortex map.

The patient we report recovered almost completely, but we could not record MEPs in the right deltoid muscle after maximal stimulation of the contralateral motor cortex, and there was no representation of the right deltoid in the left motor cortex and adjacent areas. MEPs were not evoked also by maximal stimulation of the ipsilateral motor cortex. Few other patients who recovered voluntary activity in proximal muscles but had no motor responses to contralateral and ipsilateral TMS

have been reported.² Recovery of proximal muscles in these cases may be mediated by elements other than the fast corticospinal neurones responsible for MEP generation. Slowly conducting contralateral or ipsilateral corticospinal projections or polysynaptic pathways, such as corticobulbospinal, corticoreticulospinal or corticopropriospinal projections, which cannot be probed by TMS, may compensate the reduction or even the complete loss of the fast corticospinal inputs to the motoneurones of proximal muscles after stroke.

A Uncini, C M Caporale

Department of Oncology and Neurosciences, "G. d'Annunzio" University and the Aging Research Center, Ce.SI, "G. d'Annunzio" University Foundation, Chieti-Pescara, Italy

M Caulo, A Ferretti, A Tartaro

Department of Clinical Sciences and Bio-imaging "G. d'Annunzio" University and ITAB-Institute for Advanced Biomedical Technologies, "G. d'Annunzio" University Foundation, Chieti-Pescara, Italy

F Ranieri, V Di Lazzaro

The Neurological Institute, Catholic University, Rome, Italy

Correspondence to: A Uncini, Clinica Neurologica, Ospedale "SS. Annunziata", via dei Vestini, 66013 Chieti, Italy; uncini@unich.it

doi: 10.1136/jnnp.2005.087346

Competing interests: None declared.

Informed consent was obtained for publication of the patient's details described in this report.

References

- 1 Komatsu K, Fukutake T, Hattori T. Isolated shoulder paresis caused by a small cortical infarction. *Neurology* 2003;**61**:1457.
- 2 Turton A, Wroe S, Trepte N, *et al*. Contralateral and ipsilateral EMG responses to cranial magnetic stimulation during recovery of arm and hand function after stroke. *Electroencephalogr Clin Neurophysiol* 1996;**101**:316–28.
- 3 Alagona G, Delvaux V, Gérard P, *et al*. Ipsilateral motor responses to focal transcranial magnetic stimulation in healthy subjects and acute-stroke patients. *Stroke* 2001;**32**:1304–9.
- 4 Uy J, Ridding MC, Miles TS. Stability of maps of human motor cortex made with transcranial magnetic stimulation. *Brain Topogr* 2002;**14**:293–97.
- 5 Liepert J, Hamzei F, Weiller C. Motor cortex disinhibition of the unaffected hemisphere after acute stroke. *Muscle Nerve* 2000;**23**:1761–3.
- 6 Yousry TA, Schmid UD, Alkhadhi H, *et al*. Localization of the motor hand area to a knob on the precentral gyrus. A new landmark. *Brain* 1997;**120**:141–57.
- 7 Crafton KR, Mark AN, Cramer SC. Improved understanding of cortical injury by incorporating measures of functional anatomy. *Brain* 2003;**126**:1650–9.

Rapid improvement of diffusion-weighted imaging abnormalities after glucose infusion in hypoglycaemic coma

Diffusion-weighted imaging (DWI) may detect hyperintense lesions in patients with transient hypoglycaemia-induced hemiparesis or coma, which are completely reversible after glucose

infusion.^{1–3} In vivo animal studies have documented the visualisation of such hypoglycaemia-induced changes of signal intensity and the reversal by glucose intake in detail.⁴ However, the time necessary for hyperintense lesions on DWI to disappear after glucose infusion in humans is still unclear. We treated a patient in a hypoglycaemic coma with DWI abnormalities in the splenium of the corpus callosum and the bilateral posterior limbs of the internal capsules, which were drastically improved 2 h after glucose infusion.

A 54-year-old woman was taken to an emergency department after being found in an unresponsive state. The patient had had type 2 diabetes mellitus, hypertension and hyperlipidaemia for 7 years, controlled by oral drugs. On arrival, she was comatose (Glasgow Coma Scale 4; E1, V1, M2). Her pupils exhibited isocoria (2 mm/2 mm) and were reactive to light. Tetraparesis and decerebrate rigidity to pain was evident. Her blood pressure was 91/40 mm Hg and her heart rate was 88 beats/min. She had spontaneous respiration and a normal temperature. Magnetic resonance imaging (MRI) and magnetic resonance angiography were carried out immediately, as we considered the possibility of brain stem infarction. DWI showed hyperintense lesions in the splenium of the corpus callosum and the bilateral posterior limbs of the internal capsules (fig 1A). T2-weighted and fluid-attenuated inversion recovery sequences showed no abnormal lesions, and magnetic resonance angiography showed no major abnormalities, including basilar artery occlusion. Laboratory examinations showed no metabolic abnormalities except for severe hypoglycaemia with a blood glucose concentration of 1.8 mmol/l. The patient was immediately given 40 ml of 50% glucose infusion; she recovered consciousness shortly afterwards. The abnormal hyperintense lesions had almost disappeared on DWI obtained 2 h after glucose infusion (fig 1B), and had completely disappeared 2 days later (fig 1C). Blood glucose concentrations after glucose infusion and 2 days later were 13.9 and 6.5 mmol/l, respectively. The final cognitive function was completely normal in her daily life. The patient's hypoglycaemia was due to low dietary intake and irregular medication in the 4 days prior to admission. The duration of hypoglycaemia was not apparent.

Follow-up MRI has been carried out between 8 h and 10 days in previous cases,^{1–3} hence the duration of DWI abnormalities cannot be exactly known. But in reality, the hyperintense lesions on DWI had almost disappeared in only 2 h in our patient. The in vivo animal study of hypoglycaemia found complete apparent diffusion coefficient (ADC) normalisation except in the periventricular regions by 10 min after glucose infusion.⁴ Glucose deprivation is widely assumed to lead to severe brain energy failure, reduction of cell membrane ionic pump activity and consequent shift of cerebral water from the extracellular space to the intracellular space. The fast recovery of the clinical symptoms and normalisation of imaging abnormalities presumably resulted from the immediate intravenous glucose supplementation. Although, in our patient, follow-up MRI was carried out in 2 h after glucose infusion and rapid improvement in DWI abnormalities was detected by chance, there is a possibility that the actual duration of DWI abnormalities in previous case reports also might be short, as in our patient.