ON THE

PATHOLOGY OF ACUTE PERIOSTITIS.

BY

CLINTON T. DENT, F.R.C.S.,
ASSISTANT SURGEON TO ST. GEORGE'S HOSPITAL.

(Received May 24th—Read June 14th, 1881.)

THE geographical distribution of disease is a subject which has received its due share of consideration, but a branch of this wider question, viz.:—the variations in different countries and among different nationalities of diseases everywhere common, has not perhaps met with the attention which it deserves. I am inclined to think that the disease treated of in the present paper may possibly furnish evidence that given pathological changes may vary in their progress and results according to the nationality of the patient in whom they are manifest; this is the case with other diseases. Among the Germans and the Turks, for instance, there is a remarkable susceptibility to carbolic acid poisoning.¹

Sufficient proof that the pathology of acute periostitis has been matter for difference of opinion is afforded by its varied nomenclature. Thus, it has received the name of "acute periostitis," "osteomyelitis" (Billroth),² "péri-

¹ Billroth's 'Chir. Klin.,' 1876, p. 39.

² 'Surg. Path. and Therapeutics,' London, 1877, p. 289.

ostite phlegmoneuse" (Giraldés), "ostéite phlegmoneuse diffuse,"¹ "abcès sous-périostique aigu,"² "ostéite épiphysaire," &c. The exact relation which acute periostitis and osteomyelitis bear to each other, as well as to the articular affections with which they are so often associated, have long been matters of dispute; so also have their effects with regard to the extent of necrosis which usually follows. Among French surgeons the opinion originating with Chassaignac is very generally held, that total necrosis of the shaft will not take place as a sequel of acute periostitis alone, but that osteomyelitis must have coexisted. German writers³ seem also to have adopted the same conclusion very generally. On this question there seems to be some difference of opinion between continental surgeons and those in this country; but the difference is less than it might appear at first sight, and can be well estimated from a critical *résumé* in the 'London Medical Record,'⁴ by Mr. Holmes, on the subperiosteal removal of the entire diaphysis of the bone for diffuse phlegmonous periostitis. Two points may be gathered from this paper:—(1) That osteomyelitis need not necessarily be associated with acute periostitis, and (2) that the entire diaphysis of a bone may be so completely necrosed, as a result of periostitis unaccompanied by osteomyelitis, as to render its subperiosteal removal justifiable and necessary.⁵ Thus in the case recorded by

¹ Cornil et Ranvier, 'Manuel d'Histologie Pathologique,' Paris, 1881, p. 399.

² Chassaignac, 'Mém. de la Société de Chirurgie,' tome iv, p. 281.

³ See a case of acute periostitis of the tibia reported in Billroth's 'Chir. Klin.,' 1867, p. 487, Berlin, 1869.

⁴ For 1876, p. 73.

⁵ Cases where this has been done are referred to in the 'London Med. Record' for February 15th, 1876, p. 73, by Mr. Holmes. See also the 'Lancet,' vol. i, 1866, p. 340 (referred to by Verneuil in the 'Gaz. Hebdomadaire de Médecine et Chirurgie,' 1866, No. 21, p. 321); 'Med.-Chir. Proceedings,' vol. viii, p. 434 (Mr. W. Pye's case); Weinlechner, 'Allg. Wien. Med. Zeit.,' No. 24, 1879 (referred to in the 'London Med. Record,' August 15th, 1879); Holmes, 'St. George's Hosp. Reports,' vol. x, p. 500, and 'Surgical Treatment of the Diseases of Childhood,' 2nd ed., p. 385, and elsewhere.

Mr. Holmes, in which he performed this operation, there was no trace of osteomyelitis in the part of the bone (the tibia) examined. I shall presently have to advert to similar cases.

Considerable difference of opinion prevails with regard to the exact tissue where the changes commence in acute periostitis. The majority of English writers¹ seem to hold that the affection either commences in the deeper osteogenetic layer of the periosteum, or as a superficial ostitis, and that effusion or pus raises up the tendinous periosteum, thus leading to the denudation and necrosis of the bone. I do not think that the view is with us generally entertained, that acute spontaneous osteomyelitis is an affection of common occurrence, but it would seem to be otherwise on the Continent, judging by the writings of continental surgeons, particularly of German surgeons, and from the descriptions of their cases. The term *abcès sous-périostique aigu* employed by Chassaignac, signifies partial agreement with the opinion that acute periostitis begins beneath the periosteum. Cornil and Ranvier state that the disease consists principally in diffuse suppuration, which may have its seat in any part of the bone—beneath the periosteum, in the superficial layers, in the osseous tissue, or in the central medulla, which is tantamount to saying that there are no means of distinguishing clinically between acute periostitis and osteomyelitis. Billroth and other German surgeons go further, and express very decided views on this affection. Billroth, for instance, in an exceedingly able and suggestive essay on these diseases,² states that acute periostitis commences as inflammation of the outer cellular layer of the periosteum; according to his views the fibrous periosteum is destroyed subsequently by suppuration, and thus the bone becomes exposed. Extension inwards of the inflam-

¹ Cf. for instance, Holmes's 'Principles and Pract. of Surg.,' 1st ed., p. 392, and 'A Syst. of Surg.,' 2nd ed., vol. iii, p. 741; also Crampton "On Periostitis," 'Dublin Hosp. Reports,' vol. i, p. 331.

² 'Surg. Path.,' Hackley's Transl., pp. 289—293.

mation may lead to osteomyelitis. Further, according to Billroth, if it be denied that the outer layer of the periosteum constitutes part of that membrane, but is to be looked upon as the intermuscular cellular tissue, then there is no such thing as acute periostitis, for the tendinous part of the periosteum, he remarks, is as little liable to primary inflammation as the fasciæ or tendons. I hope to be able to show directly that this does not hold good universally, but that the views of Cornil and Ranvier, which have just been adverted to, are nearer the mark.

With regard, first, to the curious affection which attacks the terminal phalanges of the fingers (panaritium periostale of the German writers) and to which the term "whitlow" is somewhat loosely applied by writers in this country, it may be remarked that this is considered by some to be due to acute spontaneous osteomyelitis.¹ Now I do not think that this opinion is generally shared in this country; and it appears rather to me that the older view is more correct.² From some slight injury—often from no apparent cause—inflammation of the cellular tissue, tending to become diffuse, commences in the pulp of the finger. Owing to the close connection which exists between all the tissues of the part, from the skin down to the periosteum and bone,³ and the unyielding nature of these tissues, necrosis of the whole phalanx readily results. It would seem *prima facie* most improbable that the affection should begin as osteomyelitis, but it is not unnatural that this complication should follow, although Chassaignac partly denies that subperiosteal abscess will give rise to osteomyelitis;⁴ but he is here speaking of the disease as

¹ Billroth writes, "Akute spontane Osteomyelitis ist mit Ausnahme der Endphalangen (panaritium periostale) an den oberen Extremitäten selten," 'Chirurg. Klin.,' 1871-76, Berlin, 1879, p. 426.

² See Crampton, *op. cit.*, p. 333.

³ Humphry 'On the Skeleton,' pp. 18 and 394.

⁴ Chassaignac, who at one time appeared to doubt whether osteomyelitis was ever produced by subperiosteal abscess, seems to allow that this may take place by a case which he records in the 'Mém. de la Société de Chirurgie,' Tome iv, 1857, p. 320. An essay on the subject by the same author, entitled

it attacks the long bones, where the conditions are widely dissimilar. In panaritium periostale, then, if we suppose what I think undoubtedly to be the case, that the outer layers of the periosteum, *i.e.* the pulp of the finger, are primarily affected, we may seem to support the view that the same may obtain elsewhere; but the anatomy of the periosteum in other parts, say over the inner surface of the tibia or the lower third of the femur, is totally different.

Microscopical observation shows, that in some cases at least the inflammation distinctly commences beneath the periosteum, and again, that the characteristic effects of this disease may result without the occurrence of any osteomyelitis. This was the case in the following instance:

A boy, *æ*t. 13, was admitted into St. George's Hospital under the care of Mr. Holmes,¹ with acute periostitis of the tibia. Free incisions were made over the inner surface of the tibia and the bone was found exposed. Ultimately, on the 19th day of the disease, it was found necessary to amputate the thigh in order to save the patient's life. The knee-joint in this case was affected. A small piece of the thickened, softened layer next to the tibia was removed for microscopical examination. Especial care was taken to select the periosteum from a part where there were no overlying muscles. Sections were cut vertically through the specimen, which showed the appearances illustrated in Plate VII. On the side next to the bone was seen a uniformly thick layer of small cells. Above this the distinct layer of the fibrous periosteum, in one part of which commencing formation of new bone could be plainly made out. Still more external, in the parts overlying the fibrous layer, were evidences of extensive inflammation. The whole specimen was enormously vascular. It may be noted here, then, that the inflammation was

"Mémoire sur l'Osteomyélite," will be found in the 'Gazette Médicale' for August and September, 1854.

¹ To whom I am indebted for kindly permitting me to refer to this and other cases.

further advanced beneath the periosteum, where completely formed pus was found, than in the outer cellular layer, where the parts were merely infiltrated. Careful examination failed to show any inflammation of the medulla or of the substance of the bone itself. In the compact tissue there was not the slightest trace of what Mr. Erichsen has well termed "laminated expansion" of the bone, nor was there even the faintest discolouration. Examination of the medulla, indeed, showed changes more characteristic of breaking down, disintegration, and necrosis of that tissue than of actual acute inflammation or the sequelæ thereof. As seen under the microscope the fat cells were indistinct, granular, and breaking down. The whole specimen, in short, resembled medulla taken from a bone after decomposition had set in; but in this case, as the preparation was made from an amputated limb, properly preserved from the first, the disintegration must have taken place before the operation. I do not think, either from the appearance of the bone or the medulla, that the changes were the result of previous osteomyelitis, for the appearance was very different to that in a case to be presently described. I rather hold that the changes seen may be explained by the cutting off of the vascular supply owing to the periosteal affection, for this was a case which distinctly showed what Chassaignac has termed "*décollement périostique*."

I am aware that this view as to the breaking down and disintegration of the medulla is altogether at variance with that held by the French surgeons. Gerdy¹ goes so far as to doubt whether destruction of the medulla, other than that due to suppuration, ulceration, or new growths, has ever been seen. He states also, that he does not know of any examples of gangrene of that tissue. Nevertheless, if the medulla be examined it will be found to present very diverse appearances in different cases. No doubt periostitis, osteitis, and inflammation of the medulla are usually so mixed up together by the time the cases

¹ "*Périostite et Médullite*," 'Arch. Gén. de Médecine,' 1853, p. 141.

come under observation in hospitals, that the changes brought about in the medulla, either from the pre-existing inflammation or from necrosis of the tissue, might be indistinguishable one from the other; but when the bones are examined at an early period, as for instance, in the case above mentioned, the changes in the medulla due to the condition of acute periostitis alone present a marked contrast to those resulting from active inflammation of the tissue.

The other specimen (Plate VIII) was taken from a man *æt.* 21, in whom the disease first showed itself as an acute inflammation of the knee-joint. Subsequently acute periostitis of the lower third of the femur became evident, and eventually the thigh was amputated. I had not an opportunity of examining the medulla in this case, but there was no appearance of inflammation at the level where the femur was sawn through. In this case also, as shown in the drawing (Plate VIII), the inflammatory changes were chiefly seen in the deeper layers. The periosteum was lifted away from the bone, thickened, infiltrated and vascular. Inflammation was more advanced in the inner than in the outer layers; the infiltration extending outwards was chiefly noticed around and about the smaller blood-vessels.

The form of acute periostitis which Billroth describes as attacking the outer layers of the periosteum is, I think, more frequently seen over the lower third of the femur than anywhere else. The following case is an example of this variety:

A boy, *æt.* 9, was admitted into the Belgrave Hospital for Children under the care of my colleague, Mr. W. H. Bennett, with slight swelling over the hip-joint and pain in moving the limb. I am unable to refer to the notes of the case, but can recollect that the affection was of very recent date, not more than a few days. Under the influence of rest in bed, the symptoms about the hip-joint subsided, but acute synovitis of the knee then set in. Very shortly afterwards a fluctuating swelling was observed in

the popliteal space. An incision was made and the finger passed in went down at once to the popliteal space, but the bone everywhere was covered. Gradually the knee-joint recovered, and the abscess healed up. Such cases are not uncommon, although it too often happens that timely treatment is not adopted, and the periosteum thus becomes destroyed and the bone exposed. I merely cite the case here as furnishing a good and uncomplicated example of the variety of the disease described by Billroth. With regard to the articular affection, it may be observed, as has been pointed out, that, genetically speaking, the capsule of the joint is a continuation of the periosteum.¹

The following case is a typical one of intensely acute periostitis :

A girl, æt. 15, was admitted into St. George's Hospital with acute periostitis of the right tibia. Four days before her admission she felt ill, and had a rigor. Two days before her admission a slight blow over the tibia from a fall called her attention to the condition of the bone. The morning after admission the temperature was 104°, the pulse 145. The whole of the right leg below the knee was swollen and painful. At the upper part was an elastic, tense collection of fluid over the tibia. Incisions let out some blood-stained fluid, but no pus. The bone was extensively denuded. She died of pyæmia nine days after her admission—probably on the thirteenth day of the disease.

Post mortem.—The entire shaft of the tibia was evidently necrosed. The periostitis, as stated, began over the upper third of the tibia. There was no trace of osteomyelitis at this point, but lower down, at the centre of the shaft, the compact tissue over a very small extent of the bone was slightly laminated and discoloured, and the medulla beneath disintegrated and breaking down at one point into pus. Still, in this case I can hardly believe that the osteomyelitis was other than secondary to the

¹ Billroth's 'Surg. Path. and Therapeutics,' p. 290. See also Du Hamel "On the Relation of the Epiphyses and the Articular Cartilage to the Periosteum," 'Mém. de l'Académie Royale,' 1743, p. 143.

periostitis,¹ nor could this little localised patch of osteomyelitis have determined the total necrosis of the shaft. As a rule, however, where the symptoms take such an acute course and lead, as in this case, to metastatic abscesses in the lungs and disorganization of the joints, it is usual to find more extensive osteomyelitis.

To sum up. Acute inflammation may occur in the tissues immediately external to the periosteum, as in the case of the boy recorded above (p. 313). Such inflammation may lead to destruction of the periosteum and exposure of the bone, unless relieved by timely treatment. The affection is often exceedingly difficult to distinguish from acute subperiosteal abscess,² *i.e.* true acute periostitis. It is frequently associated with inflammation of the neighbouring joints, and attention is often drawn to the condition by the affection of these parts.

Acute periostitis, as we know the disease, usually begins in the deeper osteogenetic layer of the periosteum³ by inflammation and exudation around the vessels; the periosteum will become more or less detached and lifted up from the bone by the exudation,⁴ and in parts it will be actually destroyed. This condition may lead simply to exfoliation, or may be the precursor of more extensive necrosis, but it is not necessarily associated with osteomyelitis, even although all the train of symptoms, as we so frequently see them, be present. Acute periostitis, according to Chassaignac, comparatively seldom affects the joints,⁵ and this is true, but the exceptions to this rule are numerous enough. Acute periostitis may lead to destruction

¹ See note 4, p. 310, *supra*.

² On the Diagnosis of the several conditions from each other, see Chassaignac, *op. cit.*, p. 326.

³ I do not see how we can distinguish clinically from this—the commoner form—the disease which commences primarily as an acute ostitis.

⁴ True pus does not commonly form so very rapidly.

⁵ *i.e.*, the joints at either extremity of a long bone attacked by the disease. Pyæmic affections of the joints are only too common, especially when osteomyelitis exists, and must be carefully distinguished from the varieties here referred to in the text.

of the medulla, without the occurrence of any acute inflammation of that tissue. Acute primary osteomyelitis is a rare disease in this country.

With regard to the age of the patients, it is well known that those about the age of puberty are chiefly liable to the affection, but many cases are recorded at the extremes of life. An instance has been published of diffuse periostitis at the age of eleven days.¹ As an illustration that the old are not exempt, I may mention the case recorded by Billroth,² of a man, æt. 70, who was attacked with acute periostitis and osteomyelitis of the humerus from no known cause, and died of pyæmia.

I need hardly point out that although it is perfectly true that these various forms of the affection become confounded together very early, yet it is not on that account any the less important to study their separate courses, in so far as this is possible. The various forms will lead severally to widely different results, especially when—as unfortunately happens far too rarely—they are modified by early treatment. With this latter branch of the subject I am not now concerned.

¹ 'Path. Soc. Trans.,' vol. vi, p. 284.

² Billroth, 'Chir. Klin.,' 1867, p. 417, Berlin, 1869.

DESCRIPTION OF PLATES VII AND VIII.

Pathology of Acute Periostitis (Clinton T. Dent, F.R.C.S.).

PLATE VII.

Vertical section of periosteum from a case of acute periostitis.
× 100.

- a.* Layer, next to the bone, of small granulation cells and pus.
- b.* Fibrous periosteum.
- c.* New bone developing in the periosteum.

PLATE VIII.

Vertical section of periosteum from a case of acute periostitis.
× 45.

- a.* Layer, next to the bone, of small granulation cells and pus.
- b.* Fibrous periosteum.
- c.* Blood-vessels, around and about which the infiltrating cells chiefly extend from the deeper to the outer parts.