A CASE
OF
INNOMINATE ANEURISM
TREATED BY
SIMULTANEOUS DISTAL LIGATURE OF THE CAROTID AND
SUBCLAVIAN ARTERIES;
WITH REMARKS
ON THE BEHAVIOUR OF A TENDON LIGATURE.

BY
C. T. DENT, F.R.C.S.,
ASSISTANT SURGEON TO ST. GEORGE'S HOSPITAL.

(Received February 8th.—Read March 22nd, 1881.)

THE patient, whose case is given below, was admitted into St. George's Hospital on April 26th, 1880, under the care of Mr. Pollock. I have to express my warm thanks to Mr. Pollock for kindly permitting me to publish the case.

The history, symptoms, and progress of this case of aneurism, and the operation undertaken for its cure will, I hope, be thought worthy the attention of the Society. Double distal ligature of the carotid and subclavian arteries, for the cure of thoracic aneurism, is an operation of sufficient rarity and importance to justify a detailed

account of any case where this somewhat formidable measure is adopted.¹

On the history of the case however, and on the general results of the operative treatment, I shall touch but lightly, preferring to invite the attention of the Society to certain pathological points in connection with the ligature of the arteries with a peculiar form of ligature.

The history of the case is as follows :

Samuel L—, æt. 37, an engineer, had always been healthy up to the commencement of the disease for which he was admitted. His habits of life were active, and for many years he had been accustomed, as “first tenor” in a choir, to much singing. On occasions he had rather strained his voice, but never felt any ill effects for long. He had never suffered from syphilis. About twelve months previously, in April, 1879, he noticed some difficulty in producing his “chest notes,” as if there existed some obstruction, which was referred to the root of the neck. In October, 1879, he first felt pain about the neck and the sterno-clavicular joint. The pain though not severe was tolerably constant; three months later, some swelling was observed in the right sterno-clavicular region, which increased rather rapidly and occasioned much pain down the arms and in the back. From this time he began to lose strength and during the next three months, lost about eighteen pounds in weight. The breathing became affected, and the voice a little husky and stridulous; he was troubled also with a constant slight clearing cough, and with what the German physicians well term “Hustenreiz.”

When admitted, he was seen to be a tall, well-formed, rather spare man. His expression was rather anxious; the breathing was a little harsh, and the voice weak and slightly husky, but he had no severe dyspnoea.

The radial pulse was the same on both sides, regular and quiet.

¹ Mr. P. Barwell, in a paper read before the Royal Med. and Chir. Soc. in December, 1878, mentions eleven cases. (*Med.-Chir. Trans.*, vol. lxii, 1879.)

Over an area extending on the right side, from a little above the clavicle to the third or fourth rib below, and behind the sternum, was situated a moderately prominent, ill-defined, pulsating swelling, which gave the sensation of being deeply situated. The pulsation, however, had no expanding character, and the tumour was at first thought to be possibly a malignant mediastinal growth; there was no thrill to be felt, nor bruit to be heard; on auscultation the heart sounds were communicated with great distinctness to any part of the swelling. The carotids on the two sides pulsated equally. Some difficulty was experienced in swallowing dry food.

Complete rest in bed and large doses of iodide of potassium brought about a marked improvement. It appeared that the symptoms had previously been relieved by this drug. As, however, the pain and the difficulty of deglutition and breathing diminished, the swelling assumed more obviously the appearance of aneurism. From May 31st to June 10th his diet was limited to half a pint of milk and half a pint of beef tea daily, but without yielding any marked benefit. The improvement resulting from rest and the iodide did not last long, and by June 10th the swelling had become much more prominent and the pulsation more marked. The aneurism had increased upwards and displaced the sternal half of the clavicle forwards. A faint bruit was now audible posteriorly, on a level with the sixth dorsal vertebra.

On June 10th, Mr. Pollock tied the right common carotid above the omohyoid, and the third portion of the right subclavian artery. For the carotid a tendon ligature was employed; in tying the second knot, the ligature broke. The first knot sufficed to keep the loop firmly applied to the artery, but it was thought insecure to leave it so. Accordingly, a thick catgut ligature was passed round the artery and tied close to the tendon. The subclavian artery was secured with a flat tendon ligature only. Both arteries seemed perfectly healthy. The operation was not performed under the antiseptic system.

The effect on the aneurismal tumour was very marked immediately ; the clavicle sank back a little and its outline became more defined ; the pulsation distinctly decreased and the prominence below the clavicle diminished.

Two days later (June 12th) he had a severe attack of dyspnœa, lasting several minutes. The next day he again had a similar attack and became rather cyanosed. The attacks were thought to be due to nervous irritation ; opium was given, with the effect of markedly relieving the dyspnœa and distress. Four days after the operation the tumour had so much decreased, or the lung collapsed, that there was decided flattening on the right side over the first four ribs. Subsequently, similar flattening was observed on the left side, and later on a concavity in this situation. The respiratory movements of the thorax were very restricted and the abdomen in the epigastric region retracted ; the respirations were 42, and very laborious. During the next six days (up to June 18th) the symptoms rather increased, though at times he seemed to rally and was in comparative comfort. The attacks of dyspnœa then became more frequent, and the expression at times was one of terrible anxiety and distress. Finally, on June 20th, ten days after the operation, the breathing once more became very laboured ; the frequency of the respirations increased till they reached 52 ; the pulse grew weaker, and he died very suddenly.

The day after the operation faint pulsation could be felt in the radial pulse, but this disappeared again three days later. The temporal pulse on the right side was felt for the first time five days after the operation.

The temperatures of the two sides were carefully taken. The day after operation the temperature of the right ear was about one degree lower than the left ; by the fifth day they were nearly identical. In the axillæ the temperatures were almost exactly the same from the very first.

The post-mortem examination was performed by Dr. Isambard Owen, to whom I am indebted for the following notes and for the preparation of the dissection :—" Lungs,

emphysematous, more solid than natural; abundant, slightly purulent fluid exudes on pressure; some of the lobules are choked with yellowish material. The apex of the right lung is compressed by the aneurism.

"Heart uncontracted, filled with soft black clot; tissues healthy.

"The aorta is covered with large patches of atheroma. In the position of the orifice of the innominate artery is an opening, the size of a sixpence, with funnel-shaped edges. This opening leads into an aneurism of lobular form, about the size of an orange, situated behind the first piece of the sternum and rising above it. The aneurism is full of laminated clot. The sternum is eroded at one point, and the sterno-clavicular joint opened. At the level of the lower border of the first rib the anterior wall of the trachea is compressed by the tumour, and the tracheal wall is eroded completely through, the opening being the size of a threepenny-piece, and bounded by the cut ends of the cartilaginous rings; the trachea thus communicates with the connective-tissue plane of the mediastinum, but there is no corresponding opening from the aneurism. The right subclavian and carotid rise from about half an inch of common innominate artery, from the right side of the aneurism."

The opportunity was thus accorded to me of examining the artery and noting the changes that had taken place at the seat of ligature in the course of ten days. On dissecting down to the carotid artery, I found that the deeper part of the wound was united. A very moderate amount of lymph had been effused in and about the carotid sheath. The knot of the ligature was completely buried, but a depression indicated its position. About an inch and a half of the artery was removed, together with the ensheathing lymph, and hardened in a mixture of chromic acid and spirit. On making a longitudinal section through the specimen, the appearance was shown which is most accurately indicated in the accompanying drawing (Plate III, fig. 2). The knot of the tendon ligature was seen at

the upper part in close contact with the artery, encysted in a small cavity in the effused lymph. The knot was almost gelatinous in appearance, but small, glistening, tendinous bands could be seen crossing the dark space. The vessel was completely occluded for a quarter of an inch, being represented by a cord of fibro-cellular tissue. A firm clot in close connection with the wall of the artery filled up the vessel on the distal side of the ligature. On the proximal side the clot was less well formed, and showed a tendency to break down in the centre.¹

For purposes of microscopical examination sections were made in a longitudinal direction, including both ends of the obliterated artery and the knot of the ligature. Other sections were made in a plane at right angles to these through the knot, so as to show the ligature encircling the vessel. My best thanks are due to my colleague, Mr. W. H. Bennett, for making the sections, and to Mr. F. C. Compton for his care and skill in preparing and mounting the same.

In some of the sections the connection of the ligature with the artery was so slight that the two were separated either in cutting the section or in mounting the specimen. In many others, however, and even in very thin sections, the connection was much firmer, and a very slight, but still appreciable, amount of force was necessary in order to detach the knot of the ligature, or the tendon itself, from the vessel. The adhesion was about as firm as that of a skin wound uniting by first intention at the end of a few hours. To this point I shall return later on.

A considerable part of the tendon ligature, where not in immediate contact with the artery, was altogether unaltered. To the naked eye it seemed unchanged, and even under the microscope it was quite normal in appearance.

¹ Senftleben has asserted that an internal clot does not usually form when catgut ligatures are used (Virchow's 'Arch.,' lxxvi; 'Gaz. Hebdomadaire de Med. et de Chirurg.,' 1880, No. 20, p. 325). But this assertion has been altogether disproved by Dr. J. F. Arnaud, in an essay entitled 'Contribution à l'étude de la ligature dans le traitement des anévrismes,' Paris, 1880.

No trace of the catgut ligature could be found. It is quite possible, however, when we look at the extent of artery obliterated in the case now under consideration, that the catgut ligature had contributed an important share towards the occlusion of the vessel. This quite accords with the results of Dr. Arnaud's experiments.¹ This observer ligatured the carotid and femoral arteries of dogs fourteen times with carbolised catgut. At the end of four days in one experiment, at the end of seven days in two, and at the end of nine in another, the parts were examined and no trace of the catgut could be discovered.²

On examining the sections under the microscope (see Plate II and Plate III, fig. 1), the following points are observable :

1. The external coat of the artery is not ulcerated. In this particular case it might be asserted that the breaking of the knot when the ligature was tied must have led to some relaxation of the ligature, and that, consequently, the non-ulceration proves nothing ; but examination of the subclavian artery, which was also tied with a tendon ligature and on which the knot was still found firm, showed no ulceration of the external coat. I conclude, therefore, that the tendon behaves like other animal ligatures, notably catgut, and that slight softening and swelling of the ligature obviates any ulcerative action. A further and highly important explanation—to be further adverted to presently—would appear to consist in the fact that new blood-vessels are early developed in those parts of the tendon which lie close to the artery.

The action of the tendon ligature, therefore, resembles that of the catgut, in so far as it does not lead to ulceration of the external coat ; but it may possibly differ from the catgut in that this effect is not wholly due to the softening

¹ Op. sup. cit.

² In other cases, however, 4, 9, and 11 days respectively, after the application of the ligature, the catgut was still distinctly recognisable, and in one instance, 16 days after the application of the ligature the catgut was found only softened and infiltrated. Op. cit., p. 25.

of the ligature. It has been asserted that blood-vessels develop in the catgut. That this at any rate is not always the case is proved by the occasionally early disappearance of the ligature.

2. The action of the tendon on the middle and inner coats of the artery is the same as that of any other ligature, that is to say, these coats are ruptured in places, and their inner walls lie in contact.

3. In some of the sections the tendon ligature is seen to be most closely connected with the artery, and but for the differentiation of the tissues by the staining fluids, the artery could hardly be distinguished from the contiguous ligature. Still, there is no evidence of destruction of the arterial coats, though they are compressed by the ligature.

4. In places where the tendon lies in close connection with the artery it was infiltrated with small, round, granulation cells or leucocytes. The cells, which are readily stained by logwood, have penetrated into the interfascicular spaces, and tend to split up the tendon into longitudinal bands. Close examination of these parts reveals the presence of what I cannot but conclude are newly developed blood-vessels. Round and about these vessels the cells are numerous, and show a tendency to elongation (see Plate III, fig. 1); the latter change has been described by numerous observers (O. Weber,¹ Bubnoff,¹ Szuman,¹ Senftleben,² Billroth,³ and others) as preliminary to the formation of blood-vessels in the clot itself.

Several objections may be raised against the supposition that these vessels seen in the specimens are newly formed. For instance, they present a somewhat different appearance to newly developed blood-vessels elsewhere, such as can be seen in the tadpole's tail, or such as have been recently described by Tillmanns⁴ and beautifully figured in his work. But here the conditions under which the blood-

¹ Quoted by Arnaud, *op. cit.*, p. 48.

² *Op. jam. cit.*

³ 'Surg. Path.,' Hackley's transl., 1877, p. 113.

⁴ Tillmanns, "Experimentelle und anatomische Untersuchungen über

vessels developed were totally different from those in the case under consideration; the process is more analogous to one form of the development of blood-vessels in granulations which Billroth¹ and others have described.

Secondly, it might be thought that these blood-vessels are merely those normally found in tendon. This view I think is altogether negatived by two facts; one is that in parts of the tendon remote from the artery absolutely no blood-vessels can be found. The second is that numerous cells are found in the neighbourhood of the new vessel besides those which directly contribute to its formation; an appearance quite different from that of a blood-vessel which has existed for any length of time. Finally, it may be said that the development of blood-vessels is astonishingly rapid, considering that the ligature had only been *in situ* for ten days. It may be so; if the answers to the other objections are deemed satisfactory, this last difficulty falls to the ground.

5. In one or two of the sections it appears possible to trace blood-vessels which pass across the line of connection and extend into the artery on the one hand, and into the tendon on the other.

6. The different behaviour of the infiltrating cells towards different tissues is well illustrated by some of the specimens in which some muscular tissue, which was probably adherent to the ligature, is seen surrounded by numerous granulation cells. The muscular bundles are being attacked and eaten away. No giant cells can be distinguished at this part, as might possibly have been expected.

Such are some of the points observed under the microscope in these specimens. It is far from my intention to draw any decided conclusion from a single case, or to assert that the tendon ligature is more reliable than even ox-aorta ligatures, of which Mr. Barwell speaks so favourably,

Wunden der Leber und Nieren," Virchow, 'Arch.,' 1879, Bd. lxxviii, Heft 3, p. 437.

¹ "Untersuchungen über die Entwicklung der Blutgefäße," Berlin, 1856. Quoted in Paget's 'Surg. Path.,' p. 160.

or chromic catgut, as recommended by Dr. MacEwen¹ and Mr. Lister.²

Acupressure has shown that it is quite sufficient to occlude an artery for forty-eight hours. For my own part I fail to see any great force in the objections urged in many quarters and by good authorities against catgut; even though this material may dissolve or soften at an early period. Still, those who mistrust the catgut ligature may be satisfied with an animal material which requires less preparation and is more constant in its behaviour when in contact with living tissues. I do not venture to assert at present that in tendon we have a ligature fulfilling these conditions; nevertheless, I trust that the subject may be thought worthy the attention of those who take an interest in the still unsettled questions connected with the ligature of arteries.

It will of course be understood that in this paper no priority at all is claimed in the use of a tendon ligature. In the 'Clinical Trans.' for 1878, vol. xi, p. 103, the late Mr. G. W. Callender suggested the possible value of the material in the course of some remarks on tendon ligatures; these tendons were, like those employed in this case, taken from the tail of a marsupial, and were supplied to Mr. Callender by Mr. J. M. Girdlestone, of Melbourne. Other references to the use of tendon ligatures will be found in the 'Australian Medical Journal' of December 15th, 1879, and in the 'London Medical Record' of April 15th, 1878. But the material has been, I believe, often used before by the older surgeons, though perhaps no such good opportunity has previously occurred as fell to my lot in the present case of ascertaining its exact mode of action on an artery.³

¹ 'British Med. Journal,' vol. i, 1879, p. 272, and Sept., 1878, p. 359.

² Lister, "On Chromic Catgut Ligature," 'The Lancet,' Feb. 5, 1881.

³ Copious references to the literature on the subject of this paper will be found in an essay by Dr. Fritz Raab, entitled "Ueber die Entwicklung der Narbe im Blutgefäss nach der Unterbindung," Langenbeck's 'Arch.,' vol. xxiii, 1879, p. 156.

DESCRIPTION OF PLATES II AND III,
Innominate Aneurism treated by Distal Ligature of Carotid and
Subclavian Arteries with a tendon ligature (Mr. C. T. Dent).

PLATE II.

Transverse section of artery showing tendon ligature *in situ*,
 $\times 45$. *a.* Carotid artery. *b.* Lumen of artery. *c.* Shows rupture
of inner coats. *d.* Tendon. *e.* Infiltration of small cells splitting
up tendon into longitudinal bands.

PLATE III.

FIG. 1.—Shows elongation of cells, and development of new blood-
vessel in the tendon. $\times 200$.

FIG. 2.—Twice natural size. Longitudinal section of artery show-
ing ensheathing lymph and tendon ligature *in situ*. The vessel
is completely occluded. The clot is less perfect on the proximal
than on the distal side of the occlusion.

Fig. 1.

Fig. 2.