For each subject two sets of determinations at 4, 8, 12, and 16 litres per minute were made. In the first set breathing was through the nose, and in the second through the mouth. Mouth-breathing appeared to make little difference, as will be seen from the table and graph (Fig. 6).

Many more analyses were made than are represented in the above table. Oxygen figures are given to the nearest whole number. It would be misleading to suggest that greater accuracy is possible, as there are important potential sources of error in both sampling and analysis. It seems probable that, taking everything into consideration, the margin of experimental error may well be up to 5 per cent.

The effect of different rates of respiration was not determined, since this has been done very thoroughly by Barker, Parker, and Wassel⁵ using the oropharyngeal technique on three normal subjects. They found that respiratory rates of 16, 30, and 60 per minute made little difference to the alveolar concentration of oxygen at the rates of flow under study. Our results accord closely with those of these workers. The graph from their paper is reproduced in Fig. 7.

FIG. 7.—Graph reproduced from paper by Barker *et al.*

Summary

1. Apparatus, new in several respects, for the nasal catheter administration of oxygen has been described.

2. Alveolar oxygen concentrations, at flows from 4 to 16 litres per minute, ranged from 29 to 58 per cent. (mean values). Thus the fact that optimal concentrations of oxygen for prolonged administration can be obtained by the apparatus has been demonstrated.

We wish to express our thanks to Professor Samson Wright for great help and encouragement. Mr. D. Russell Davis most kindly consented to act as one of the subjects in the investigation. Mr. S. A. Sewell has taken considerable trouble with the illustrations.

The catheter carrier may be obtained from Messrs. Bell and Croyden of Wigmore Street, London, W.1.

REFERENCES

- ¹ Campbell, A., and Poulton, E. P.: *Oxygen and Carbon Dioxide Therapy*, 1934, London.
- ² Potts, W. H.: *Amer. Journ. Med. Sci.*, 1932, clxxxiv, 616.
- ³ Boothby, W. M.: *Journ. Amer. Med. Assoc.*, 1932, xcix, 2026 and 2106.
- ⁴ Wineland, A. J., and Waters, R. M.: *Arch. Surg.*, 1931, xxii, 67.
- ⁵ Barker, M. H., Parker, D. M., and Wassel, G.: *Journ. Amer. Med. Assoc.*, 1934, ciii, 244.
- ⁶ Bourne, G.: *Lancet*, 1922, ii, 23.
- ⁷ *Journ. Amer. Med. Assoc.*, 1934, cii, 1870.
- ⁸ Beaumont, G. E., and Dodds, E. C.: *Recent Advances in Medicine*, seventh edition, 1934, London, p. 123.

The Anti-noise League, 66, Victoria Street, S.W.1, announces that in response to many inquiries it has published further leaflets, which form part of a series now in preparation. They include a note on that branch of hospital planning which is directed to the prevention of noise, and similar pamphlets refer to noise in schools and in domestic buildings. There are notes also on the protection of night workers from noise during the day. Suggestions are made as to the best methods of dealing with the more common noise complaints.

SPLENECTOMY FOR THROMBOCYTOPENIC PURPURA

BY

GWENDOLINE SMITH, M.D., F.R.C.S.

ASSISTANT SURGEON, SOUTH LONDON HOSPITAL FOR WOMEN

The disease known as purpura haemorrhagica embraces a number of different conditions the aetiology of which is not clearly understood, but essential thrombocytopenic purpura haemorrhagica has now been separated from the general group of purpuric diseases, and cases of it can be diagnosed with some certainty. The characteristic features are purpura, haemorrhages from various organs, a diminished platelet count, and prolonged bleeding time. The condition may be acute or chronic, and occurs in children and adults. There is much interest in the blood changes and in the differential diagnosis of the disease, but the surgical importance lies in the treatment of the established condition. Surgical treatment was only considered just less than twenty years ago, when Kaznelson first performed splenectomy for it, and a review of the literature and medical textbooks reveals considerable divergence of opinions on the indications for the operation, and also on the late results of medical and surgical treatment.

In view of this it seems important that yet more isolated cases should be published, as the condition is uncommon, and a large series must eventually be collected before the best method of treatment can be established. A brief record is therefore given of a case treated by splenectomy over a year ago, whose progress has so far been satisfactory, and can be reported on again at a later date.

Case Record

A married woman, aged 41, was admitted to the South London Hospital under the care of Dr. Pickard on February 17th, 1934.

History.—For six weeks the patient had noticed a rash on the chest and back, bruising of the limbs, and occasional bleeding from the gums. Epistaxis had occurred once. Menstrual periods had previously been normal, lasting seven days, but the one just prior to admission had commenced ten days early, lasted ten days, and had been unduly profuse. She had no children, and had had no miscarriages. She had been in good health up till this time.

On admission the general condition was quite good; colour fair. The gums were unhealthy, and were bleeding at several points. The abdomen was covered with a purpuric rash, which was also visible on the back. The spleen could not be felt. Heart and lungs, pelvic organs, and urine were all normal. Blood count: red blood cells, 4,190,000; haemoglobin, 70 per cent.; colour index, 0.83; white blood cells, 6,000; some anisocytosis.

Progress.—On February 22nd uterine haemorrhage recommenced, and was very profuse. More subcutaneous haemorrhages appeared. Blood count: red blood cells, 3,100,000; platelet count, 21,700. March 3rd: retinal haemorrhages occurred in both eyes. March 5th: blood count—red blood cells, 3,333,000. March 9th: bleeding from gums more severe; blood transfusion given. March 15th: blood transfusion repeated. March 19th: uterine haemorrhage commenced again, with profuse loss. March 23rd: blood count—red blood cells, 2,470,000; platelets, 17,500. General condition much worse, patient very pale, pulse rapid and of poor volume. As all types of haemostatic drugs had been given without effect, it had already been decided that splenectomy was necessary, but the patient had become so much worse during the last twenty-four hours that it was obvious that she would die very soon, and that operation must be undertaken at once, if at all.

Operation.—Emergency splenectomy was performed. The spleen was about one and a half times the normal size, and there were no adhesions. The operation was rapidly com-

pleted as the patient's condition was extremely poor. Bleeding from the abdominal wall was easily controlled as soon as the splenic pedicle had been ligated.

Post-operative Progress.—Saline infusions were given for twelve hours, during which time the condition was critical, but there was no more uterine bleeding. After this there was no further anxiety, and progress was steadily maintained. The gums healed rapidly, and necessary dental extractions were performed before discharge from hospital without any undue bleeding. The patient was sent to a convalescent home on May 22nd.

The following summary of some of her post-operative blood counts shows the rate of recovery of red cells and platelets.

		Red blood cells	Platelets per c.mm.
March 26th	...	2,100,000	26,700
March 28th	...	2,360,000	50,000
April 7th	...	2,800,000	67,000
April 18th	...	3,870,000	121,000
May 18th	...	4,320,000	121,000
November 15th	...	4,200,000	140,000
December 5th	...	4,300,000	200,000
May 7th, 1935	...	4,250,000	250,000

Haemoglobin, 90 per cent.

Menstruation did not return until three months after operation, and the periods have since been regular and not excessive. In May, 1935, the general condition was very good. The colour was excellent, and there had been no return of haemorrhages or purpura.

Pathological report of the spleen is as follows: Spleen enlarged and of normal shape. Microscopical examination shows dilated venous sinuses with thickened walls. The Malpighian bodies are of normal size; large mononuclear cells, some being degenerate, are seen in the centre of the bodies. Giant cells are not seen.

Discussion

This case would be properly classed as one of acute thrombocytopenic purpura haemorrhagica. As far as can be ascertained the acute cases form about 20 to 25 per cent. of the total number, and the immediate mortality after splenectomy is given by Price¹ as 80 per cent. Pemberton² reports eleven cases treated by splenectomy, with two deaths, while Marzullo³ mentions three deaths in four cases, but in the same series there were four patients treated medically, and these all died. It is probable that, with an earlier diagnosis and decision as to the necessity for surgical treatment, the immediate mortality can be considerably reduced, and should not be higher than 15 to 20 per cent. in the acute cases. Undoubtedly in the case here reported surgical treatment saved the patient's life, and probably should have been undertaken some days sooner.

It is more difficult to trace the subsequent history of those patients who recover from operation and to say what proportion of them are liable to suffer from a recurrence of the symptoms. In Pemberton's² series, of the nine patients who recovered only one is stated to have suffered from later haemorrhages, the others remaining well for periods up to eight years after operation at the time of reporting.

The value of splenectomy for chronic cases is more controversial. The immediate operation mortality rate is much lower than in the acute cases. Price¹ gives 8 per cent., and Pemberton's² series from the Mayo Clinic shows thirty-four patients with no deaths directly attributable to the operation. There was one death five years later, from severe uterine bleeding at the onset of puberty. Recurrent attacks of bleeding occurred in ten cases, but showed a tendency to get less and less severe, only persisting eventually in four cases. Spence⁴ reviewed 101 cases in the literature in 1928, and found that splenectomy was beneficial in 16.6 per cent. of acute cases and in 80.9 per cent. of chronic cases. As some of these reports were only made a year after operation the figures are not

so valuable, as recurrences may take place later. Myers⁵ reports three cases well after periods of eight years (two patients) and three and a half years (one patient).^{*} Hartfall and Oldfield⁶ published an interesting account with full blood counts, in a woman of 25, whose only symptom was menorrhagia. Splenectomy was successfully performed, and the patient had remained well for a year at the time of reporting.

Until more knowledge of the pathology of the disease is gained the value of any one method of treatment must be based on records such as these. Evidence is in favour of the spleen being only one part of a diseased reticulo-endothelial system, and some observers do not recommend splenectomy on the grounds that only partial removal of the diseased tissues is thereby effected. It seems that the spleen is certainly one of the largest single structures affected, and if clinical results show that patients are improved and remain well, it would appear wiser to base treatment on these results.

It is clear that the actual platelet count is not a certain index of the severity of the disease, nor of the degree of recovery after operation. The platelet count may rise immediately the splenic vessels are secured, but may not show any increase for some days later. The criterion of permanent recovery is freedom from haemorrhages and purpura over an indefinite number of years. The presence of accessory spleens is occasionally a cause of return of symptoms, and they should be looked for at operation and removed if found.

The figures quoted above show that in a certain number of cases a long period of good health has been assured by operation in the acute and chronic states of the disease. In acute cases which fail to respond to medical measures the mortality without surgical intervention is in the region of 100 per cent. In chronic cases the indications are not so clear, but splenectomy would appear advisable in those of moderate severity which fail to improve with drugs and blood transfusions, or suffer from acute exacerbations of purpura, with a persistent low platelet count. Cases of mild severity will respond to treatment of anaemia and removal of septic foci. A certain percentage of recurrences must be expected, but one may anticipate that even then the symptoms will gradually lessen and eventually disappear in the majority.

Conclusions

1. A case of acute purpura haemorrhagica in a woman of 41 years is recorded. Splenectomy was successfully performed, and the patient remains in very good health fifteen months after operation.

2. The immediate and remote results of splenectomy for this condition are discussed.

3. Conclusions based on published cases show that splenectomy is certainly indicated in acute cases with severe bleeding, and delay in operating adds to the operative risks. In chronic cases splenectomy should be considered in all except the mild ones, but though the immediate mortality rate is much lower, permanent cure cannot be expected in every case.

My thanks are due to Dr. Ellen Pickard for asking me to treat this case, and to Dr. Whittingham, pathologist to the South London Hospital, for numerous blood examinations and reports.

REFERENCES

- 1 Wits, L. J.: *Price's Textbook of Medicine*, fourth edition, 1933, p. 806.
- 2 Pemberton, J. de J.: *Amer. Journ. Surg.*, 1934, xxiv, 793.
- 3 Marzullo, E. R.: *Ibid.*, 1933, xxii, 92.
- 4 Spence, A. W.: *Brit. Journ. Surg.*, 1928, xv, 466.
- 5 Myers, B.: *St. Bart's Hosp. Journ.*, 1933, xli, 44.
- 6 Hartfall, S. J., and Oldfield, C.: *British Medical Journal*, 1934, ii, 8.

* See also *British Medical Journal*, September 7th, 1935, p. 445, published after this paper was submitted.