for valuable advice in the preparation of the paper; and Professor T. F. Hewer for assistance in the interpretation of histological material. We gratefully acknowledge the generous help received from the Medical Research Fund of the United Bristol Hospitals towards the expenses of this study.

REFERENCES

- Bennett, I. L., and Cluff, L. E. (1957). *Pharmacol. Rev.*, **9**, 427.
 Chasis, H., Ranges, H. A., Goldring, W., and Smith, H. W. (1938). *J. clin. Invest.*, **17**, 683.
 Gillespie, W. A., Linton, K. B., Miller, A., and Slade, N. (1960). *J. clin. Path.*, **13**, 187.
 Hare, R. S. (1950). *Proc. Soc. exp. Biol. (N.Y.)*, **74**, 148.
 Houghton, B. J., and Pears, M. A. (1957). *Brit. med. J.*, **1**, 622.
 Hutt, M. S. R., Chalmers, J. A., MacDonald, J. S., and de Wardener, H. E. (1961). *Lancet*, **1**, 351.
 Katz, Y. J., and Moore, R. S. (1960). *Excerpta Med.*, International Congress Series, **29**, 48.
 Leather, H. M., and Gault, H. M. (1962). *Lancet*, **1**, 746.
 — and Hutchings, M. T. (1960). *Ibid.*, **2**, 654.
 Montgomerie, J. Z., and North, J. D. K. (1961). *Ibid.*, **2**, 824.
 Pears, M. A., and Houghton, B. J. (1958). *Ibid.*, **2**, 128.
 — (1959). *Ibid.*, **2**, 1167.
 Taylor, R. D., and Page, I. H. (1943). *J. Amer. med. Ass.*, **121**, 754.

BACTERIOLOGICAL EXAMINATION OF URINE

BY

W. R. CATTELL, M.B., M.R.C.P., M.R.C.P.Ed.

Lecturer in Medicine

M. J. LEFFORD, M.B., B.S.

Registrar

From the Professorial Medical Unit and the Department of Pathology, St. Bartholomew's Hospital, London

In recent years there has been a reappraisal of the bacteriological methods in use for the diagnosis of infection of the urinary tract. Appreciation of the risk of infection following the catheterization of female patients has led to several techniques being developed to allow collection of mid-stream samples of urine from women (Hart and Magee, 1957; Boshell and Sanford, 1958; Wood *et al.*, 1960; Leather and Hutchings, 1960). In conjunction with these, quantitative urine culture has been advocated as a means of distinguishing between contaminated urine and genuine infection (Kass, 1955, 1956; Sanford *et al.*, 1956; MacDonald *et al.*, 1957; Jackson *et al.*, 1958; Monzon *et al.*, 1958).

Unfortunately most of the methods suggested for the collection of mid-stream specimens are elaborate, time-consuming, and often impracticable because of age or infirmity on the part of the patient. The standard method for performing bacterial counts on urine is expensive in time and material, and demands either immediate culture of the specimen or refrigeration at 4° C. There is also some doubt about the need for elaborate bacterial counts for the detection of significant bacteriuria (100,000 organisms/ml.).

In this paper the different methods recommended for the collection of "clean" specimens of urine from women have been re-examined, and a simplified procedure is suggested. In addition a rapid method for performing quantitative urine culture is described and compared with the standard technique. Finally, the results obtained by quantitative culture and from routine laboratory reports are compared.

Material and Methods

Urine specimens were obtained by the nursing staff of busy general medical wards. So far as possible these were obtained from consecutive admissions, irrespective of the reason for admission or the presence or absence of suspected renal disease.

Collection of Specimens

Several methods were attempted, but eventually two were adopted for collecting urine specimens. Clean specimens were obtained by preparing the patient as for catheterization. The perineum was first washed with soap and water. This was followed by cleansing of the vulva with a sterile swab soaked in 1% cetrimide solution, the labia being separated and cleansed from before backwards. The patient was then sat on a sterile bed-pan and encouraged to pass urine. The sample

collected in the bed-pan was used for bacteriological examination. No attempt was made to obtain a mid-stream specimen. The second method employed a sterile glass tube as described by Leather and Hutchings (1960). The same preparation was carried out, but this was followed by placing the bell-shaped end of the tube over the urethral orifice. The patient was encouraged to void urine, and a mid-stream specimen was collected in a sterile container. All samples collected were divided between two sterile specimen jars and immediately taken to the laboratory or refrigerated at 4° C. until this was possible.

Where the results of urine culture indicated infection, catheter specimens were obtained within 24 hours, and before any treatment was begun.

Bacteriological Methods

One-half of each specimen was examined in the routine manner. This involved seeding a blood-agar plate with a urine inoculum of variable size, depending on the number of bacteria seen on the Gram film of the centrifuged deposit. The inoculum was distributed with a glass rod spreader. The plate was incubated at 37° C. overnight and then reported in the conventional way. The other half of each specimen was taken to a separate laboratory and quantitative cultures were performed. Two methods were employed. In the first of these nutrient-agar pour plates were prepared, using undiluted urine and urine diluted 1 in 100 in normal saline. Colony counts were made after overnight incubation at 37° C. The second method involved the use of a fused platinum loop of 2-mm. internal diameter. This loop was immersed vertically in the undiluted urine and discharged fully on to the surface of a well-dried blood-agar plate. This inoculum was distributed with a glass spreader, was incubated at 37° C. overnight, and the surface colonies were counted.

Since bacterial counts were performed on specimens which were also being examined independently in the routine manner, a comparison of the results obtained by the two methods was undertaken. It has been the practice here to report the results of urine cultures on a roughly quantitative basis thus: sterile; scanty growth; mixed growth of doubtful significance; pure growth of "X." In this study such reports were analysed by a clinician familiar with this practice and allocated to one of the three groups representing sterile or contaminated urine, infected urine, or urine which yielded an

equivocal result requiring further study. This was done without prior knowledge of the bacterial count.

Results

In order to allow comparison between our results and those reported elsewhere (Kass, 1956, 1957; Jackson *et al.*, 1958) we have adopted the convention of regarding bacterial counts of less than 10,000/ml. of urine as negative, by which is meant sterile urine or simple contamination during voiding and collection. Counts of more than 100,000/ml. we call positive and consider indicative of infection of the urinary tract; while counts between 10,000 and 100,000 were regarded as equivocal and requiring repetition.

Quantitative Culture

Bacterial counts performed by the pour-plate technique were found to be satisfactory, although occasional high counts due to diphtheroid bacilli were obtained, so representing false-positive results on the basis of bacterial counts alone.

The second method used was designed to investigate the possible use of surface plates for quantitative culture using a standard inoculum of such size that accurate colony counts could be carried out over the critical range of 10,000–100,000 bacteria per ml. The 2-mm. loop was found to discharge an inoculum of average size 0.00176 ml. (S.D. 0.00024). Calculation shows that urine containing 100,000 bacteria per ml. should give a colony count of 176, and urine containing 10,000 bacteria per ml. should give a count of 18. It was found that, using this technique, colony crowding was no problem. Further, where culture of the urine produced a mixed growth, colony counts of each type of organism were possible and identification of the bacteria from their growth characteristics and Gram-stained film could be done from the same plate. Reproducibility of counts on the same specimen were found to be satisfactory at all levels of bacteriuria, but not as good as with the pour-plate technique. This is to be expected from sampling errors incurred by using a much smaller amount of the original urine. We have been unable to demonstrate any significant loss of organisms on the spreader used in this technique. As with the pour plates, occasional high counts due to contamination with diphtheroids were obtained.

When the counts are divided into the three critical groups—that is, negative, equivocal, and positive, on the basis of counts of $0-10^4$, 10^4-10^5 , and $>10^5$ bacteria per ml. respectively by the pour-plate technique, or 0–18, 19–176, and >176 colonies per plate by the surface-count method—a good correlation is obtained (Table I). There is complete agreement between the two methods in 89% of instances. There was one false positive by the surface plate count. The number of equivocal results increases from 13 by the pour-plate technique to 19 by the surface count. There were no false-negative results by the surface plates.

TABLE I.—Comparison of Results from Surface Culture with Those from Pour Plates

Bacterial Counts per ml. by Pour-Plate Method	Surface Colony Counts (per 0.00176 ml. of Urine)		
	0–18	19–176	>176
$0-10^4$	100	9	1
10^4-10^5	5	8	0
$>10^5$	0	2	36

Total number of specimens, 161.

Urine Collection

The clean-specimen technique was found easy to perform and applicable to all patients with very few exceptions. It was no more time-consuming for the nursing staff than catheterization. The results obtained from the first specimen of urine from 102 consecutive patients are shown in Table II: 72% gave negative

TABLE II.—Results Obtained by Different Methods of Urine Collection

		Bacterial Counts per ml. of Urine			Total
		$0-10^4$	10^4-10^5	$>10^5$	
Clean-specimen technique	No. of specimens	74	12	16	102
	% of total	72	12	16	100
Tube method	No. of specimens	52	6	12	70
	% of total	74	9	17	100
Quinn and Kass (1960), mid-stream specimen, % of total		73	19	8	100

results, 12% were equivocal, and 16% were positive. Of the 12 patients whose urine samples gave equivocal results, two were subsequently shown to have infected urine, in nine the bacterial counts were negative, and one had persistently equivocal counts both in clean and in catheter specimens.

Fewer specimens were obtained by the tube technique, owing to the fact that over half of the 150 patients in whom this method was attempted were unable to pass urine while using the tube. It was found necessary to employ a bed stool, and this, plus the time spent in explanation and persuasion, greatly increased the work of the nursing staff. The results obtained from 70 consecutive patients' specimens are shown in Table II. These are similar to those obtained by the clean-specimen technique. Again there were no false-positive results. All positive cultures were subsequently confirmed by a catheter specimen of urine. For comparison, figures obtained by Kass (1960) are included in Table II; these show a good correlation. When the bacterial counts of 10^4 bacteria per ml. are examined in detail (Table III) it is apparent that there is a higher proportion of sterile cultures (56% as opposed to 32%) by the tube method, and that 91% of counts in this group are under 1,000/ml., as opposed to 81% with the clean-specimen technique.

TABLE III.—Comparison of Results Obtained from Clean and Tube Specimens of Urine with Bacterial Counts Less than 10^4 ml.

		Bacterial Counts per ml. of Urine			Total
		0	$0-10^3$	10^3-10^4	
Clean-specimen technique	No. of specimens	24	36	14	74
	% of total	32	49	19	100
Tube method	No. of specimens	29	18	5	52
	% of total	56	35	9	100

Comparison of Bacterial Counts with Conventional Reports

The results obtained when bacterial counts were compared with the reports given on the same specimen of urine using conventional methods of culture are shown in Table IV. There is agreement between these in 90% of urine samples. There were no false-positive results by either method, but the number of equivocal results is doubled when the conventional method alone is used. The two instances in which bacterial counts indicated infected urine but conventional reports were equivocal

were subsequently shown to be positive by both methods. No false-negative results were obtained on conventional reporting. The single negative result by this method which was equivocal on bacterial counting was subsequently shown to be negative by both methods. These results show that the conventional method in use at this hospital compares well with precise quantitative cultures, but does increase the number of equivocal results. It does not fail to detect "significant bacteriuria."

TABLE IV.—Comparison of Results Obtained by Bacterial Counting (Pour-plate Technique) and by Conventional Reporting

		Result on Conventional Reporting		
		Positive	Equivocal	Negative
Result on bacterial count	Positive	18	2	0
	Equivocal	0	8	1
	Negative	0	7	64

Discussion

The rationale of quantitative urine culture depends on the fact that when infection of the urinary tract occurs bacteria multiply in the bladder, so that urine samples regularly contain more than 10^5 organisms per ml. By contrast, bacteria from the urethra and vulva rarely produce counts in excess of 10^4 organisms per ml. provided care is taken in the collection of such specimens to ensure that contamination is reduced to a minimum.

We have studied several methods which have been suggested for the collection of mid-stream specimens from women. The method of Boshell and Sanford (1958) is too complicated for routine practice and has been justifiably criticized (*Lancet*, 1958). That of Hart and Magee (1957) we have also found impracticable. The Queen Charlotte's Hospital method described by Wood *et al.* (1960) is applicable to ambulant patients only, and demands considerable intelligent co-operation if a true mid-stream specimen is to be obtained. Furthermore, most patients consider the procedure distasteful. The method devised by Leather and Hutchings (1960) has the advantage that it can be used for ambulant and bed-ridden patients, and allows the collection of uncontaminated specimens during menstruation. Unfortunately, it is time-consuming, and successful in only 50% of our patients.

Kaitz and Williams (1960) have pointed out that the degree of bacteriuria due solely to contamination depends on three things: the volume of the urine specimen, the volume of the contaminating material, and the concentration of organisms in the latter. Thus a sterile urine sample of 100 ml. requires 10 ml. of contaminant containing 10^6 bacteria per ml. to give an overall bacterial count of 10^5 organisms per ml. Contamination of this degree is unlikely if the patient is carefully cleaned and then allowed to micturate into a sterile bed-pan. Special circumstances such as the presence of profuse vaginal discharge may militate against the success of this technique, but may be controlled by the insertion of sterile vaginal packs. We have compared the results obtained from mid-stream and "clean" specimens. The results show that the two methods give comparable results in respect of the proportion of infected, contaminated, or equivocal counts. No false-positive results were obtained by either method, but a larger number of sterile specimens were obtained by using the tube. These studies indicate that the sterile bed-pan technique not only provides satisfactory specimens but is more generally applicable to routine hospital

admissions, and is less time-consuming than methods designed to obtain mid-stream samples.

A number of methods for quantitative urine culture using surface media have been described (Hoeprich, 1960; O'Sullivan *et al.*, 1960; Bates, 1961). These all employ a standard inoculum which is distributed with a loop. The method we have adopted employs a glass rod spreader. This technique produces an even distribution of colonies on the plates which allows rapid and accurate enumeration. Colony crowding, which is a feature of all loop-spreading techniques, does not occur.

The use of the standard loop provides a rapid method for estimating the bacterial count with adequate accuracy over the critical range of 10^4 – 10^6 /ml. These advantages more than offset the single disadvantage of slightly decreased accuracy compared with the pour-plate technique.

Quantitative bacterial culture of urine has been widely accepted in America. Some authors (Kleeman *et al.*, 1960) go so far as to state that it is the only acceptable method of assessing the presence or absence of infection of the urinary tract. Kaitz and Williams (1960) state that 75% of patients shown to have bacteriuria exceeding 10^5 per ml. were missed by routine laboratory methods. However, quantitative urine culture has not been widely introduced in Britain, as few bacteriologists will accept that conventional techniques for the bacteriological examination of the urine fail to demonstrate urinary infection. To our knowledge no direct comparison has ever been made between bacterial counts and conventional methods of reporting the results of urine culture. The present study clearly demonstrates that the results obtained by these different techniques agree very closely. No cases of significant bacteriuria were overlooked by the conventional method; neither were any false-positive results obtained. The incidence of equivocal results is higher by the conventional method but is resolved by examination of further specimens.

It is of interest that the results using the standard loop were not significantly better than those obtained by the routine procedure. We would point out that assessment of routine cultures was carried out by an experienced bacteriologist, and we feel that with the use of a standard loop the enumeration of surface colonies may be more accurate where less experienced staff is available. Even so, it is apparent from our investigation that bacteriological methods in common use in this country do not fail to detect significant bacteriuria. There is thus no reason to believe that research programmes previously carried out in respect of infections of the urinary tract are in any way invalidated by failure to use bacterial counts.

Summary

The bacteriological examination of urine from female in-patients has been investigated with regard to (a) collection of specimens, (b) methods of quantitative culture, and (c) comparison of quantitative culture with routine laboratory reporting.

Methods recommended for collecting mid-stream urine specimens from female patients were investigated and, in general, found to be unsatisfactory. Collection of clean voided specimens in a sterile bed-pan after vulval cleansing is described. This method is suitable for patients of all degrees of infirmity. Urine samples collected in this way from 102 consecutive patients

proved as satisfactory as 70 specimens obtained by the more elaborate tube method described by Leather and Hutchings (1960). Furthermore, failure to obtain a specimen using the tube technique occurred in over 50% of the patients in whom it was attempted.

A technique for performing bacterial counts on surface plates using a fused platinum loop is described, and it compared with the standard pour-plate technique. Counts obtained by both methods agree in 89% of specimens examined, and all but one of the discrepancies were of a minor character. Thus the loop technique provides a simplified method of quantitative urine culture suitable for routine laboratories.

Bacterial counts have been compared with the independent routine laboratory reports on 100 consecutive urine specimens. The results agree in 90% of samples. In no case was significant bacteriuria overlooked by the routine laboratory. However, the use of bacterial counts halved the number of equivocal results. These findings cast doubt on the necessity for quantitative urine culture in routine practice.

We are pleased to acknowledge our indebtedness to Professor L. P. Garrod, Professor E. F. Scowen, Professor R. A. Shooter, and Dr. A. G. Spencer for advice and support in this work. Also to Sister K. Rawlinson and her nursing staff, whose enthusiastic co-operation made the study possible.

REFERENCES

- Bates, J. H. (1961). *J. Ark. med. Soc.*, **57**, 404.
 Boshell, B. R., and Sanford, J. P. (1958). *Ann. Intern. Med.*, **48**, 1040.
 Hart, E. L., and Magee, M. J. (1957). *Amer. J. Nurs.*, **57**, 1323.
 Hoepfich, P. D. (1960). *J. Lab. clin. Med.*, **56**, 899.
 Jackson, G. G., Griebel, H. G., and Knudsen, K. B. (1958). *J. Amer. med. Ass.*, **166**, 14.
 Kaitz, A. L., and Williams, E. J. (1960). *New Engl. J. Med.*, **262**, 425.
 Kass, E. H. (1955). *Amer. J. Med.*, **18**, 764.
 — (1956). *Trans. Ass. Amer. Physns*, **69**, 56.
 — (1957). *Arch. intern. Med.*, **100**, 709.
 — (1960). In *Biology of Pyelonephritis*, edited by E. L. Quinn and E. H. Kass, p. 399. Little, Brown, Boston.
 Kleeman, C. R., Hewitt, W. L., and Guze, L. B. (1960). *Medicine (Baltimore)*, **39**, 3.
Lancet, 1958, **2**, 515.
 Leather, H. M., and Hutchings, M. T. (1960). *Lancet*, **2**, 654.
 MacDonald, R. A., Levitin, H., Mallory, G. K., and Kass, E. H. (1957). *New Engl. J. Med.*, **256**, 764.
 Monzon, O. T., Ory, E. M., Dobson, H. L., Carter, E., and Yow, E. M. (1958). *Ibid.*, **259**, 764.
 O'Sullivan, D. J., Fitzgerald, M. G., Meynell, M. J., and Malins, J. M. (1960). *J. clin. Path.*, **13**, 527.
 Sanford, J. P., Favour, C. B., Mao, F. N., and Harrison, J. H. (1956). *Amer. J. Med.*, **20**, 88.
 Wood, E. C., Pinkerton, J. H. M., and Calman, R. M. (1960). *Brit. med. J.*, **1**, 961.

"A dean is, I suppose, like all others charged with authority over academic institutions. The list of qualities demanded of him by his colleagues will be varied and quite remarkably comprehensive. He will be fortunate if colleagues expect as the absolute minimum qualities something between the wisdom of Solomon, the patience of Job, the character of a saint, and the questionable knowledge of a sea lawyer. He will be unfortunate if he does not possess also the hide of a rhinoceros. He will be expected to swallow all the advice given to him by anybody at any time and yet make bold decisions with which nobody will disagree, and while being required to meet other conditions he must also act speedily and decisively. I sometimes think he is rather like the description of the Almighty by a Scottish preacher, who said: 'The Almighty is compelled to do many things in his official capacity which he would scorn to do as an individual.'" (Sir John Lockwood, inaugural address, 104th session, 1962-63, Royal Dental Hospital of London School of Dental Surgery.)

POSITIONS DURING LABOUR

BY

DONALD P. C. CHAN, M.B., F.R.C.S.Ed.
 F.R.F.P.S., M.R.C.O.G., M.M.S.A.

Lecturer in Obstetrics and Gynaecology,
 Hong Kong University

Even though various positions of the mother during the second stage of labour have been described and their respective advantages and disadvantages discussed (Moir, 1956; Claye, 1959), her position during the first stage of labour has been given little consideration in most of the standard obstetrical textbooks. She is usually encouraged to walk about or sit in a chair in the early stage of labour and before the membranes rupture, but most of the time she is allowed to lie in bed either in the dorsal or in one of the lateral positions. Theoretically, however, the erect posture appears to possess some advantages over that of lying flat in bed. In the sitting or standing-up position, the fundus of the uterus is thrown forward and thus the longitudinal axis of the birth canal is more straightened out. The uterine force can work at a better advantage and may therefore be more efficient. Furthermore, the descent of the foetus and the progressive dilatation of the cervix may be encouraged or assisted by gravity. The risk of inhaling vomitus, and hence of aspiration pneumonia or even death, would be reduced should general anaesthesia be found necessary during labour. On the other hand, the erect (sitting or standing-up) posture may be uncomfortable or even intolerable to the mother, especially during the first stage, and inconvenient or even impossible to the obstetrician, particularly during the second stage of labour. An attempt has been made to determine clinically if the erect posture is of any advantage, and, secondly, whether the maintenance of the erect posture during the first and second stages of labour is practicable.

Material and Method

Alternate primigravidae, in whom elective caesarean section would not be done, were assigned at random to one of two groups, A and B. The patients in group A were kept in the erect position: they were allowed to walk about, to sit in a chair, or to sit up in bed propped up with a bed-rest. The patients in group B, the controls, were kept in bed in the supine or lateral position. During the second stage the patients in group A were propped up to 45-60 degrees in the delivery bed and

TABLE I.—General Features of the Two Groups of Patients

	Group A	Group B
Age { Range	17-36	16-35
Average	25.16	25.23
Maturity { Range	33 4/7-43 1/7	32 5/7-45 4/7
Before 36 weeks	10	11
36 weeks or more	90	89
Complications { Uncomplicated oedema	25	19
Pre-eclampsia	10	12
Anaemia	13	15
Contracted pelvis	7	7
Twin pregnancy	1	0
Others	6	11
Dilatation of cervix { Closed-< 3 cm.	47	44
on admission { 3-5 cm.	44	40
5 cm. or more	9	16
Condition of membranes { Intact	68	67
on admission { Leaking	26	26
Ruptured	6	7
Presentation of foetus { Occipito-anterior	97	91
on admission { Occipito-posterior	2	5
Breech	2	4