first four years of an epidemiological study of 612 families randomly selected from two populations in South Wales.

No evidence is found in our data to suggest that arterial pressure is not a continuously variable characteristic, and in the absence of a sufficiently precise definition of essential hypertension to allow its genetic investigation in the general population we have used the appropriate system of quantitative analysis to investigate the simpler concept of multifactorial inheritance.

If arterial pressure is determined by graded polygenic inheritance it would be expected that the relationship in pressure between relatives and *propositi* would be similar at all ranges of pressure, and that the data would fit a linear regression more and more closely as the errors were eliminated. This is what we have found. On the basis of Mendelian inheritance of a specific gene for essential hypertension it would be reasonable to expect that the accumulation of more data, and more reliable data, would reveal internal inconsistencies in such an analysis and would segregate the population into at least two groups in which the relationship in pressure between relatives and *propositi* was different. We found no evidence of these.

It is possible that single-gene inheritance plays an important part in determining the arterial pressure of a very small fraction of the community, but our analyses suggest that in a representative sample of a general population which is experiencing what we take to be the usual morbidity and mortality associated with high blood-pressure this fraction is too small to be detected. For the majority of the population it appears that a regression of 0.287 for systolic pressures, and 0.224 for diastolic best summarizes the genetic resemblance of first-degree relatives, leaving between 55% and 77% of the systolic variance, and between 70% and 87% of the diastolic variance, for environmental factors to explain.

We gratefully acknowledge the continued co-operation of the families who are taking part in these studies, and the help of members of the survey team of the M.R.C. Epidemiological Research Unit who have assisted in the collection and analysis of the findings.

REFERENCES

- Hamilton, M., Pickering, G. W., Roberts, J. A. F., and Sowry, G. S. C. (1954). *Clin. Sci.*, **13**, 273.
 Miall, W. E. (1962). M.D. Thesis, London University.
 — and Oldham, P. D. (1955). *Clin. Sci.*, **14**, 459.
 — (1957). *Acta genét. (Basel)*, **7**, 114.
 — (1958). *Clin. Sci.*, **17**, 409.
 Morrison, S. L., and Morris, J. N. (1960). *Lancet*, **2**, 829.
 Oldham, P. D., Pickering, G., Roberts, J. A. F., and Sowry, G. S. C. (1960). *Ibid.*, **1**, 1085.
 Platt, R. (1959). *Ibid.*, **2**, 55.
 — (1960). *Ibid.*, **1**, 1189.
 — (1961). *Ann. intern. Med.*, **55**, 1.
 Roberts, J. A. F. (1959). *An Introduction to Medical Genetics*, 2nd ed. Oxford Univ. Press, London.
 Söbye, P. (1948). *Op. dom. Biol. hered. hum. Kbh.*, Vol. 16.

"The Air Force Surgeon General's office [U.S.A.] has ordered Air Force hospitals and clinics to stop distributing free cigarettes to patients. A similar prohibition has been ordered on the inclusion of cigarettes in the packaged lunches that are prepared for service personnel on long flights. Major-General R. L. Bohannon, the Deputy Surgeon General, said in a strongly worded directive that the 'ever-increasing evidence' of a link between cigarette smoking and cancer and certain other diseases 'no longer can be ignored.'"—*New York Times*, October 8.

BRONCHIECTASIS IN CHILDHOOD

BY

NORMAN S. CLARK, M.D., B.Sc., M.R.C.P., D.C.H.
Senior Lecturer in Child Health, University of Aberdeen

Bronchiectasis, while not a very common disease in childhood, is by no means rare, often passes unrecognized for many years, and is responsible for a sum-total of chronic and recurrent ill-health quite out of proportion to the actual number of its victims. The natural history of this disease, as of most chronic conditions, is not easy to study, and our knowledge thereof is incomplete and based more upon impressions than objective observations.

So far as I know, only two comprehensive surveys of bronchiectasis in childhood have been published from this country in recent years (Field, 1949; Strang, 1956). While both contain much useful information, neither is based on a completely unselected series of cases. Field's cases were collected at the Hospital for Sick Children, Great Ormond Street, and at University College Hospital, London. Neither of these hospitals draws all the cases from any defined population group and both probably tend to attract the difficult or unusual case. Strang's report is based on the results of a follow-up of children admitted to Newcastle Regional Thoracic Surgery Centre: such a series is likely to be deficient both in cases with disease too extensive for surgery and in cases with few symptoms or symptoms well controlled by medical treatment.

The survey here reported was undertaken in the hope of building up a comprehensive picture of bronchiectasis in childhood and of assessing the end-results of treatment in an unselected series of cases.

Material

This survey covers all children aged 0 to 11 years inclusive in whom a diagnosis of irreversible bronchiectasis was first established in the Royal Aberdeen Hospital for Sick Children during the 10-year period 1946–55. This is the only children's hospital in the North-eastern Hospital Region of Scotland, an area which includes the City and County of Aberdeen and the Counties of Kincardine, Banff, Moray, Orkney, and Shetland. It is certain that during this period no children have been admitted for bronchography direct to the Regional Thoracic Surgical centre or to any other hospital in the Region. The regional boundaries are well defined and it can be claimed with some confidence that few, if any, cases living in this region have been investigated in adjacent regions. Indeed, our figures include a number of cases transferred from the Northern Hospital Region as possible candidates for thoracic surgery. This series therefore includes all cases of bronchiectasis confirmed in children from the North-eastern Hospital Region during a 10-year period. I have personally examined all these children and have taken part in their supervision throughout the whole period covered by the survey.

I have excluded from further consideration three cases of bronchiectasis due to mucoviscidosis (fibrocystic disease) and one case of congenital cystic disease of the lung presenting with tension cysts in the neonatal period (Clark *et al.*, 1956). No other case has been excluded as "congenital cystic disease."

Incidence

During this 10-year period the diagnosis of bronchiectasis has been confirmed in 116 cases (57 boys, 59 girls), by necropsy in one case and by bronchography in the remainder. For reasons which will be apparent, it was clear that the great majority had irreversible disease when the diagnosis was made, and time has confirmed the irreversibility of the remainder.

Of these 116 children, 10 were transferred from the Northern Hospital Region and 106 belonged to the North-eastern Hospital Region. The child population of this region aged 0 to 11 years at the 1951 census was 99,556, so that the average annual incidence rate was 1.06 per 10,000 children. If the incidence in this region is representative of that in Great Britain as a whole, approximately 1,000 new cases should be recognized in childhood each year in Great Britain.

Clinical Features

Age at Onset.—Table I shows the age at onset of persistent respiratory symptoms in these children. The evidence suggests that bronchiectasis has in fact been

TABLE I.—*Age at Onset of Symptoms in 116 Cases*

Age	0-2	3-5	6-10	Uncertain
No. of cases ..	50	34	18	14*

* At least 5 had symptoms before age 6.

present from this age in most cases, though proof of this is possible in only a few. These figures, like those of Strang (1956), again emphasize that bronchiectasis most commonly arises in the early years of childhood. It is, however, unusual for bronchiectasis to be recognized in its early stages. Of these 102 children, 35 had a history of symptoms continuing for over five years before the diagnosis was made, 15 had a history of four to five years, and in only 25 had the symptoms arisen in the year prior to diagnosis. Field (1949) also comments on the number of children whose symptoms had persisted for years before they were referred for investigation.

Initial Illness

The respiratory symptoms were said to date from an attack of pneumonia in 26 cases, pertussis in 24, and measles in 16. Post-operative atelectasis appeared to have initiated the bronchiectasis in three cases and inhalation of a foreign body in two. In the remaining cases no clear history of an initial illness was given. These figures are in accord with the general views on aetiology and emphasize once again the importance of pneumonia and pertussis. The proportion of cases following measles in this series is higher than is sometimes suggested: of Field's (1949) cases only 3% followed measles.

Consideration of the 25 cases with a history of symptoms for less than a year before the diagnosis was established does not support the suggestion of Williams and O'Reilly (1959) that there are two distinct groups of cases, one in which bronchiectasis is the result of subacute pyogenic pulmonary collapse and one in which it is a sequel to non-specific infective bronchiolitis and/or interstitial pneumonia. Of these 25 cases, 22 had a well-defined initial illness, but nine of them showed no evidence of atelectasis: the bronchiectasis subsequently demonstrated in these 22 cases varied from the very localized to the very diffuse, and the subsequent

course of the disease could not be related to the presence or absence of initial atelectasis nor to the type of the initial illness.

Although calcification of the hilar nodes is present in three children, there is no evidence that tuberculous infection has played any part in the genesis of bronchiectasis in these cases. This is in keeping with the findings of both Field (1949) and Strang (1956). Though it is generally agreed that tuberculous hilar adenitis can lead to atelectasis and subsequent bronchial dilatation, it has been shown (Kent, 1942; Graham and Huichison, 1947) that such dilatations commonly remain free from secondary infection and are consequently asymptomatic, and that tuberculosis plays a very small part in the aetiology of infected bronchiectasis.

Symptoms and Signs

Cough was, of course, the commonest symptom and was described as persistent in 94 and intermittent in 14. In the remaining eight cough, though present at times, was not a prominent symptom. It is commonly stated in textbooks that children with bronchiectasis do not produce sputum until trained to do so, but in this series 67 had a history of purulent expectoration. Haemoptysis had occurred in five cases only and in all of them had been slight. Field's (1949) figures for this symptom are similar (7%); but Strang (1956) found an incidence of 25%, and in some textbooks (Brenneman, 1959) haemoptysis is described as a frequent symptom. This difference of opinion is surprising in respect of a symptom such as haemoptysis which is immediately obvious and which commonly alarms parents so that they are unlikely to forget it.

The only other symptoms complained of were dyspnoea on exercise in 12 cases and wheezing in nine. The low incidence of wheezing in the present series is remarkable; both Field (1949) and Strang (1956) noted wheezing as a symptom in approximately 25% of their cases.

Medium rales, often described as "crackling" or "bubbling," localized to one or more areas of one or both lungs, and heard at the same site or sites on repeated examination, were present in 105 cases. It is the persistence rather than the character or location of these adventitious sounds which should suggest the diagnosis of bronchiectasis. In many cases no other abnormal sign was noted on physical examination of the chest. Alteration of the breath sounds either in character or in intensity was present in 24 cases and impaired percussion note in 20. Deformity of the chest wall, usually amounting only to slight flattening of one side of the chest, was present in eight cases, and in one of them the deformity was probably due to rickets. Five children showed clinical evidence of mediastinal shift. Persistent signs of coexisting generalized bronchitis found by Helm and Thompson (1958) in 10 out of 32 children were rare in the present series.

Persistent cyanosis was not seen in any case in this series and finger-clubbing was present in only 39, including seven with doubtful minimal clubbing. While there is no doubt that clubbing is more often seen in those with heavily infected cystic bronchiectasis and in those with extensive bilateral disease, this relationship is not constant, and definite clubbing was present in one child with tubular disease affecting only two basal segments of the left lower lobe. Field found clubbing in 44% of her cases and Strang in 51%.

The general physique of these children varied greatly, but most of those with extensive disease were much below their expected weight. In 45 children the weight on first examination was below that of the 10th percentile in the Iowa weight and height tables (Nelson, 1959). Nevertheless, bronchiectasis of moderate extent was found in children of good and indeed better than average physique.

The degree of disability varied considerably and bore no close relationship to the extent, site, or type of bronchiectasis present. Thirty-five children were reported to suffer from acute respiratory illnesses sufficient to incapacitate them several times a year; 75 complained only of persistent cough with occasional acute flare-up; three appeared to have no real disability; and in three the history is not clear on this point.

Associated Upper Respiratory Infection.—The high incidence of chronic or frequently recurrent upper respiratory infection in bronchiectatic children has long been recognized. Many of the children in this series had a long history of upper respiratory symptoms and a considerable number had undergone investigation and treatment in E.N.T. departments before the presence of bronchiectasis was recognized. Bilateral antral lavage was carried out in 87 of these children in the course of their initial investigation and in 77 (88%) of them pus or mucopus was found in one or both antra.

Investigations

Allibone *et al.* (1956) found *Haemophilus influenzae* the predominant organism in bronchoscopic aspirations and/or sputum in all of 32 bronchiectatic children investigated, and concluded that this organism was responsible for keeping the chronic inflammatory process smouldering. The present investigation lends no support to this theory: cultures were made from 114 specimens of sputum obtained from 68 children during their first hospital admission for bronchiectasis: *H. influenzae* was isolated on only 14 occasions, the remaining specimens showing a mixed, mainly coccal, flora with no one organism predominant.

The results of the initial plain x-ray films of the chest were as follows: in one case the whole left lung was collapsed and airless apart from several cystic spaces which showed fluid levels; cavitation with or without evidence of atelectasis was the most prominent feature in 12 cases; lobar atelectasis was present in 34 cases and segmental atelectasis in five; ill-defined patchy opacities were present, usually in the basal areas, in 46; the films were passed as normal or as showing only doubtful increase of the basal markings in 16 cases; and in two the original films were missing.

Bronchoscopy was performed only in the small group of cases with a short history who showed atelectasis, presumably of recent origin. While it may play a useful part in the treatment of the early reversible stage of bronchial dilatation it probably adds no useful information in the investigation of the established disease (Field, 1949).

Bronchography

Bronchography was carried out in every case except that of the patient who was admitted unconscious and died within a few hours (see below). In seven children in whom the diagnosis was in no doubt from a very early age bronchography was postponed for several years, but in all others this procedure was carried out

during the initial investigation. It has been our almost invariable practice to outline the bronchi of each lung at separate sessions so that true lateral films could be taken as well as obliques. Satisfactory outlines of the whole bronchial tree were obtained in all but three patients, for whom incomplete bronchograms are available. To obviate observer variation I have personally reviewed all the bronchograms without reference to the original radiologist's report, and the findings to be detailed are all based on my own interpretation of the films.

Bilateral bronchiectasis was demonstrated in 61 cases, and the disease was confined to the left lung in 40 and to the right lung in 12 cases. In the remaining three bronchiectasis was demonstrated on one side only, but the bronchograms are incomplete. Bilateral disease was almost as common in Field's (1949) series, but Strang (1956) found both lungs involved in only 28% of his cases: this low incidence is almost certainly the result of selection, as his cases were seen in a thoracic surgical unit at a time (1935–48) when surgery was less often employed in the treatment of bilateral disease.

Table II shows the relative frequency of involvement of the different lobes. (In two cases of situs inversus the right lung was recorded as "left" and vice versa for the purpose of this table.)

Involvement of:	No. of Cases
Right upper lobe	23 (20%)
" mid-lobe	58 (50%)
" lower lobe	45 (39%)
Left upper lobe	12 (10%)
Lingula	85 (73%)
Left lower lobe	89 (77%)
Total lobes involved	312

The distribution of the disease shown in Table II is generally similar to that previously reported, though the incidence of right lung involvement is lower and of lingular involvement higher than that found by Field (1949). It is generally accepted that bronchiectasis is predominantly a disease of the left lower lobe, lingula, and right mid-lobe and that this distribution is largely determined by the difficulty with which secretions drain from these lobes. In only 18 cases was the disease confined to one lobe, and in 57 cases more than two lobes were involved. No case showed the disease confined to one upper lobe, but with this exception every lobe and nearly every possible combination of lobes is represented in this series. By far the commonest combination was left lower lobe and lingular involvement, which was present alone or in combination with other lobes in 71 cases.

Bronchiectasis is often further subdivided according to whether the dilatations are predominantly "tubular," "saccular," "varicose," etc. I do not believe that such subdivision serves any useful purpose. Tubular and saccular bronchiectasis co-existed in 12 out of the 23 cases which showed saccular dilatation, and in four cases dilatations which appeared tubular at the initial bronchogram had become saccular at repeat bronchography several years later. I have found no evidence in this series to suggest that the different morphological types are produced in different ways, and in my view saccular or cystic dilatations indicate only a more severe degree of damage to the bronchial wall. In most cases saccular dilatations certainly appear to be more heavily infected and to produce more purulent sputum, but this was reflected in the degree of disability only in one or two cases with very extensive disease.

Management

During the whole of the 15-year period from the start of this survey our policy has been to recommend lung resection except in children with few symptoms and limited dilatations of slight degree and in cases with disease so extensive that resection of all affected parts of the lungs was clearly out of the question. Limited resection of the most severely affected areas has been offered to a few children in the second category. There has been close co-operation throughout between the staff of this hospital and that of the Regional Thoracic Surgical Unit at Woodend Hospital, Aberdeen, where all the surgical treatment has been carried out.

When the diagnosis has been established and the distribution of the disease determined, it has been our practice to interview the parents, explain the nature of the disease, and demonstrate to them the appropriate position for postural drainage. The enthusiasm and efficiency with which postural drainage has been carried out at home has, of course, varied considerably from case to case, but for the best results it is essential for the physician to spend some time explaining and demonstrating to parents the proper method. It is useless simply to suggest that the child should hang over the side of the bed. It is a considerable athletic feat for anyone to maintain himself in an adequate head-down slope over the side of a bed for any length of time. Support must be provided, and in the majority of patients, who can best be drained in a face-down or partly rotated face-down position, a tipping-board is essential.

We have shown the parents a simple tipping-board and advised them to make one for the child's use at home. This is well within the powers of even the most unhandy father, and the cost is only that of a few pieces of wood and two pairs of hinges. It is not, however, sufficient to provide a tipping-board and leave it at that. The novelty of postural drainage palls rapidly and the average child left to himself on a tipping-board will shortly be found comfortably propped on his elbows. It is essential that an adult should supervise the procedure from beginning to end, encouraging the child to cough actively and applying firm percussion intermittently to the lower ribs. We have advised at least two 20-minute periods of postural drainage daily continued throughout childhood or until lung resection is completed. Posturing during sleep by raising the foot of the bed has been used only in one or two cases with much purulent secretion. In my experience this is seldom a very useful procedure in childhood, as the child usually wriggles round during sleep into a more or less transverse position.

We have advised the family doctor to restrict the use of antibiotics to the treatment of any acute flare-up of infection in order to minimize the risk of persistent antibiotic-resistant infection.

With increasing experience our surgeons have tended in recent years to postpone segmental or lobar resections to later childhood or adolescence in order to lessen the risk of post-operative atelectasis in the remaining lobe or segments. Pneumonectomy has been performed on five children aged between 4 and 10 years. We have been less hesitant in advising early operation in these cases, as post-operative complications in the remaining healthy lung are uncommon: in all five a heavily and

persistently infected lung was an additional argument for early operation.

Except in children who have come to operation within 12 to 18 months of the initial bronchography, we have repeated the bronchograms on one or both sides shortly before operation. Repeat bronchography has also been performed in some of the medically treated cases. In addition, post-operative bronchograms of the affected side or sides have been carried out at intervals of six months to one year after operation in all but 13 of the surgically treated cases. Including the initial bronchograms a total of 532 separate bronchographic examinations were performed in these children up to the end of July, 1961 (each examination covering one lung only). As with the initial bronchograms I have personally reviewed all the films without reference to the original radiologist's report in order to keep the standards of interpretation as constant as possible.

Results of Repeat Bronchography

In 79 children bronchography has been repeated on one or both sides (in 14 of them on two occasions) without operation intervening since the original examination. The interval between the first and the last bronchogram was between six months and three years in 33 patients and between three and eight years in 46. The results of these examinations throw some light on the still debated question of whether bronchiectasis is a progressive disease.

In several paediatric textbooks (Grulee and Eley, 1948; Paterson and Moncrieff, 1949; Ellis, 1956; Nelson, 1959) the view is expressed that in bronchiectasis there is commonly a tendency for the dilatations to increase in calibre and to spread to hitherto unaffected parts of the lung, though no evidence is advanced in favour of this view. The opposite opinion that bronchiectasis is rarely a spreading disease has been repeatedly advanced in a succession of original papers on the subject, again in some cases with remarkably little concrete evidence in its favour.

Churchill and Belsey (1939) express the view that "bronchiectasis usually has reached its full extent and distribution at the time the diagnosis is made" and suggest that the opposite view has had undeserved support as a result of "incomplete bronchograms in varying portions of the lung at different examinations." Perry and King (1940) reviewed 114 patients after an interval of 1 to 19 years and found evidence of minimal spread in only six. It is not clear, however, whether this review included complete repeat bronchographic examination. Diamond and Van Loon (1942) followed 75 bronchiectatic children for periods of up to eight years. Though they give no details it appears that bronchography was repeated in many and possibly in all of them: two showed some increase and one some decrease in the size of the affected bronchi, but in none was there evidence of spread to previously uninvolved bronchi. Franklin (1944) followed 24 affected children for periods of 4 to 12 years and found that in one case tubular dilatation had become saccular and that in another the disease had spread to the other lung. Again, it is not clear whether bronchography was repeated in every case.

Strang (1956) in a follow-up of 209 children and Wynn-Williams (1957) in a report on 217 patients, mostly adult, were of the opinion that bronchiectasis was rarely

a spreading disease but quoted no bronchographic evidence in support of this.

On the other hand, Field (1949), found as a result of a considerable number of repeat bronchograms that there was "a greater tendency for the dilatation to get worse or remain stationary rather than improve." Ginsberg *et al.* (1955), reviewing the results of surgical treatment of 221 patients, nearly all adult, noted that about one-eighth of them had shown extension of the bronchiectasis but do not say in what proportion the extension may have been a post-operative phenomenon. Of the authors quoted only Diamond and Van Loon (1942) and Field (1949) seem to have performed repeat bronchograms on any considerable number of children, and their results appear to be mutually contradictory. I have so far found no other report which presents any evidence bearing on this problem.

In the present series of 79 children the bronchograms in 46 cases showed no significant change in the extent or degree of the dilatations. Twenty-seven cases showed deterioration and seven improvement. (One case appears in both the last two groups as one lung showed deterioration and the other improvement.) Of the 27 cases which showed deterioration this consisted of an increase in calibre of already affected bronchi in 12; this cannot be considered an important change, and is, in any event, difficult to assess, as allowance must be made for growth and in some cases for slight differences in the phase of respiration at which the films were taken. The remaining 15 cases showed involvement of 17 segments which had previously appeared normal; the newly affected segments were distributed as follows: right mid-lobe 3; right lower lobe, apical 2; right lower lobe, basal 3; lingula 1; left lower lobe, apical 3; left lower lobe, basal 5. In seven of these 15 cases the extension of the disease was shown to persist at a third bronchogram one year or more later.

In the seven cases showing improvement, eight previously affected segments now appear normal—in the right upper lobe 2, right mid-lobe 2, right lower lobe 1, lingula 1, left lower lobe 2. In five of them it is fairly certain that bronchiectasis had been present for some years before the first bronchogram, but in the other two it is possible that the condition was of six months' duration or rather less. In spite of the improvement all are left with bronchiectasis of significant extent.

When factors which might be related to a tendency to deterioration are considered it appears that early onset is not significant. Deterioration is certainly not confined to the early years of childhood, as in seven cases the first bronchogram was performed at the age of 9 years or over. As might be expected, changes are rather commoner in cases where the disease appears to be of relatively recent onset, though changes are by no means confined to the earlier years. Deterioration is not related to the original extent of the disease, but when the distribution is considered it appears that a higher proportion of those showing deterioration had original involvement of an upper-lobe bronchus (Table III).

If we discard the 12 cases showing only increase in calibre of the affected bronchi as being of doubtful significance, there remain 15 of the 79 cases who have shown definite spread of the disease. This is surely not surprising if it be accepted that the combination of bronchial occlusion and infection is the usual precursor of bronchial dilatation. Sticky purulent secretion is

almost constantly present in the affected bronchi in bronchiectatic children and must not infrequently spill over into normal bronchi with changes of posture. While this often causes a bout of coughing, it would seem probable that from time to time it will fail to

TABLE III

No Change (46 Cases) Lobes Originally Involved				Worse (27 Cases) Lobes Originally Involved			
R.U.L.	..	..	6 (13%)	R.U.L.	..	..	8 (30%)
R.M.L.	..	..	26 (57%)	R.M.L.	..	..	16 (59%)
R.L.L.	..	..	17 (37%)	R.L.L.	..	..	9 (33%)
L.U.L.	..	..	1 (2%)	L.U.L.	..	..	5 (19%)
Lingula	..	..	39 (85%)	Lingula	..	..	19 (70%)
L.L.L.	..	..	35 (76%)	L.L.L.	..	..	19 (70%)
Totals .. 124				76			

clear sticky secretion from the small-calibre bronchus of a child. If the bronchial occlusion persists everything is then in favour of the development of a fresh patch of bronchiectasis. This, like all bronchiectasis, probably passes through a reversible stage, and if the airway is re-established at this stage the bronchus may revert to normal. Presumably in the seven cases which showed improvement the bronchi concerned were in this stage at the time of the first bronchogram though the disease as a whole was irreversible. The relative frequency of upper-lobe disease in cases showing deterioration is in keeping with this hypothesis, for infected secretions from an upper lobe are more likely to drain into the lower part of the lungs.

It is suggested, therefore, that bronchiectasis in children, even when it is known to have been present for some years, should not be regarded as a static condition in which the only change to be expected is waxing or waning of the infection in an already finally determined area of bronchial dilatation. Both spread and regression of the disease are not uncommon, and the findings in this series support the view of Field (1949) that the condition is more likely to become worse than better. Increase in the extent of the lung involved does not, of course, imply that the symptoms will become worse. Nevertheless, if surgical treatment is contemplated the possibility of extension of the disease is a factor that must be borne in mind. In most cases the risks of spread are probably less than the risks of post-operative complications if segmental resections are carried out in a young child, but in the occasional case with upper-lobe involvement and already extensive disease the question of expediting operation should be seriously considered. It is quite clear that in all cases it is wise to repeat bronchography before surgical treatment unless this is carried out soon after the diagnosis is established.

Results

All patients, except those from the Orkney and Shetland Islands and from the Northern Hospital Region, have been seen at intervals at the follow-up clinic at the Royal Aberdeen Hospital for Sick Children until age 12 or until operation at an earlier age. Thereafter, most have been seen at intervals at the thoracic surgical unit. For the purpose of the present investigation I have endeavoured during the period June, 1959, to December, 1960, to trace and examine personally all surviving patients in the series, and the assessment of the final state of these patients is based on the results of this examination and consideration of the most recent bronchograms. It proved impossible to visit or to bring

back to Aberdeen for examination most of the patients living in the Northern Hospital Region or in the Orkney and Shetland Islands. Questionnaires were sent to the family doctors of these patients and in most cases adequate information was obtained for assessment. A similar questionnaire in respect of one boy who had gone to live in England was completed by the local chest clinic, which he had attended regularly.

This method of final assessment by a physician or by the family doctor rather than by the thoracic surgeon was chosen deliberately for two reasons. Firstly, the same standards were applied, in most cases by the same person, irrespective of whether the case had been treated medically or surgically; and, secondly, it was felt that there was some substance in the criticism: "A surgeon's estimate of his own results is liable to be unconsciously influenced not only by his own enthusiasm but also by the natural wish of many patients to gratify one who had made strenuous and dramatic efforts to relieve them" (*Lancet*, 1955). Twenty cases could not be assessed in this way either because they could not be traced or because they refused to attend for examination and no report was received from their family doctors. The final state of these 20 patients has been assessed on the record of their last visit to the thoracic surgical unit and on the most recent bronchogram. I had intended to exclude these cases from the final analysis of the results of surgical treatment, but in fact this proved unnecessary as their inclusion made no significant difference to the final results. Indeed, if anything, the results of surgical treatment appeared slightly poorer in this group than in the cases reassessed by myself or their family doctors.

Deaths

Three children are known to have died from the direct effects of the disease, all in the earlier years of the survey. A girl aged 11 years was admitted moribund and died within 24 hours from rupture of a large cerebral abscess into the lateral ventricle. She had no history suggestive of bronchiectasis, but necropsy showed heavily infected atelectatic bronchiectasis of the left lower lobe resulting from bronchial obstruction by an inhaled fragment of pine branch. Two boys aged 5 and 6 years died at home of "pneumonia" six months and three years respectively after establishment of the diagnosis. Both had heavily and persistently infected bronchiectasis, in one involving the right mid and lower lobes, and in the other involving the whole of both lungs except for the upper lobes.

This is probably the total mortality from the disease during the period of the survey apart from two post-operative deaths, which are considered in a later section. Only three children were completely untraced after their initial admission; two children were followed for only six months after successful operations in the early years of the survey and were thereafter untraced. Nine were followed for varying periods and were well when last seen in 1955 or later. The remaining 97 children were known to be alive at the final follow-up in 1959-60.

The three fatal cases have been considered separately as they cannot fairly be included in either the medically treated or the surgically treated groups. The girl was moribund on admission and the two boys failed to report back, and we have no information on whether medical treatment was in fact carried out at home.

D

Of the 110 children remaining after excluding the three who died and three untraced, 75 have completed surgical treatment, 5 have had unilateral resection and are awaiting operation on the other side, 2 are awaiting operation, and 28 have been treated medically.

Results of Surgical Treatment

Deaths.—A total of 104 separate resections have been carried out on the 80 children who have so far come to operation, with two post-operative deaths (operative mortality 1.9%, case mortality 2.5%).

Post-operative Complications

The frequency of significant non-fatal post-operative complications is shown in Table IV. Only 46 operations were followed by a completely uncomplicated convalescence.

TABLE IV

Segmental or lobar collapse	..	..	42
Pneumothorax	..	..	6
Sterile effusion	..	..	5
Empyema	..	..	1
Broncho-pleural fistula	..	..	1
Severe tracheitis	..	..	1
			<hr/> 56

Bronchoscopic aspiration and physiotherapy resulted in rapid re-expansion of the collapsed lobes or segments in all except three cases. In these three bronchiectasis developed in the collapsed area, which was subsequently successfully resected. The incidence of post-operative collapse was higher in younger children and in children with bronchiectasis in the contralateral lung. If the five pneumonectomies, none of which was followed by collapse, are excluded, 11 out of 19 operations on those aged 6 to 8 years were followed by collapse, as compared with 16 out of 37 on those aged 9 to 11 years, and with 15 out of 41 on those aged 12 years and over. In 42 cases in which bronchiectasis was present in the opposite lung at the time of operation, 24 developed post-operative collapse. In 55 cases in which the opposite lung was healthy or in which bronchiectatic segments on the opposite side had previously been successfully removed, only 18 developed post-operative collapse.

Broncho-pleural fistula occurred in one child after resection of the lingula and basal segments of the left lower lobe. When the fistula was outlined with lipiodol it was found to communicate with the apical segment of the left lower lobe, which now showed gross bronchiectasis: the fistula was excised with the affected segment without further complications, and subsequent bronchography showed a normal very well expanded left upper lobe.

None of the other complications was followed by any lasting sequelae.

Post-operative Bronchograms

Bronchograms were repeated within one year of operation in 62 children. The 13 patients in whom no post-operative bronchograms were carried out include the five treated by pneumonectomy, in all of whom the opposite lung had been free from disease. In 33 cases the post-operative bronchograms were normal and complete—that is, if resection was performed on both lungs bronchography was repeated on both sides. In a further three cases subjected to bilateral operation bronchography on one side gave a normal result but the other lung was not re-examined after operation. In 18 cases residual bronchiectasis was demonstrated, and in

eight there had been spread of the disease to previously healthy segments in the same lung as the most recent resection. In 12 of the cases with residual disease this was minimal and confined to one lung. Their symptoms after the original resection on the more severely affected side were not thought sufficient to warrant operation on the second side. In six cases the disease was too extensive to allow of complete extirpation, but resection of the most severely involved lung on one or both sides was carried out as a palliative measure. In the eight cases that showed spread of the disease the symptoms were regarded as severe enough to justify further operation in three (see previous section): in one the apical segment of the right lower lobe was removed and in two the left upper lobe after previous removal of the left lower and lingula. In five cases the dilatations were slight, were confined to one segment, and produced no significant symptoms.

Final State

In one patient who showed considerable improvement after her first operation, a second resection had been carried out too recently to allow of final assessment. Final examination of the remaining 72 patients was made between 6 months and 12 years after completion of surgical treatment. The follow-up period was from six months to three years in 24 cases, from four to six years in 24 cases, and from 7 to 12 years in 24 cases. Patients were classed as "recovered" only if the following criteria were all satisfied: (1) There was no persistent cough. (2) No sputum was produced except during an acute respiratory infection. (3) Acute respiratory infections were neither more frequent nor more severe than in the average healthy person. (In assessing this point the parents were asked to compare the child with his friends, with other members of his class, and with other members of his family. Whether the answer was "Yes" or "No," few parents were in any doubt when they were questioned on this point.) (4) Exercise tolerance was good—that is, the patient was able without distress to take part in games and activities normal for his age. (5) There were no abnormal signs on clinical or radiological examination other than those indicating diminution of lung volume. No patient was included in this group unless he was himself entirely satisfied with his present state. Those classed as "much improved" would have qualified for the "recovered" group but for slight dyspnoea on exercise or a slight but persistent cough with occasional mucoid or mucopurulent sputum. In a number of cases the latter symptom might more properly be ascribed to persistent nasal sinus infection than to residual pulmonary disease; nevertheless it probably gives a fairer picture if such cases are excluded from the "recovered" group. The group classed as "improved" still had significant symptoms though these were in no case disabling, and all were satisfied that there had been a real improvement in their symptoms. Most of them produced a little purulent sputum from time to time and a number of them still had persistent moist sounds in the chest. The one patient classified as "unchanged" still had a persistent productive cough nine years after operation: signs of infection in his chest were minimal and some of his symptoms might be attributable to persistent nasal sinus infection.

Table V shows the end-results in the 75 cases surgically treated. In Table VI the results in the 72 survivors who were followed up are further analysed according to

whether there is bronchographic evidence of residual bronchial dilatation. Those with no such evidence are further subdivided into cases with normal post-operative bronchograms and those in whom post-operative bronchography was not carried out but in whom all the affected lung tissue was resected. The latter group includes the five cases in which pneumonectomy was performed.

TABLE V

Bronchiectasis	Dead	No Follow-up	Un-changed	Improved	Much Improved	Recovered	Total
Unilateral	0	0	1	7	5	24 (65%)	37
Bilateral	2	1	0	11	7	17 (45%)	38
All cases	2	1	1	18 (24%)	12 (16%)	41 (55%)	75

TABLE VI

	Un-changed	Improved	Much Improved	Recovered	Total
Residual or new bronchiectasis demonstrated	1	9	7	9 (35%)	26
Post-operative bronchograms normal	0	6	2	25 (76%)	33
No post-operative bronchograms but all affected lung tissue removed	0	3	3	7	13
All cases	1	18	12	41	72

Results of Medical Treatment

One patient cannot be classified as her initial bronchograms were technically unsatisfactory. She has complete transposition of the viscera, with collapse of the right lower lobe and cystic bronchiectasis in the right upper lobe and possibly also in the other lung. She lives in the Shetland Isles and has not been seen here since her original admission. Her doctor reports that now at the age of 16 she is considerably incapacitated with very poor exercise tolerance and persistent cough productive of 6 oz. (170 ml.) of purulent sputum daily.

The remaining 27 patients treated medically do not form a homogeneous group and cannot, of course, be directly compared with the surgically treated group. Six patients were thought to be suitable for operative treatment, but this was declined: in 12 the symptoms were not sufficient to warrant surgical treatment, and in nine the disease was so widespread that operation was considered impracticable. Three additional patients originally placed in this last group were after reconsideration submitted to resection of the most severely affected segments. It is possible that one or two others will ultimately graduate from this group to the surgically treated group.

These 27 children have been followed for periods of between 3 and 13 years from the time of diagnosis. The follow-up period was from three to six years in 10 cases and from 7 to 13 years in 17. Table VII shows their status at the time of the last examination. These figures are based on the same criteria as were used in the surgically treated cases.

TABLE VII

	Worse	Un-changed	Im-proved	Much Improved	Re-covered	Total
Refused operation	1	0	4	0	1	6
Operation not indicated	0	1	6	2	3	12
Inoperable	2	1	6	0	0	9
	3	2	16	2	4	27

The number of cases in this group is unfortunately too small for any significant conclusions to be drawn. Twenty-two have shown significant symptomatic improvement, but with one exception only those in the group with disease of limited extent have reached the "recovered" or "much improved" categories. Bronchography has been repeated in 16 cases: eight showed no change, three improvement, and five deterioration (these cases are included in the 79 discussed in the previous section on the results of repeat bronchography).

Discussion

In a succession of leading articles in the *Lancet* (1955, 1958a, 1958b) it has been suggested that the number of cases of bronchiectasis suitable for and requiring resection is relatively small, that the results of surgery are frequently disappointing, and that medical treatment gives adequate results in the majority of cases. While those articles are concerned with the problem in all age-groups it is suggested (*Lancet*, 1958b) that "in childhood lengthy observation is essential, and at the end of it the surgical harvest is likely to be small."

With the first half of this statement there would be fairly general agreement in view of the high rate of post-operative complications in young children following lobar or segmental resection. I would, however, make exception in the occasional case (five in this series) with extensive heavily infected cystic disease in one lung and a normal lung on the opposite side. Pneumonectomy was well tolerated by these five children, there being only one significant post-operative complication (empyema), with complete recovery in one case and very striking improvement in the other four. The possibility of spread to the other lung is, I think, sufficient to justify early operation in such cases. The second half of the statement that "at the end of it the surgical harvest is likely to be small" is, I believe, much more debatable. Surgical treatment was certainly possible in 90% of the cases in this unselected series and was considered both possible and advisable in 75%. Of the cases submitted to surgery 55% were classed as completely recovered and a further 16% as much improved, a total of 71% with what may fairly be termed very satisfactory results. Against this an operative mortality of 1.9% and a case mortality of 2.5% cannot be regarded as prohibitive risks and can almost certainly be improved upon with further experience.

Though there are many reports in the literature of the result of surgical treatment of bronchiectasis in adults, I have found comparatively few which refer only to children. Of 202 children followed up by Field (1949), 93 had been treated surgically, with eight post-operative deaths. In 50 patients in whom resection of affected lung tissue was complete, 54% were classed as cured and 32% as improved. In 43 cases with incomplete removal of affected lung tissue none was cured and 58% were improved. In Strang's (1956) series of children 163 were treated surgically with 16 post-operative deaths. Forty-eight children were classed as cured, 56 as greatly improved, and 19 as slightly improved. Swierenga (1957) claimed "good results" (this term is not further defined) in 155 out of 176 children treated surgically with four post-operative deaths. Dyggve and Gudbjerg (1958) followed up 17 children treated by resection and found 12 symptom-free, four improved, and one dead (18

months after her last operation). Post-operative bronchography in 11 cases showed no abnormality in seven, new dilatations in three, and residual disease in one. Helm and Thompson (1958) reported the results of 174 resections in 159 patients with three post-operative deaths. In the whole series 41% were cured, 23% much improved, and 20% improved. Only 32 of their patients were children, and of these 15 were cured, six much improved, and nine improved.

When the present series is combined with the figures quoted above a total of 556 children have been treated by resection, and of these 372 (67%) are classed as "cured" or "much improved." While the operative mortality was high in the earlier reported series it has now with increasing experience fallen to a more acceptable level.

On the basis of the present study it may be claimed that resection can be carried out on 75% of bronchiectatic children with symptoms severe enough to bring them to hospital for investigation and that 70% of those submitted to operation can expect a very satisfactory result. This does not, of course, prove that these children have in fact done better with surgery than they would have done with medical management. Unfortunately, I have found no reports of a really comparable series of children who have been treated medically. Franklin (1958) reported that out of 20 children followed for between 6 and 22 years 11 were "well," three were well "despite large amounts of sputum," two were well "except for occasional attacks of pneumonia," three were "invalids," and one was dead. Wynn-Williams (1957) followed 217 patients, of whom only 15 were treated by resection: only 39 of his patients were aged 15 years or less and of these two had died, two had deteriorated, 19 were unchanged, and 16 had improved. Of Field's (1949) series 109 were treated medically and 12% recovered and 54% improved, but it is not clear what proportion of these cases had disease too extensive for surgery and what proportion were cases with symptoms too mild to justify surgery. In the present series it will be seen from Table VII that of the 18 operable cases treated medically only six reached the "recovered" or "much improved" categories.

Only in Franklin's (1958) small series have the results of medical treatment approached those of operation, and until more convincing evidence based on a larger series of cases is forthcoming I believe that in children resection is still the treatment of choice in all cases in which complete removal of affected lung is possible except for those with very localized dilatation of slight degree and only minor symptoms. I believe also that palliative resection should be considered in all cases with disease too extensive for complete extirpation if there are localized areas of gross cystic disease or complete lobar atelectasis. Such areas are in any case functionless, and if they can be removed without excessive risk the benefit is often striking even though complete cure cannot be expected. If these criteria are adopted I believe that the "surgical harvest" will be by no means "small" and that the crop will be of satisfactory quality.

Finally, I would emphasize that these conclusions apply only to bronchiectasis discovered during childhood. In the adult who develops bronchiectasis or who has passed through childhood and adolescence with little trouble from his bronchiectasis the situation may well be different.

Summary

Study has been made of 116 consecutive cases of irreversible bronchiectasis diagnosed in the Royal Aberdeen Hospital for Sick Children during the years 1946-55. The average annual incidence of discovered bronchiectasis in this region was 1.06 per 10,000 child population.

The history suggested that in almost 50% of the cases bronchiectasis had developed before the third birthday but that in only 25% of all cases was the child referred for investigation within one year of the onset of persistent symptoms. In nearly two-thirds of the cases the symptoms dated from an attack of pneumonia, pertussis, or measles.

A history of purulent expectoration was common but haemoptysis was rare. The presence of persistent localized rales was noted in almost every case. Other abnormal physical signs in the chest were present only in a minority of cases and finger-clubbing was seen only in one-third. Antral lavage was performed in 87 children and produced purulent secretion in 77. In 16 cases the original x-ray films were passed as normal or as showing only a doubtful increase of basal markings. Bronchography showed bilateral disease in 61 cases and involvement of more than two lobes in 57.

Bronchography was repeated in 79 children after an interval varying from six months to eight years and without operation intervening. In 15 cases bronchiectasis was demonstrated in 17 segments previously considered normal and in seven cases eight previously affected segments now appeared normal. The possibility of spread must be remembered when surgery is contemplated, and the interval between bronchography and operation should not be too long.

Three children have died as a result of their bronchiectasis and three are untraced. Surgical treatment has been completed in 75 cases, five have had unilateral resection and are awaiting operation on the other side, two are awaiting operation, and 28 have been treated medically.

A total of 104 resections have been carried out on 80 children, with two post-operative deaths. Post-operative atelectasis was common, particularly in younger children, and lobar or segmental resection is now usually deferred to later childhood or adolescence. At final follow-up examination, between 6 months and 12 years after operation, 55% were considered to have recovered completely and a further 16% had minimal symptoms only.

The cases treated medically were not a homogeneous group and cannot be directly compared with the surgical group. In 18 of these cases complete resection of the affected lung would have been technically possible: at follow-up examination six of them were classed as "recovered" or "much improved."

These results are compared with previous reports of follow-up studies of childhood bronchiectasis, and it is suggested that lung resection offers the best chance of a satisfactory result in a large proportion of children with bronchiectasis.

I wish to thank Mr. F. J. Sambrook Gowar and Mr. P. L. Brunnen for carrying out all the surgical treatment in these cases and for giving me access to their records. I am also indebted to Dr. P. Macarthur, Inverness, who arranged for the follow-up examination of the cases from the Northern Hospital Region, to numerous family doctors who sent me

reports on the present state of patients whom I was unable to examine myself, and to Professor John Craig and Mr. F. J. Sambrook Gowar for advice and criticism on the preparation of the paper.

ADDENDUM.—Since this paper was completed Field (1961) has presented a further report on the cases described in her original papers. After a follow-up period of from 8 to 21 years 35 out of 104 patients treated medically and 36 out of 121 treated surgically were classed as "well." In the surgical series a further 79 were improved, but in the medical series only 42. The two groups were not considered comparable, and the medical group included 38 "mild cases, some of which were of doubtful permanency."

REFERENCES

- Allibone, E. C., Allison, P. R., and Zinnemann, K. (1956). *Brit. med. J.*, **1**, 1457.
 Brenneman, J. (1959). *Practice of Pediatrics*, vol. 2, ch. 54, p. 48. Prior, Hagerstown.
 Churchill, E. D., and Belsey, R. (1939). *Ann. Surg.*, **109**, 481.
 Clark, N. S., Nairn, R. C., and Sambrook Gowar, F. J. (1956). *Arch. Dis. Childh.*, **31**, 358.
 Diamond, S., and Van Loon, E. L. (1942). *J. Amer. med. Ass.*, **118**, 771.
 Dyggve, H., and Gudbjerg, C. E. (1958). *Acta pediat. (Uppsala)*, **47**, 193.
 Ellis, R. W. B. (1956). *Disease in Infancy and Childhood*, 2nd ed., p. 558. Livingstone, Edinburgh.
 Field, C. E. (1949). *Pediatrics*, **4**, 21, 231, 355.
 — (1961). *Arch. Dis. Childh.*, **36**, 587.
 Franklin, A. W. (1944). *Proc. roy. Soc. Med.*, **37**, 576.
 — (1958). *Arch. Dis. Childh.*, **33**, 19.
 Ginsberg, R. L., Cooley, J. C., Olsen, A. M., Kirklin, J. W., and Clagget, O. T. (1955). *Surg. Gynec. Obstet.*, **101**, 99.
 Graham, S., and Hutchison, J. H. (1947). *Arch. Dis. Childh.*, **22**, 162.
 Grulee, C. G., and Eley, R. C. (1948). *The Child in Health and Disease*, p. 246. Williams and Wilkins, Baltimore.
 Helm, W. H., and Thompson, V. C. (1958). *Quart. J. Med.*, **27**, 353.
 Kent, E. M. (1942). *Amer. Rev. Tuberc.*, **46**, 524.
 — (1955). *Lancet*, **1955**, **2**, 489.
 — (1958a), **1**, 631.
 — (1958b), **2**, 734.
 Nelson, W. E. (1959). *Textbook of Pediatrics*, 7th ed., p. 808. Saunders, Philadelphia and London.
 Paterson, D., and Moncrieff, A. (1949). *Diseases of Children*, 4th ed., vol. 1, p. 661. Arnold, London.
 Perry, K. M. A., and King, D. S. (1940). *Amer. Rev. Tuberc.*, **41**, 531.
 Strang, C. (1956). *Ann. intern. Med.*, **44**, 630.
 Swierenga, J. (1957). *Dis. Chest*, **32**, 154.
 Williams, H., and O'Reilly, R. N. (1959). *Arch. Dis. Childh.*, **34**, 192.
 Wynn-Williams, N. (1957). *Tubercle (Lond.)*, **38**, 133.

A new Radiotherapy Department at Northampton General Hospital was opened last November by Sir George Godber, Chief Medical Officer, Ministry of Health. The new building will house not only all the existing radiotherapy services but also a cobalt unit, and room will also be available for future developments, such as other supervoltage machines. The new building contains on the main floor two supervoltage treatment rooms, conventional deep x-ray therapy and superficial x-ray therapy rooms, examination and planning rooms, operating theatre for radium work, consulting suites, waiting space, and reception office. On the ground floor are fully equipped radiobiological research laboratories, physics laboratory, workshop, mould room, library, staff recreation room, and other services. The cost of the building and equipment was approximately £120,000, a great proportion of which was provided by a donation by the late Mrs. Alice Barratt and endowments of the Barratt family. Among the audience addressed by Sir George Godber were local personalities and many representatives of radiotherapy departments in Great Britain.