this dosage. But in view of the fact that no other treatment was available, and she was making such marked improvement, the continued dosage seemed justifiable if it held out any hope of cure. The interesting aspect of the treatment was that, in spite of the high and continuous dosage, living worms were occasionally seen four months after treatment began. They gradually became fewer in number, and finally none were seen. This suggests that diethylcarbamazine acts on the *lagochilascaris* in the same way that it acts on the adult filarial worms, sterilizing them rather than killing them outright, and that treatment should be continued for the natural life of the remaining worms.

DISCUSSION

Lagochilascaris minor is a small worm, 1–1.5 cm. in length, resembling a threadworm on superficial examination. Winkel and Treurniet (1956) reported a similar case in Surinam; a few other cases have been reported, some from Trinidad. Nothing is known of the life-cycle of this worm, and according to these authors Pawan's report that the cloudy leopard (*Felis nebulosa*) is a host is unfounded. The present case is of interest in view of the fact that the patient had been known to eat raw meat, and one might speculate that this was the source of her infection. In Tobago the (Manakou) opossum is highly favoured as a food, and in this particular household it was often eaten. Very little other wild meat was eaten. However, none of the local people questioned on this matter had ever seen any meat containing worms of this type, although they are familiar with intestinal infestations. The girl had no other peculiar habits as a child, and it is difficult to see how else she could have become infected.

Direct infection seems to be unlikely, as the patient has a family, none of whom so far has produced any worms. The youngest daughter, however, does have nasal obstruction, and this is a worry to her mother and her doctor; but in spite of careful observation nothing has been seen to suggest that she is similarly infected.

J. W. DRAPER, M.R.C.S., L.R.C.P.

Tobago, West Indies.

REFERENCE

- Winkel, W. E. F., and Treurniet, A. E. (1956). *Docum. Med. geogr. trop. (Amst.)*, **8**, 23.

Abdominal Pregnancy Complicated by Acute Appendicitis

The following report describes the very rare association of advanced extrauterine pregnancy with acute appendicitis. It also underlines the considerable difficulty in diagnosis in such a case.

CASE HISTORY

A 24-year-old Chinese woman was brought to me with a history of three months' amenorrhoea, abdominal pain, and constipation. There was an obvious pregnancy corresponding roughly to the period of amenorrhoea. Blood-pressure, pulse, and temperature were normal. The cervix, which was normally soft, was somewhat small. The only unusual finding was some diffuse abdominal tenderness, especially on displacing the uterus laterally. The symptoms were regarded as probably the minor troubles of the first trimester in a primipara, and a reassuring diagnosis was given.

Three weeks later there was still some tenderness and constipation; but the fundus had risen to the umbilicus and the foetal heart was normal. After a further fortnight I was called urgently to the patient's home. She had been

taken acutely ill and had vomited twice; the pain was more severe. She was in moderate distress, with a pulse rate of 100. Rectal temperature was 101.2° F. (38.4° C.). The abdomen was not distended, the fundus was easily palpable at the level of the umbilicus, but the uterus was in stony-hard contraction and extremely tender: displacing the uterus laterally, especially to the right, was very painful. There was no cervical dilatation or bleeding, but the foetal heart could not be heard. It was explained that this was probably a case of acute appendicitis, that operation was essential, and that the foetus might be lost. After a blood count had shown 14,000 white cells with 90% polymorphonuclears, I operated without further delay and came with difficulty, through an extended McBurney incision, on a long rigid appendix about to burst. It lay in a deep pit against what appeared to be the postero-lateral aspect of the pregnant uterus, congested with clusters of purple veins, and was removed with some effort. Troublesome venous bleeding was controlled with a pack left *in situ*. The post-operative course was smooth. The pack was out in 48 hours, the wound healed uneventfully, and the patient left the clinic on the tenth day.

She was readmitted two weeks later for removal of the foetus. However, oxytocin having failed to produce either uterine contractions or cervical dilatation, and a laminaria tent proving equally unsuccessful, an attempt was made to introduce a uterine sound. This passed for only 8 cm., and it was evident that only laparotomy could solve the problem. It did so, for at operation a macerated 4-months foetus was found free in the abdomen. The placenta was inserted on the posterior parietal peritoneum of the right side of the pelvis, on the postero-lateral aspect of the (normal-sized) uterus, and on the back of the right uterine tube and broad ligament as well as on some adjacent loops of small bowel. A small flap of placenta close to the caecum had separated from its attachment, accounting for the haemorrhage at the end of the appendectomy. What had then been taken for a cluster of veins on a pregnant uterus was now seen to be the edge of the placenta viewed through the amniotic sac. Both uterine tubes were intact. It was easily possible to detach the placenta, except in a few places on the intestine, where removal was not attempted. Bleeding was readily controlled and the post-operative course uneventful.

CONCLUSIONS

This case is a reminder that abdominal pregnancy occurs from time to time, that it may unfairly mimic normal pregnancy even to the extent of simulating a contracted uterus in the presence of peritonitis, and that an abdominal placenta seen through a small laparotomy incision looks very like a pregnant uterus.

I have been unable to find any record of a similar case. The exhaustive studies of Hreshchychyn *et al.* (1961) and of Cavanagh (1958) make no mention of such a complication. It is possible, therefore, that the present report of this rare combination of acute appendicitis and abdominal pregnancy may be the first to be published.

Geneva.

L. A. BERETVAS,* B.A., M.D

REFERENCES

- Cavanagh, D. (1958). *Amer. J. Obstet. Gynec.*, **76**, 523.
Hreshchychyn, M. M., Bogen, B., and Langheran, C. H. (1961). *Ibid.*, **81**, 303.

*Formerly stationed at Phnom-Penh, Cambodia, as a private gynaecologist.

Over 190 public libraries are now providing library services for their local hospitals, says the Library Association. Altogether, 730 hospitals in the country are getting this service, and topping the list is the Kent County Library, serving 69 hospitals of one kind and another.