M. H. GOUGH *ET AL.*: LYMPHANGIOGRAPHY

FIG. 5.—Case 3. Abdominal and iliac lymph-node enlargement due to Hodgkin's disease.

FIG. 6.—Iliac glandular enlargement in a man suffering from chronic lymphatic leukaemia.

P. S. GUPTA *ET AL.*: VENO-OCCLUSIVE DISEASE OF LIVER

FIG. 1.—Case 1. Showing acute V.O.D. with centrolobular lagoons of blood. (H. and E. $\times 100$.)

FIG. 2.—Case 1. Showing acute V.O.D. with occlusion of centrolobular vein by subendothelial oedema of intima. (Reticulin stain. $\times 400$.)

FIG. 3.—Case 2. Showing acute V.O.D. with occlusion of centrolobular vein (arrow) by connective-tissue fibres and distension of sinusoids. (Reticulin stain. $\times 200$.)

in the absence of clinical adenopathy: it is in these patients that endolymphatic radiotherapy may be most effective in destroying early malignant emboli in the sinuses of the node. Iodine has a longer half-life than gold and may prove to be a more satisfactory isotope.

With intralymphatic ^{198}Au , doses for the leg have been from 50 to 70 mc., and in the arm from 30 to 40 mc. These dosages are infused in a volume of approximately 30 mc./ml. at a rate of about 0.4 ml./minute. More rapid infusion should be avoided, as any extravasation through the lymphatic will produce a high local concentration with risk of later radionecrosis. During the next few days there may be a brisk local reaction at the gland sites—for example, inguinal region—which subsides after a period of 6 to 10 days to leave the glands somewhat enlarged and indurated. The results in a small group of patients followed up for periods extending to three years have been encouraging.

There have been no published reports of the clinical use of endolymphatic cytotoxic agents in the treatment of lymphomas or secondary deposits. The possibility that these agents, some of which are rapidly hydrolysed on contact with tissues, might produce an acute inflammatory reaction and later lymphatic obstruction will have to be carefully considered.

Summary

Clinical lymphography using water-soluble dyes was described in 1951. Following the recent development of oily contrast media which are slower and more difficult to inject, a new operative technique was required.

Oily media are retained in the lymph nodes for longer periods of time than water media. This enables a fuller study of both the channels and the nodes themselves to be made. Animal experiments and clinical trials have shown that in the recommended dosages no harmful effects arise from these new media.

Experiences with oily media are discussed in relation to the diagnosis of oedema of obscure origin and in the determination of the presence or degree of spread of lymphomas.

Lymphography may also be used to detect the presence of secondary deposits in regional glands in conditions such as epithelioma or melanoma.

The use of endolymphatic ^{198}Au is described as a method of delivering high local doses of radiation to such regional glands. Preliminary results of a clinical trial are encouraging.

We are grateful to Mr. D. Ruff for technical assistance, to the Photographic Department of St. Thomas's Hospital for the illustrations; to Messrs. Bengué & Co. Ltd. for supplies of ultrafluid lipiodol, and for a grant towards our expenses; and to Mr. L. Crook, United Kingdom Atomic Research Authority, Wantage, Berks. for help in sterilizing lymphatic cannulae by gamma-radiation.

REFERENCES

- Clementz, B., and Olin, T. (1961). *Acta radiol. (Stockh.)*, **55**, 109.
 Colin, R. (1961). *L'Exploration du Système lymphatique en Pathologie Chirurgicale*. Union Française d'Impression, Bordeaux.
 Fisher, H. W., Laurence, M. S., and Zimmerman, G. R. (1961). *J. Amer. med. Ass.*, **175**, 327.
 Hudack, S. S., and McMaster, P. D. (1933). *J. exp. Med.*, **57**, 751.
 Jantet, G. H. (1958). Ninth Scientific Meeting, Surgical Research Society, London.
 — (1962). *Brit. J. Radiol.*, **35**, 692.
 Kinmonth, J. B. (1951). Communication to Medical Research Society, London.
 — (1952). *Clin. Sci.*, **11**, 13.
 — (1954). *Ann. roy. Coll. Surg. Engl.*, **15**, 1901.
 — Rob, C. G., and Simeone, F. A. (1962). In *Vascular Surgery*. Arnold, London.
 — and Taylor, G. W. (1954). *Ann. Surg.*, **139**, 129.
 — Tracy, G. D., and Marsh, J. D. (1957). *Brit. J. Surg.*, **45**, 1.
 Schobinger, R. A. (1960). *Intraosseous Venography*. Grune and Stratton, New York.
 Tjernberg, B. (1962). *Acta radiol. (Stockh.)*, Suppl. No. 214.
 Wallace, S., Jackson, L., Schaffer, B., Gould, R. J., Greening, R. R., Weiss, A., and Kramer, S. (1961). *Radiology*, **76**, 179.

VENO-OCCLUSIVE DISEASE OF LIVER

BY

P. S. GUPTA, M.D. G. D. GUPTA, M.D. M. L. SHARMA, M.R.C.P., D.T.M.

Department of Medicine, Irwin Hospital, and Maulana Azad Medical College, New Delhi, India

[WITH SPECIAL PLATE]

Veno-occlusive disease of the liver (V.O.D.) has been recently recognized as a separate entity. The disease is seen predominantly in children, but it has also been reported in adults (Stuart and Bras, 1957). In the acute stage the small hepatic veins are involved and to a less degree the medium-sized vessels. The blockage is due to subendothelial oedema, as a result of which massive centrilobular congestion occurs which leads to a varying degree of destruction of liver cells (Jelliffe *et al.*, 1954; Bras and Hill, 1956).

Clinically, the acute stage is characterized by rapid development of abdominal discomfort, pain in the right hypochondrium, ascites, and an enlarged, tender liver. Some patients recover in four to six weeks, the results of liver-function tests and the histology becoming normal. A number, however, die in the acute stage from liver failure, while others pass into a subacute or chronic stage. The chronic phase is characterized by a picture of cirrhosis of the liver which is indistinguishable from that of other forms of cirrhosis (Stuart and Bras, 1957).

In the acute stage the diagnosis is usually based on the history, the characteristic clinical picture, and the liver-biopsy findings (Stuart and Bras, 1957).

V.O.D. has a world-wide distribution in both man and animals (Hill, 1960), and in human beings the disease occurs in Jamaica (Jelliffe *et al.*, 1954), Jerusalem, Egypt (Hashem, 1939), and Johannesburg (Higginson *et al.*, 1957). It is possible that many of the reported cases of "idiopathic" Budd-Chiari syndrome are examples of V.O.D. (Sherlock, 1958). Selzer and Parker (1951) have reported a series of cases from South Africa which began as acute Budd-Chiari syndrome and were due to eating bread made from inadequately winnowed wheat contaminated by a species of weed, *Senecio*. There is an increasing weight of evidence that a number of bush-teas taken for medicinal and other purposes in Jamaica are responsible for V.O.D. (Bras *et al.*, 1957). Tirumurti and Radhakrishna Rao (1934) hold the view that the childhood cirrhosis seen in India may be of similar aetiology, but no case of V.O.D. in adults has

so far been reported from that country. However, that it does occur in adults in India is shown by the two cases described below, which followed the use of herbal drugs.

Case 1

A Hindu male aged 25 years was admitted to Irwin Hospital, New Delhi, on February 11, 1960, with a diagnosis of acute abdomen. The history was that two months before admission he started taking some herbal pills obtained from a "hakim" as a remedy for ichthyosis. Within a few days he lost his appetite, became disinclined to work, and developed swelling of his elbow-joints and a generalized itching rash. With treatment the rash and joint pains subsided, but loss of appetite and weakness persisted. He continued taking the pills for another six weeks, when he experienced sudden epigastric pain and abdominal distension which increased in severity till the day of his admission to hospital. The bowels were sluggish, but he had no urinary symptoms. There was no history of a similar illness and he had never been jaundiced. He took a vegetarian, fairly nutritious diet. Personal and family histories were non-contributory.

The patient, who was fairly well built, looked ill and anxious. There was no jaundice, no rise in jugular venous pressure, no lymph-node enlargement, and no oedema. Pulse rate 96 per minute; temperature 99.4° F. (37.4° C.); blood-pressure 130/80 mm. Hg. The abdomen was distended, with no venous prominence over the surface. There was vague tenderness all over the abdomen, more marked in the right hypochondrium. The liver was felt, by dipping, three fingerbreadths below the right costal margin and was smooth, firm, and tender. The hepato-jugular reflux was absent. The spleen was not palpable, and marked ascites was present. The other systems were normal.

Investigations

Haemoglobin 13 g. per 100 ml.; R.B.C. 4.1 millions per c.mm.; total W.B.C. 12,100 per c.mm. (polymorphs 76%, lymphocytes 22%, and monocytes 2%); E.S.R. 4 mm. in the first hour (Westergren). Urine and stool normal. Ascitic fluid: straw-coloured; specific gravity 1016; protein concentration 2.88 g. per 100 ml.; microscopical examination revealed 102 cells per c.mm., consisting of erythrocytes, lymphocytes, and endothelial cells. Total serum proteins 5.4 g. per 100 ml., with albumin 2.8 g. and globulin 2.6 g. (A/G ratio 1:1); serum bilirubin 0.6 mg. per 100 ml.; alkaline phosphatase 8 K.A. units; zinc sulphate turbidity 4 units; blood urea 44 mg. per 100 ml.; true blood glucose 78 mg. per 100 ml.; serum cholesterol 150 mg. per 100 ml. Blood V.D.R.L., P.P.R., and Meinicke tests negative. Chest radiograph showed minimal pleural effusion on the right side; fluoroscopy of chest for cardiac enlargement and pulsations, normal. There was no evidence of oesophageal varices. Plain radiograph of abdomen normal. Electrocardiogram within normal limits. Venous pressure in arms 10.8 cm. of water, and in legs 9.8 cm. Liver biopsy revealed venous congestion with centrilobular distension of sinusoids giving an appearance of lagoons of blood resembling haemorrhage and cell necrosis (Special Plate, Fig. 1). Connective-tissue stains showed occlusions of centrilobular veins by sub-endothelial oedema of the intima (Special Plate, Fig. 2) or a fine reticulum.

Treatment and Progress

He was given injections of streptomycin with penicillin, chlorothiazide, and a salt-free diet. Ascitic fluid was tapped twice to relieve the abdominal discomfort. The general condition improved and the ascites diminished. The liver, however, remained enlarged, and the patient was discharged from hospital. On follow-up examination two months later the ascites had disappeared. The liver was enlarged two fingerbreadths below the costal margin and was smooth and firm in consistency. Eighteen months later the patient was well, though the liver was palpable one fingerbreadth below the costal margin, and was smooth and firm. The histological appearance of the liver on biopsy was normal except for some focal scarring.

Case 2

A Hindu farmer aged 35 was admitted on December 15, 1960, complaining of progressive distension of the abdomen for 10 days and some difficulty in breathing and abdominal discomfort for a week. He had had generalized relapsing psoriasis for about 12 years. Three months before the present episode he started taking treatment from a "hakim" for the psoriasis in the form of a decoction made from fresh weeds. This he took for about two months, at the end of which the skin condition settled down, but he suddenly lost all appetite. With additional treatment the appetite returned within a week, but he suffered moderate aching pain in the right upper abdomen, mild generalized abdominal discomfort, and sluggish bowels. The abdomen began to swell, and in 10 days became so large as to lead to respiratory embarrassment. At no time was there jaundice, highly coloured urine, change in stool colour, fever, or definite oliguria. There was nothing significant in the past or family history. The patient took a good vegetarian diet; he did not smoke or take alcohol.

On examination he was seen to be of average build, but with a greatly distended abdomen. Pulse rate 80 per minute; blood-pressure 120/80 mm. Hg.; temperature 99° F. (37.2° C.). Positive findings were: moderately tense ascites, tenderness in the right upper quadrant, doubtful hepatomegaly on dipping, and receding generalized extensor psoriasis. There was no oedema of the face, sacrum, or feet, no jaundice or lymphadenopathy, and no rise in jugular venous pressure or venous prominence elsewhere.

Investigations

Haemoglobin 10 g. per 100 ml.; R.B.C. 4 millions per c.mm.; total W.B.C. 5,600 per c.mm. (polymorphs 70%, lymphocytes 28%, monocytes 2%); E.S.R., 20 mm. in first hour (Westergren). Urine: no sugar, albumin, bile pigments or salts, or urobilinogen detected; sediment normal on microscopical examination. Stool normal. Ascitic fluid: clear, light-yellow coloured, with protein 1.08 g. per 100 ml., sp. gr. 1012; microscopical examination revealed 24 cells (10 lymphocytes, 4 eosinophils, and 10 red cells) per c.mm. Total plasma proteins 6.4 g. per 100 ml., with albumin 3.5 g. and globulin 2.9 g. (A/G ratio 1.2:1) zinc sulphate turbidity 4 units; alkaline phosphatase 12 K.A. units; serum sodium 136 and potassium 4.5 m.Eq/l.; CO₂-combining-power 55 ml.%; plasma chlorides (as NaCl) 650 mg. per 100 ml.; serum cholesterol 250 mg. per 100 ml.; serum bilirubin 0.4 mg. per 100 ml.; blood urea 32 mg. per 100 ml.; prothrombin time (Quick's one-stage) 24 seconds. Radiograph of chest normal. Barium swallow did not reveal oesophageal varices. Fluoroscopy of heart did not reveal any abnormality.

Venous pressure in the antecubital vein was 12 cm. of normal saline, no difference occurring on hepatic pressure. In the femoral vein (after abdominal paracentesis) the pressure was 15 cm. of normal saline.

Liver biopsy 9 days after admission showed appearances similar to Case 1 with centrilobular lagoons of blood, due to sinusoidal dilatation following occlusion of the central veins by a connective-tissue reticulum (Special Plate, Fig. 3) or intimal oedema.

Progress and Treatment

He was given a low-salt, high-protein (100 g.) diet, with mersalyl 2 ml. twice weekly and multivitamin tablets. As there was no diuretic response at the end of a week, abdominal paracentesis was performed, 130 oz. (3.7 l.) of fluid being removed. At this stage the liver was palpable three fingerbreadths below the costal margin; it was firm, smooth, and tender. The spleen was not palpable. Ascites was rapidly forming in spite of the addition of chlorothiazide to the treatment, and abdominal paracentesis was repeated. Symptomatic treatment for abdominal pain and dyspepsia was given.

Liver biopsy repeated after one month showed marked improvement, there being few central haemorrhages. Liver cells appeared normal, but central veins and surrounding sinusoids were still dilated. Clinically, the liver was still two fingerbreadths enlarged, and firm though less tender.

One month after admission the ascites gradually disappeared, and did not re-form when diuretics were withdrawn. After another month the liver was only one fingerbreadth enlarged; it was not tender but was slightly firmer. The spleen was not palpable. The patient felt very well. The total duration of the illness had been three months. On follow-up eight months from the start of the illness findings were similar and the patient felt well.

Discussion

The history and the clinical picture of these two cases were very similar, and were fairly characteristic of an acute, severe, passive venous congestion of the liver confirmed by liver biopsy. In the absence of evidence of cardiac failure and of occlusion of the inferior vena cava the lesion responsible for the liver pathology appears to be thrombosis of one or more hepatic veins, or of the smaller tributaries at any level. The former is known to be the essential pathological basis of Budd-Chiari syndrome, and the latter of the "veno-occlusive disease of the liver" reported from Jamaica and other places (Jelliffe *et al.*, 1954; Bras and Hill, 1956). In Budd-Chiari syndrome the obstruction in the main hepatic vein or veins occurs mostly secondarily to other disease processes such as pressure and invasion from adjoining hepatic neoplasms, vena-caval spread of renal carcinoma, thrombophlebitis migrans, portal pyaemia (Chiari, 1899), and polycythaemic states (Sohval, 1938). In many cases, however, no such cause or association can be found, and Chiari (1899) favoured primary inflammation of one or more hepatic veins, leading to thrombosis, as a cause for such cases.

The entity described as veno-occlusive disease of the liver from Jamaica and other places differs from Budd-Chiari syndrome in that the cause is ingestion of toxic weeds and other substances, and the essential pathology is an obliterative endophlebitis of the small tributaries of the hepatic venous system (Stuart and Bras, 1957).

The two cases described here did not, either at the time of presentation or subsequently during follow-up at 1½ years and 8 months respectively, show any evidence of such disease states as are known to produce Budd-Chiari syndrome. Moreover, the acute onset and the progressive improvement are suggestive of some noxious process which operated acutely for only a certain length of time.

The history of the use of indigenous medicines in both cases immediately preceding the hepatic disease, in the absence of any other recognizable cause for the latter, leads to the suspicion that some ingredient of the herbal medicine is possibly the aetiological agent. Obviously the skin conditions present—ichthyosis in one case and psoriasis in the other—could not have any relation to the liver disease. The mechanism of action could be similar to that of the plants *Crotalaria* and *Senecio*, known to produce the hepatic disease in Jamaica and other countries (Selzer and Parker, 1951; Bras *et al.*, 1957) and also in the experimental animal (Berry and Bras, 1957; Hill and Martin, 1958; Hill *et al.*, 1958; Bruun-Hanssen, 1961). With this tentative presumption we have labelled our cases "veno-occlusive disease of the liver." Confirmation of such a presumption will be possible only by experimentally reproducing the disease

with these plants, and we are now working along these lines.

The importance of recognizing the basic pathology in these cases lies in the fact that, though in the early stages the disease may be diagnosed merely as cirrhosis, there is a risk that eventually it will pass into the stage of chronic veno-occlusive disease. It is possible that many of the cases of so-called "portal cirrhosis" that occur in India, at any rate, have this aetiological and pathogenetic background, as indigenous treatment with herbal medicines is not uncommon and many patients suffering from "portal cirrhosis" give a history of abdominal pain and ascites months or years before from which they had recovered (unpublished personal observation).

Our purpose in reporting these two cases is to confirm that this type of "iatrogenic" liver disease does exist in India, and to suggest that its occurrence should be borne in mind.

Summary

The clinical features and incidence of "veno-occlusive disease" of the liver, which has recently been recognized as a distinct entity, are briefly described. Two cases which occurred after the indigenous treatment with herbal medicines of ichthyosis and psoriasis respectively are reported. The patients presented with the picture of acute "veno-occlusive disease" of the liver, and almost completely recovered within three months. One and a half years and eight months respectively after the start of the illness they were found to be well and symptomless, though their livers continue to be palpable and somewhat firm. The significance of encountering this disease in India and its possible relation to cirrhosis of the liver are discussed.

Our thanks are due to Dr. D. N. Gupta, Professor of Pathology, Maulana Azad Medical College, New Delhi, for his study of the liver tissue; and to Colonel B. L. Taneja, Medical Superintendent, Irwin Hospital, and Principal, Maulana Azad Medical College, for permission to report these cases.

REFERENCES

- Berry, D. M., and Bras, G. (1957). *N. Amer. Vet.*, **38**, 323.
 Bras, G., Berry, D. M., and Gorgy, P. (1957). *Lancet*, **1**, 960.
 — and Hill, K. R. (1956). *Ibid.*, **2**, 161.
 Bruun-Hanssen, C. F. (1961). *Acta path. microbiol. scand.*, **53**, 329.
 Chiari, H. (1899). Quoted by Sherlock, S. (1958).
 Hashem, M. (1939). *J. Egypt med. Ass.*, **22**, 319.
 Higginson, J., Grobbelaar, B. C., and Walker, A. R. (1957). *Amer. J. Path.*, **33**, 29.
 Hill, K. R. (1960). *Proc. roy. Soc. Med.*, **53**, 281.
 — Stephenson, C. F., and Filshie, I. (1958). *Lancet*, **1**, 623.
 Jelliffe, D. B., Bras, G., and Stuart, K. L. (1954). *Ann. trop. Med. Parasit.*, **48**, 386.
 Selzer, G., and Parker, R. G. F. (1951). *Amer. J. Path.*, **27**, 885.
 Sherlock, S. (1958). *Diseases of the Liver and Biliary System*, 2nd ed. Blackwell, Oxford.
 Sohval, A. R. (1938). *Arch. intern. Med.*, **62**, 925.
 Stuart, K. L., and Bras, G. (1957). *Quart. J. Med.*, **26**, 291.
 Tirumurti, T. S., and Radhakrishna Rao, M. V. (1934). *Indian J. Pediat.*, **1**, 153.

The 1963 List of Products approved under the Agricultural Chemicals Approval Scheme is now available free of charge. This list is the most comprehensive yet published and contains details of almost 600 products based on more than 100 chemicals or mixtures of chemicals used for pest, disease, and weed control. The introductory pages of the list give hints on the safe handling of chemicals and names those which require special precautions in use. The principal pests, diseases, and weeds controlled and the crops which may be treated are given for each different chemical. Copies can be obtained free from the Ministry of Agriculture, Fisheries, and Food (Publications), Ruskin Avenue, Kew, Surrey, or any of the Ministry's regional and divisional offices.