*Papers and Originals***NEOPLASTIC TRANSFORMATION BY POLYOMA VIRUS
AND ITS WIDER IMPLICATIONS***

BY

M. G. P. STOKER, M.D., F.R.S.Ed.
Professor of Virology, University of Glasgow

[WITH SPECIAL PLATE]

Infection with a virus is one of the few ways of introducing exogenous genetic material into a cell, and viruses ought therefore to be useful for the investigation of heredity and genetic expression at the cell level. This is possible, however, only if the cell survives the infection at least long enough for identification of any genetic change. When this occurs, as with lysogenic cultures of bacteria, the scientific rewards have been considerable.

Unfortunately animal cells are immediately killed by almost all known viruses which infect them, and this prevents the study of genetic changes which may result.

The most promising exceptions are the viruses which produce tumours, and the pioneer work of Rubin (1955) and Temin and Rubin (1958) with Rous sarcoma virus of chickens has shown how such studies can be made using cell cultures rather than the intact bird. Recently a whole group of new viruses have been isolated which will induce tumours in small mammals, and one of these, polyoma virus, turns out to be particularly suitable for experimental study of the neoplastic process. I propose to review current knowledge about malignant transformation caused by polyoma virus and to consider some implications without putting forward any general theories about the aetiology of cancer.

Polyoma virus was first isolated in Washington by Stewart *et al.* (1957) from the tissues of leukaemic mice. They reported a filterable agent which, after propagation in tissue culture, caused parotid and several other tumours after inoculation of newborn mice. In retrospect it seems likely that this same agent was responsible for the parotid tumours previously reported by Gross (1953) during studies of his leukaemic agent.

Natural Distribution

Polyoma infection has been found in mice in the U.S.A., Canada (Axelrad *et al.*, 1960), and Britain (Negroni *et al.*, 1959), and following the work of Rowe *et al.* (1959) it is now clear that the virus is a common commensal organism in stocks of laboratory mice. Some mouse colonies are free of infection, but in infected colonies a high proportion of adult mice are carriers. They excrete the virus in the saliva and urine, which in turn infects the young during the early months of life. The virus is not transmitted *in utero* or via the milk, however. Infection is followed by appearance of specific antibodies which provide an easy method of detecting the presence of the virus.

Rowe *et al.* (1958) have extended their studies to wild mice (*Mus musculus*) and have found polyoma infection in mice captured in both urban and rural areas. Polyoma virus has been found in abundance in farms and grain-stores contaminated by the excreta of wild mice, and this may be a source of infection for labora-

tory mice through the processed grain foods (Huebner *et al.*, 1961). There is no evidence of natural infection of man or any other animal.

Mice which harbour polyoma virus show no obvious signs of disease, and it is not known if the naturally acquired infection can give rise to tumours. From data obtained by ringing and recapturing mice, however, it appears that *Mus musculus* rarely survives longer than six months, while the incubation period before tumours appear may be very long, especially with low doses of virus and routes of infection associated with natural transmission. Moreover, maternal antibody may prevent infection in the early days of life when susceptibility to experimental tumour production is highest.

The Virus

Polyoma virus grows well in cultured mouse embryo cells, yielding virus stocks of high titre, which may be used for purification and characterization of the virus particle. Such virus stocks contain large numbers of spherical particles 450 A.U. in diameter (Special Plate, Fig. 1). The surface of each particle is composed of subunits arranged so as to show icosahedral symmetry, but subunit arrangements giving long filaments have also been found (Wildy *et al.*, 1960; Howatson and Almeida, 1960).

It is assumed that these subunits are protein; in addition, the virus contains 6 to 8 million molecular weight units of deoxyribose nucleic acid (D.N.A.) (Smith *et al.*, 1960; Crawford, 1963), presumably located in the centre of the particle. There is no evidence of any other constituents. The polyoma virus is similar in size and structure to three other tumour viruses of mammals—rabbit papilloma virus (also known to contain D.N.A.) (Watson and Littlefield, 1960), the virus found in human warts (Williams *et al.*, 1961), and the vacuolating agent SV40 from monkey kidney. The chicken-tumour viruses, such as Rous sarcoma virus, contain ribonucleic acid (R.N.A.) (Crawford and Crawford, 1961) and are quite different in structure.

Polyoma virus has two valuable attributes which allow detection and quantitative estimation of the virus and thus assist experimental investigation. First, infection of mouse embryo cells results in cell death with a consequent cytopathic effect. This allows counts of infective particles by plaque assay or by end-point titration. Secondly, the virus adsorbs to red blood cells, causing visible agglutination in the same way as influenza virus (Eddy *et al.*, 1958a). This provides an estimate of total physical particles without resort to electronmicroscope counts. It also gives a test for polyoma antibody, which will prevent the viral haemagglutination. So far all strains have shown antigenic similarity.

The virus has a very high thermal resistance (Eddy *et al.*, 1958b), and the consequent low rate of spontaneous

*Based on a lecture delivered to the Royal Society of Edinburgh on February 5, 1962.

inactivation no doubt contributes to the ease of natural spread of the virus among mice.

Tumour Induction in Animals

Polyoma virus will induce tumours in mice (Stewart *et al.*, 1957), hamsters (Eddy *et al.*, 1958c), rats (Eddy *et al.*, 1959b), rabbits (Eddy *et al.*, 1959a), and ferrets (Harris *et al.*, 1961). The action of the virus is most pronounced when the animals are inoculated within the first two days after birth, but tumours have also been induced with a longer incubation in hamsters inoculated after weaning at three weeks (Stewart *et al.*, 1958).

Table I shows examples of the remarkable variety of different tumours which appear as a result of inoculation with polyoma virus. The virus also causes non-malignant proliferation of renal epithelium in the mouse and rapid necrosis of renal interstitial cells in the

TABLE I.—*Different Tumours Induced by Polyoma Virus*

Mouse (Stewart <i>et al.</i> , 1957)
Pleomorphic carcinoma (parotid)
Adenocarcinoma (breast)
Epidermoid carcinoma (skin)
Spindle-cell sarcoma (various sites)
Haemangioma (liver)
Hamster (Eddy <i>et al.</i> , 1958c)
Spindle-cell sarcoma (kidney and various other sites)
Haemangioma (liver)
Rat (Eddy <i>et al.</i> , 1959b)
Spindle-cell sarcoma (kidney and various other sites)
Rabbit (Eddy <i>et al.</i> , 1959a)
Fibroma (at inoculation site)
Ferret (Harris <i>et al.</i> , 1961)
Fibrosarcoma (at inoculation site)

hamster (Ham *et al.*, 1960). It thus appears that the virus can act on many different target cells in various animals and cause malignant growth, simple proliferation, or degeneration. This could be due to a mixed population of virus particles, each with a specific effect on a certain type of cell, or alternatively each infectious virus particle may be competent to produce the whole range of effects. Virus purified by repeated plaque isolation should be genetically pure, and it has been reported that such plaque-purified virus does induce a variety of effects (Stoker *et al.*, 1960). Unfortunately, there are technical difficulties in isolating pure lines of polyoma virus by plaque isolation, and it is doubtful if we can yet be quite sure that pure clones of virus are multipotential.

Multiple tumours generally occur in each animal inoculated with a large dose of virus, and there is evidence to suggest that, in the hamster at least, this is due to multicentric action of the original virus inoculation rather than to secondary spread of tumour cells (Stoker, 1960). In other ways the tumours of mouse and hamster have the usual criteria of malignancy, and they can be transplanted to other animals. Transplanted tumours usually contain virus, but after several serial transplants virus may no longer be detectable, though the behaviour of the tumour is unaltered. The role of virus in the continuation of the tumour is discussed below.

Tumours take three or more months to develop after inoculation of newborn mice (Stewart *et al.*, 1957), but the same virus dose will kill hamsters with renal sarcoma in two weeks or less (Stoker, 1960; Ham *et al.*, 1961). Examination of the kidneys as early as four days after inoculation will show tumour cells in the intertubular tissue (Stoker, 1960; Ham *et al.*, 1961). Under certain conditions the number of foci is directly related to virus dose, so it is suggested that each is a colony resulting from a single initial virus-cell contact. The number of cells in each focus suggests that the primary

change occurs within a day of inoculation of the virus (Stoker, 1960).

From virus counts it is estimated that one tumour-producing dose is given with an inoculation of 70 plaque-forming units of virus into a hamster, but reduction in virus causes an increasing delay in the appearance of tumours, and this limits the sensitivity of the estimate (Stoker, 1960).

There is a great deal to be learned from studies on the tumour-producing activity of polyoma virus in animals, but for investigation of the mechanism of virus action in the initial change from a normal to a neoplastic cell the intact animal has obvious drawbacks. Fortunately, the effects of the virus may be reproduced and examined under controlled conditions in cultured cells *in vitro*, thus removing many of the variables of the *in vivo* experiments and allowing deductions about individual cells.

Behaviour of Polyoma Virus in Tissue Culture

When polyoma virus is added to mouse-embryo fibroblasts the virus grows, with accompanying cell degeneration, and several thousand virus particles are formed per cell. The site of growth of the virus is the nucleus, where electronmicrographs show masses of particles sometimes condensed into small crystals (Howatson *et al.*, 1960). About 20% of cells survive, however, and these may be grown and propagated. Hamster fibroblasts exposed to virus, on the other hand, show no gross cell damage, and little or no new virus is produced.

Vogt and Dulbecco (1960) made the important observation that cultures grown from hamster cells or surviving mouse cells, after exposure to large doses of virus, were recognizably different and showed characteristics of tumour cells in culture. The new or "transformed" character of the cells persisted indefinitely on subculture. Similar changes were observed by Sachs and Medina (1961), and also in our own laboratory (Stoker and Macpherson, 1961).

It thus appears that the virus has two distinct effects on fibroblastic cells in culture: (1) cytotoxic, with virus growth; and (2) transforming, resulting in viable but genetically altered cells.

Transformation *in vitro* by Polyoma Virus

Transformation was originally observed in mass cultures of cells some weeks after exposure to virus, and it was not possible to draw conclusions about the initial events. Various improvements have now been made in the experimental system which allow more precise analysis. Thus estimates of the number of cells transformed by a single short exposure to virus can be made by allowing separated cells to grow into isolated colonies which can be identified as normal or transformed (Stoker and Macpherson, 1961).

A continuous line of fast-growing hamster fibroblasts has also been isolated; this can be propagated continuously, stored at low temperatures, and recovered with ease, and it can be used as a source of pure lines or clones derived from single cells. This line of cells and its subclones are all transformable by polyoma virus (Macpherson and Stoker, 1962).

The experimental system (Stoker and Abel, 1962) at present used in our laboratory is shown in Fig. A. A clone is derived from a single cell, and samples of this population are exposed to virus to allow virus

attachment. After one hour the mixture is diluted to prevent further attachment, and a thousand or more separated cells are added to each Petri dish. A proportion of cells grow to form colonies, which after one week can be identified as normal or transformed and counted. Individual colonies can be removed if required and grown into larger populations.

FIG. A.—Experimental system for assay of neoplastic transformation by polyoma virus.

Characteristics of Transformed Cells

Comparison with Normal Cells from Same Clonal Population

Studies of the characteristics of neoplastic cells have in the past been hindered by the lack of a satisfactory control population of cells for comparison. The populations which have been compared have almost invariably been mixed, and differences found could always be due to secondary changes in the proportions of a particular cell type. Ideally, the normal and neoplastic cells should be like litter mates—that is, genetically pure lines derived from the same single parent cell. This is now possible by comparing polyoma-transformed hamster cells with untransformed cells from the same clonal population after a limited number of generations.

Cultures of transformed cells are indistinguishable from cultures of cells from tumours induced *in vivo*. Such cells have not been seen in normal cultures and they differ markedly from normal cells (see Table II). Normal fibroblasts are usually elongated bipolar cells oriented in parallel with one another. Crossing of the cells in an uncrowded plate is rare through the effect

of contact inhibition (Abercrombie and Ambrose, 1962). Transformed or tumour cells are often triangular and they lie at random, criss-crossing one another and piling up several layers thick, presumably owing to loss of

TABLE II.—Some Characteristics of a Clone of Normal Hamster Fibroblasts, With Cells from the Same Clone After Transformation by Polyoma Virus

	Normal Hamster Fibroblasts (BHK21 Strain)	Hamster Fibroblasts (BHK21 Strain) Transformed <i>in vitro</i> by Polyoma Virus
Arrangement of cells in culture	Oriented. Little overlap (contact inhibition)	Chaotic. Much overlap. (loss of contact inhibition)
Transplantability into hamsters (subcutaneous)	Poor. Minimum $>10^6$ cells	Good. Minimum $>10^4$ cells
Glycolysis	Normal	Increased
Alkaline phosphatase content	Normal	Increased
Electrophoretic mobility (Forrester <i>et al.</i> , 1962)	Homogeneous	Heterogeneous and increased
Morphology of isolated cell	No regular difference detected	
Chromosomes	Both divide every 12 hours	
Growth rate in culture	No difference detected	
Viral susceptibility		

contact inhibition (Special Plate, Figs 2 and 3). Transformed and tumour cells also show a higher rate of glycolysis than normal cells. When freshly isolated they have the same number of chromosomes as normal hamster cells, but this characteristic is less stable in the normal cells. On continued propagation the transformed cultures contain an increasing number of tetraploid cells (Macpherson, 1963).

Normal cells can sometimes be transplanted successfully if large numbers (more than one million) are injected into hamsters, but they grow slowly. If the cells are reisolated they grow as normal fibroblasts. Transformed cells, or tumour cells, grow rapidly from a smaller number of cells (less than 10,000) into progressive sarcomata, and the recultured cells have the typical random growth already described (Macpherson, 1963).

It is a reasonable working hypothesis that the transformation arising from virus action in tissue culture is equivalent to the initial stage of tumour induction *in vivo*. We continue to refer to the *in vitro* change as neoplastic transformation, or simply transformation, rather than *in vitro* tumour induction.

Evidence for a New Antigen in Transformed Cells

Sjögren *et al.* (1961) and Habel (1961) have independently observed that immunization of mice or hamsters with live polyoma virus also protects the animals against transplantation with tumour cells. Killed virus has no effect, and it is suggested that the growth of the virus gives rise to a new antigen in infected cells which is not incorporated in the virus particle itself. Appearance of this new cell antigen would be followed by sensitization, and therefore rejection of grafts of polyoma-induced tumour cells which themselves contain the antigen. Such autoimmunization does not lead to the regression of the original tumours induced by the virus, but, unlike the grafted cells, these tumours would already be well established before the immunity developed.

It is possible that the new properties which appear in the transformed cell are due to the appearance of one or more new proteins corresponding to the antigen found in the tumour cells. In fact, the antigen has now been found in cells transformed *in vitro*.

The appearance of new antigens, consequent on virus infection but not incorporated into the particle, are reminiscent of conversion by bacteriophage leading to

new somatic antigens in certain salmonellae (Iseki and Sakai, 1953), or toxin production by *Corynebacterium diphtheriae* (Groman, 1953). We will now consider the relationship between polyoma-transformed cells and lysogenic bacteria.

Relationship to Bacterial Lysogeny

Transformed cultures, or cultures of tumour cells, though apparently showing no cytopathic degeneration, may continue to yield a constant low level of virus (Winocour and Sachs, 1961). This is released discontinuously as bursts from single cells, and it was at first believed that it might be due to spontaneous induction occurring at a low rate as in lysogenic bacteria-carrying prophage. In lysogenic cultures of bacteria every cell has the potentiality to yield fully infective virus, either spontaneously or after induction by various agents such as ultra-violet light; moreover, the cells are immune to superinfection with related viruses.

On further examination by Vogt and Dulbecco (1962) the analogy has broken down for the following reasons. (1) When clones are derived from single cells they retain all the characteristics of transformed cells, but infective virus particles can no longer be detected. (2) Such cultures can be reinforced with polyoma virus, again giving a persistent infection with constant low yield, which can in turn be cured by single-cell isolation. (3) Attempts to induce virus production and release with substances active on lysogenic bacteria have completely failed.

It has been concluded by Vogt and Dulbecco that the persistence of virus in populations of neoplastic cells is coincidental, and due to a cytotoxic infection spreading at a slow rate in a partially resistant population of cells, resembling the carrier cultures of poliovirus, herpes virus, etc. This leads to the important conclusion that infectious virus or potentiality to make it is not essential to maintenance of the neoplastic state.

Proportion of Cells Affected

When cells are exposed to concentrations of 1,000 infective units per cell—that is, equivalent to about 70,000 particles per cell—about 3% of cells are transformed (Stoker and Abel, 1962). With less virus the number of transformed cells is proportionately reduced, but with more virus the number is not increased. This low efficiency of transformation is not due to deficient virus uptake, because we know that the virus adsorbs and enters the cytoplasm of all the cells (Fraser and Gharpuré, 1962), so the limitation probably occurs at a subsequent stage. A large proportion of cells might be genetically resistant to transformation, but this was ruled out by showing that several clones representing single cells in the original population all had the same low rate of transformation (Macpherson and Stoker, 1962). There still may be physiological differences, however, such as stage of mitosis, affecting transformability in a genetically uniform population of cells.

Induction, Not Selection

It has been suggested that carcinogenesis involves creation of suitable conditions for selection of pre-existing neoplastic cells, which are always present in small numbers. Alison (1961) has pointed out that this could occur in polyoma infection by release of interferon, which selects cells able to use glycolytic pathways.

The selection theory implies that transformed cells pre-exist in the population, and in a clone they must

presumably be mutants. Such cells have never been found in the absence of virus, however, and since we know from the behaviour of artificial mixtures that growth of transformed cells is not inhibited by normal cells we can conclude that if they exist there must be less than 1 in 100,000 transformed cells pre-existing among the normal cells. As soon as virus is added, however, about three cells in 100 grow into transformed colonies in the absence of any selection by killing of other cells. This leaves little doubt that transformation is induced directly or indirectly by the virus-containing suspension and is not due to selection.

Evidence for Direct Action of Virus on Cells Transformed

It is still conceivable that the action of the virus is always cytotoxic, but that some product of virus-cell interaction induces the neoplastic change in neighbouring cells. It will be remembered that degenerative changes as well as neoplasia are produced *in vivo* after inoculation with the virus. There is plenty of cytotoxic activity in mouse fibroblasts, and failure to detect it in hamster cells could still allow at least 10% of cells to be killed by the virus. Is it possible that cytotoxic infection releases a cell product which is the real transforming agent? No one has yet found such a product which is not polyoma virus, and it is, anyway, unlikely for the following reason. After release from a virus-infected cell the transforming cell product would have to reach a susceptible cell and so its activity would be markedly affected by cell density. Once the cells have been exposed to virus and diluted, however, we find that the proportion transformed is not dependent on the cell density in the culture unless confluent sheets are formed (Stoker and Abel, 1962). It may still be argued that a cell product is transferred from a virus-infected to a normal cell in a clump of cells unaffected by dilution. We have found, however, that the proportion of transformed cells is not increased when the number of cell clumps rises.

There therefore seems to be no evidence in favour of an indirect action of the virus through an intermediate cell, and we tentatively assume that the virus acts directly on the cell which is transformed.

Possible Role of the Virus

We will now consider how neoplastic transformation may be caused by polyoma virus, but it will be apparent that we are almost at the end of factual data available at present, and what follows is largely hypothetical.

Following the evidence for a new antigen in tumour cells, we will assume that transformation is due to synthesis of one or more new proteins, and that the structure—that is, amino-acid sequence—of the protein is encoded in genetic material which will be referred to as the T gene or region. The following possibilities may be considered.

Mutation

Polyoma virus may act as a mutagen which leads to an altered base sequence and thence to an altered amino-acid sequence and protein of a new specificity. A mutagen would be expected to act on more than one site and therefore to produce several sorts of genetic change. It is possible that transformation is the only type of mutation which is expressed and that many others are undetected. The transformation rate itself is so high that this is unlikely. Alternatively, transformation and a T protein may be a common phenotypic

expression of a variety of different mutations. This could be investigated by searching for differences between transformed clones and by attempts to produce transformation of the same type with other mutagens. A mutation induced by the virus might also act by damaging a regulating gene rather than by directly leading to an altered T protein. This is discussed below.

Addition

The T gene may be introduced as part of the virus D.N.A., and integrated so that it replicates in step with the rest of the genetic material of the cell. The mechanism for such integration as we have seen must be somewhat different from that for prophage in lyso-genic bacteria, but this does not rule it out, particularly in a mammalian cell. Complete failure to induce such D.N.A. to form mature virus suggests that it must be defective; that is, lacking in part of the information required to build the complete virus particle, or that a repression is present which is not sensitive to the usual inducing agents. An alternative would be the acquisition of a new R.N.A. copy of the viral D.N.A. rather than the D.N.A. itself, but there is no evidence from other cell systems to support the maintenance of a permanent hereditary character through R.N.A. alone.

It should be noted that the added genetic material need not be attached to a chromosome. It could be a free particle, possibly in the cytoplasm, resembling the episome of the bacterial cell. Since the whole change could be due to production of one new protein the amount of added D.N.A. would be small relative to that of the original virus. Such additional material must be duplicated and distributed efficiently at each cell division, however; otherwise reversions to the normal character would occasionally appear.

Release

In the third hypothesis we suppose that the information for the new protein is not added by the virus genetic material, but is already present. The normal cells may all carry a T gene and have the potentiality to grow as neoplastic cells, but this potential behaviour may not be normally expressed. In current thinking this would be due to the activity of another part of the genetic material—that is, a regulator gene which provided a repressor substance (Jacob and Monod, 1961). Neoplastic transformation would then be due to release of the neoplastic potentiality through removal or inactivation of the repressor. In this way the virus would switch on the neoplastic behaviour of the cell by releasing the necessary information already contained in the cell.

Mechanisms of this type occur in the bacterial cell and are responsible for enzyme induction. The operative substance could be a small molecular weight derivative of the virus and the initial event would not necessarily require virus multiplication.

The principal objection to this hypothesis is that it does not allow for perpetuation of the change unless the virus also multiplies so as to continue the supply of the antirepressor substance. As we have seen, however, virus growth cannot be detected in transformed cells which continue to maintain their new character.

Is it possible that the virus removes the repressor permanently by destroying the regulator gene? This might be caused by a mutation or by genetic recombination involving replacement of the regulator gene by

non-functioning virus genetic material. Many cytotoxic D.N.A. viruses cause an early breakdown of the host cell D.N.A., and so alternatively we may postulate that polyoma-virus infection may give rise to an enzyme which destroys the relevant regulator gene without affecting the remainder of the genetic material of the cell. This would be difficult to conceive if the regulator gene concerned were part of a whole chromosome. If it were extrachromosomal—that is, an episome—and in particular cytoplasmic, however, it is possible that a viral-induced deoxyribonuclease of short life might be sufficiently selective.

Another possible mode of perpetuation would depend on a mechanism somewhat similar to that described for production of β -galactosidase by the bacterial cell (Jacob and Monod, 1961). Thus the antirepressor—that is, the inducing substance—may be present in the growth medium but unable to penetrate the cell wall. Virus infection may make the cell temporarily leaky, so that the inducing agent enters, combines with repressor, and allows the T-genetic region to operate. This in itself may alter the permeability of the cell surface to the inducing agent, so that the cycle can then be perpetuated indefinitely, though the virus has disappeared.

Possible Role of D.N.A.

At present it is impossible to choose between the "mutation," "addition," and the "release" hypotheses. The demonstration that the polyoma-induced T antigen differs from the antigen acquired in non-polyoma-induced tumours certainly suggests the addition of virus-specific genetic information. There remains, however, the possibility of several genes in normal cells, which when released by one carcinogen or another lead to different proteins but the same phenotypic change. It is important to know if neoplastic transformation can be effected by D.N.A. alone extracted from the virus. It has been shown that viral D.N.A. will give rise to a virus-multiplication cycle involving cell degeneration, and also that the D.N.A. will cause tumours in hamsters (DiMayorca *et al.*, 1959). It is not known if the tumours are the direct result of transformation by D.N.A. or whether they are due to complete virus produced first by the D.N.A. with hamster-cell degeneration. Abel and Crawford (1963) have shown that transformation requires virus particles with their full complement of D.N.A. "Empty" particles which are deficient in D.N.A. are not transforming. This observation, and the ability of free D.N.A. from rabbit papilloma to induce tumours (Ito and Evans, 1961) suggests that D.N.A. is the transforming agent.

Wider Implication of Virus Transformation

Any studies on the nature of the neoplastic process must obviously have some relevance to tumours in humans. Basically a neoplasm results from a change in the genetic behaviour of one cell, and ideally we should study that cell before and immediately after the change in character induced by an efficient carcinogen of rapid action. It is almost impossible to do this in the intact animal, let alone human, because most carcinogens act relatively slowly in the presence of a complex and changing population of different cell types, only one of which may be converted into a tumour cell, though the remainder may undergo confusing changes of no immediate relevance. For example, tumour cells may have a high and abnormal complement of chromosomes,

but the polyoma studies have shown that the essentials of the neoplastic change are not accompanied by a change in chromosome numbers, and that such changes occur later after prolonged cultivation. Moreover, it is not possible to observe the details of the whole sequence of events in one individual.

Induction of neoplasia in cultured cells removes many difficulties, particularly when this can be carried out on unmixed—that is, clonal—populations of cells. Of course, the objection can always be raised that this *in vitro* change may be different from the process occurring *in vivo*. Nevertheless, it has not been possible to distinguish between tumour cells induced *in vivo* and transformed cells produced *in vitro* with polyoma virus. The *in vitro* system therefore appears at present to be relevant to the initiation of a tumour, always remembering that subsequent development of the new cell population will naturally be very different in an intact host.

At present two viruses, Rous sarcoma virus and polyoma virus, are the only carcinogenic agents which will act sufficiently rapidly and precisely in tissue culture to allow such studies. Two other chicken-tumour viruses can now be detected by the *in vitro* activity, however, and should yield further information in due course (Baluda and Goetz, 1961; Rubin and Vogt, 1962). The recent observation that the virus of avian myeloblastosis will induce *in vitro* differentiation of granulocytic cells is especially relevant to the study of human leukaemia.

The polyoma system has certain technical advantages as a model system, such as its action on cells from animals (notably mice) whose genetics have already been the subject of intensive study, but polyoma virus and the chicken-tumour viruses are, anyway, very different. Polyoma is a small icosahedral D.N.A. virus assembled in the nucleus, and Rous virus is a larger R.N.A. virus, perhaps resembling influenza virus with assembly at the cell surface and a lipid-containing envelope. As recently pointed out by Rubin, the mechanism of neoplastic transformation may differ considerably in these two types of virus, and further study may indicate whether the final genetic change is also different. This would be important because it would remove the need for an all-embracing theory of carcinogenesis.

The studies on the action of these viruses under controlled conditions are important as opening up a method of studying other carcinogens in action apart from viruses. Studies on carcinogens *in vitro* are of particular relevance to human cancer because potential carcinogens can possibly be tested on human cells in culture though they cannot be used experimentally on humans. Several viruses have recently been isolated from laboratory-maintained human tumours and it will be difficult to assess their causal relationships except by an *in vitro* transformation test of this type.

Tumour Viruses and Man

Polyoma virus has been found to be a commensal of wild mice, but these mice do not commonly develop tumours, because they do not live long enough or because they do not receive a sufficiently large dose early in life. Thus we have a ubiquitous agent of potentially high activity but with a low probability of effect under natural conditions. Twenty or more other viruses are known as commensals of mice, and five of these can induce neoplasia under laboratory conditions. Moreover, 18 viruses have been found in the kidneys

of apparently healthy monkeys, and one of these, SV40, a vacuolating virus, will produce tumours under laboratory conditions (Eddy *et al.*, 1961).

There is no evidence to suggest that polyoma virus can produce tumours in humans. Polyoma virus is probably eaten by humans through contamination of food with mouse excreta rich in virus. Larger doses have probably been acquired by the increasing number of laboratory workers handling the virus since 1957. No human disease has been attributed to the virus so far, and polyoma antibodies have not been reported in humans. It has recently been reported, however, that the monkey virus SV40 (which closely resembles polyoma virus) will cause a transformation in cultured human cells (Koprowski *et al.*, 1962; Shein and Enders, 1962). It is too early to be sure that this is a neoplastic change, but it is clearly of great importance when vaccines for human use are prepared from monkey tissues.

In microbial flora man differs from animals only in kind, and if mice and monkeys carry many commensal viruses it is safe to assume that humans do as well, and one common human virus, adenovirus type 12, has now been found to cause lung tumours in hamsters. We may forecast that other human viruses may be found to induce tumours in experimental animals and perhaps neoplastic changes in human cells. It may be difficult, however, to assess the relevance of potentially active viruses as carcinogens under natural conditions. Some viruses, like polyoma virus, are not required for continued growth of the tumour cells once the genetic change has been induced. This means that a tumour may, anyway, be the wrong place to look for the virus which caused it.

If virus-induced neoplasia is generally accompanied by an antigenic change, as with polyoma virus, it is possible that tumour cells are frequently forming but are rejected in adults. Persistence of such cells would be more likely if they first appeared as a result of virus infection in infancy or before birth, and thus it may be more profitable to search for a virus aetiology in tumours of childhood rather than of adult life. The recently described malignant lymphoma in African children has epidemiological features which have suggested a virus infection (Burkitt, 1962), and this possibility is under intensive investigation.

It has been my resolve not to put forward any general theory of carcinogenesis and practically no reference has been made to other, non-viral, but highly active, carcinogens which have been extensively studied. If ubiquitous viruses were the prime cause, such agents might act by imposing selection conditions in favour of genetically altered cells with a low survival, as has been suggested by Alison (1961). At present it seems just as likely that there are several prime causes, even among the viruses.

I will conclude by stressing the need for quantitative studies at the cell level such as are possible with *in vitro* systems, and to suggest that, in view of recent developments, we should at least be cautious before excluding viruses completely in the aetiology of human tumours.

REFERENCES

- Abel, P., and Crawford, L. V. (1963). *Virology*. In press.
Abercrombie, M., and Ambrose, E. J. (1962). *Cancer Res.*, **22**, 525.
Alison, A. C. (1961). *Virology*, **15**, 47.
Axelrad, A. A., McCulloch, E. A., Howatson, A. F., Ham, A. W., and Siminovich, L. (1960). *J. nat. Cancer Inst.*, **24**, 1095.

M. G. P. STOKER: NEOPLASTIC TRANSFORMATION BY POLYOMA VIRUS

FIG. 1. — Electronphoto-
micrograph of polyoma
virus particles. ($\times 175,000$.)

FIG. 2.—Hamster fibroblasts
(BHK 21 strain). (Unstained.
 $\times 145$.)

FIG. 3.—Hamster fibroblasts
(BHK 21 strain) transformed
in vitro by polyoma virus.
(Unstained. $\times 145$.)

- Baluda, M. A., and Goetz, I. E. (1961). *Virology*, **15**, 185.
- Burkitt, D. (1962). *Postgrad. med. J.*, **38**, 71.
- Crawford, L. V. (1963). *Virology*. In press.
- and Crawford, E. M. (1961). *Ibid.*, **13**, 227.
- DiMayorca, G. A., Eddy, B. E., Stewart, S. E., Hunter, S. W., Friend, C., and Bendich, A. (1959). *Proc. nat. Acad. Sci.*, **45**, 1805.
- Eddy, B. E., Borman, G. S., Berkeley, W. H., and Young, R. D. (1961). *Proc. Soc. exp. Biol. (N.Y.)*, **107**, 191.
- Rowe, W. P., Hartley, J. W., Stewart, S. E., and Huebner, R. J. (1958a). *Virology*, **6**, 290.
- Stewart, S. E., and Grubbs, G. E. (1958b). *Proc. Soc. exp. Biol. (N.Y.)*, **99**, 289.
- Young, R., and Mider, G. B. (1958c). *J. nat. Cancer Inst.*, **20**, 747.
- Kirschstein, R. L., and Young, R. D. (1959a). *Nature (Lond.)*, **183**, 766.
- Stanton, M. F., and Marcotte, J. M. (1959b). *J. nat. Cancer Inst.*, **22**, 161.
- Forrester, J. A., Ambrose, E. J., and Macpherson, I. A. (1962). *Nature (Lond.)*, **196**, 1068.
- Fraser, K. B., and Gharpuré, M. (1962). *Virology*, **18**, 505.
- Groman, N. B. (1953). *J. Bact.*, **66**, 184.
- Gross, L. (1953). *Proc. Soc. exp. Biol. (N.Y.)*, **83**, 414.
- Habel, K. (1961). *Ibid.*, **106**, 722.
- Ham, A. W., McCulloch, E. A., Axelrad, A. A., Siminovitich, L., and Howatson, A. F. (1960). *J. nat. Cancer Inst.*, **24**, 1113.
- Siminovitich, L., and Howatson, A. F., and Axelrad, A. A. (1961). In *Tumour Viruses of Murine Origin*, pp. 284–295. Ciba Foundation Symp.
- Harris, R. J. C., Chesterman, F. C., and Negroni, G. (1961). *Lancet*, **1**, 788.
- Howatson, A. F., and Almeida, J. D. (1960). *J. biophys. biochem. Cytol.*, **8**, 828.
- McCulloch, E. A., Almeida, J. D., Siminovitich, L., Axelrad, A. A., and Ham, A. W. (1960). *J. nat. Cancer Inst.*, **24**, 1131.
- Huebner, R. J., Rowe, W. P., Hartley, J. W., and Lane, W. T. (1961). In *Tumour Viruses of Murine Origin*, pp. 314–328. Ciba Foundation Symp.
- Iseki, S., and Sakai, T. (1953). *Proc. Japan Acad.*, **29**, 1271.
- Ito, Y., and Evans, C. A. (1961). *J. exp. Med.*, **114**, 485.
- Jacob, F., and Monod, J. (1961). *Cold Spr. Harb. Symp. quant. Biol.*, **26**, 193.
- Koprowski, H., Ponten, J. A., Jensen, F., Ravdin, R. G., Moorhead, P., and Saksela, E. (1962). *J. cell. comp. Physiol.*, **59**, 281.
- Macpherson, I. A. (1963). *J. nat. Cancer Inst.* In press.
- and Stoker, M. (1962). *Virology*, **16**, 147.
- Negroni, G., Dourmashkin, R., and Chesterman, F. C. (1959). *Brit. med. J.*, **2**, 1359.
- Rowe, W. P., Hartley, J. W., Brodsky, I., Huebner, R. J., and Law, L. W. (1958). *Nature (Lond.)*, **182**, 1617.
- Law, L. W., and Huebner, A. J. (1959). *J. exp. Med.*, **109**, 449.
- Rubin, H. (1955). *Virology*, **1**, 445.
- and Vogt, D. K. (1962). *Ibid.*, **17**, 184.
- Sachs, L., and Medina, D. (1961). *Nature (Lond.)*, **189**, 457.
- Shein, H. M., and Enders, J. F. (1962). *Proc. Soc. exp. Biol. (N.Y.)*, **109**, 495.
- Sjögren, H. O., Hellström, I., and Klein, G. (1961). *Cancer Res.*, **21**, 329.
- Smith, J. D., Freeman, G., Vogt, M., and Dulbecco, R. (1960). *Virology*, **12**, 185.
- Stewart, S. E., Eddy, B. E., Gochenour, A. M., Borgese, N. G., and Grubbs, G. E. (1957). *Ibid.*, **3**, 380.
- Stanton, M. F., and Berkeley, W. M. (1958). *Proc. Amer. Ass. Cancer Res. (Abstract)*, **2**, 348.
- Stoker, M. (1960). *Brit. J. Cancer*, **14**, 679.
- and Abel, P. (1962). *Cold Spr. Harb. Symp. quant. Biol.* In press.
- and Macpherson, I. A. (1961). *Virology*, **14**, 359.
- Temin, H. M., and Rubin, H. (1958). *Ibid.*, **6**, 669.
- Vogt, M., and Dulbecco, R. (1960). *Proc. nat. Acad. Sci.*, **46**, 365.
- (1962). *Virology*, **16**, 41.
- Watson, J. D., and Littlefield, J. W. (1960). *J. molec. Biol.*, **2**, 161.
- Wildy, P., Stoker, M. G. P., Macpherson, I. A., and Horne, R. W. (1960). *Virology*, **11**, 444.
- Williams, M. G., Howatson, A. F., and Almeida, J. D. (1961). *Nature (Lond.)*, **189**, 895.
- Winocour, E., and Sachs, L. (1961). *Virology*, **13**, 207.

CORRELATION OF RADIOLOGICAL AND HISTOLOGICAL FINDINGS IN IDIOPATHIC STEATORRHOEA

BY

J. W. LAWS, M.B., M.R.C.P., F.F.R.

H. SHAWDON, M.B., M.R.C.P., D.M.R.D.

C. C. BOOTH, M.D., M.R.C.P.

J. S. STEWART, M.B., M.R.C.P.

From the Departments of Radiodiagnosis and Medicine, Postgraduate Medical School of London, Hammersmith Hospital, London

[WITH SPECIAL PLATE]

There have been many studies of the radiological changes in the small bowel of patients judged on clinical and biochemical grounds to have idiopathic steatorrhoea. In this study the diagnosis has been supported by histological evidence obtained by biopsy, and the radiological changes are correlated with the mucosal appearances. A similar study has been carried out in tropical sprue (Floch, Caldwell, and Sheehy, 1962), but not hitherto in idiopathic steatorrhoea.

The early reports of radiological abnormalities in the small bowel in idiopathic steatorrhoea (Mackie, 1933; Snell and Camp, 1934; Mackie, Miller, and Rhoads, 1935; Golden, 1936, 1941; Ettinger, 1949) were chiefly descriptions of "flocculation" (Golden, 1941) or "clumping" (Berridge, 1951) of a barium sulphate suspension. It was then shown that flocculation could occur in normal children (Zwerling and Nelson, 1943; Teall, 1945) and in newborn infants before they have received milk (Astley and French, 1951) and that it could be induced in normal subjects by the presence of excess fatty acids and mucus in the small bowel (Frazer, French, and Thompson, 1949). It was not until the development of a flocculation-resistant barium preparation that the additional radiological abnormalities of dilatation of the small bowel and thickening of the

transverse mucosal folds were clearly demonstrated in idiopathic steatorrhoea (Ardran, French, and Mucklow, 1950), coeliac disease (Astley and French, 1951; Anderson, Astley, French, and Gerrard, 1952), and tropical sprue (French, Gaddie, and Smith, 1956; Paterson and Baker, 1958).

This paper correlates the radiological findings with clinical, biochemical, and biopsy data in 28 patients with idiopathic steatorrhoea.

Patients Studied.—Twenty-eight patients (11 male and 17 female) were studied. Their ages ranged from 14 to 83 years. Six of these had a history of coeliac disease in childhood. The remainder presented in adult life with no childhood history of intestinal disturbance. Intestinal biopsy showed partial or subtotal villous atrophy at the time of the study in all patients.

Methods

Radiological

Radiological examination of the small bowel was carried out at or near the time of the patient's first visit or admission to hospital. In almost all cases 100 ml. of undiluted "micropaque," a partially stabilized micro-suspension of barium sulphate, was given at the first