THE SHAPE OF A PLEURAL EFFUSION

BY

STANLEY DAVIS, M.R.C.P., F.F.R.

FRANCES GARDNER, F.R.C.P.

AND

GEORGE QVIST, F.R.C.S.

From the Royal Free Hospital, London

[WITH SPECIAL PLATE]

The upper limit of a free pleural effusion is usually described as having the shape of a curved line higher in the axilla than at the front and back of the pleura, this concept being based on clinical and radiological evidence.

The clinical sign frequently quoted is a rising line of dullness to percussion extending from the front and back of the chest up into the axilla. Ellis (1874) is usually credited with the original description of this and it has been known as the Ellis S-shaped line, though in a later publication (1876) this author gave full credit to Damoiseau, who had described a similar finding in 1843.

Radiological evidence appears to confirm the Ellis line. In a postero-anterior radiograph of good penetration a pleural effusion produces an area of increased density bounded superiorly by a line which rises in the lateral portion of the chest.

The literature on pleural effusion contains very little criticism of this classic description. It was suggested by Fleischner (1926) that the rising line of radiographic density can be explained by the axillary rays hitting the body more tangentially and having to pass through a greater thickness of fluid than the medial rays. Kaunitz (1935) demonstrated by the injection of "lipiodol" (iodized oil) in "gomenol" (purified niaouli oil) that this light opaque oil floats on the surface of the fluid at a higher level than the curved dense shadows seen on radiography.

In this paper it is suggested that neither the Ellis line of dullness nor the radiological oblique line of density corresponds to the true upper limit of a free pleural effusion. It is submitted that in fact the true upper limit is horizontal and, indeed, that if it were not so the typical x-ray appearances of a pleural effusion would not be obtained.

Physical Considerations

Various physical factors have been proposed to explain the upper curved line of a pleural effusion. Capillary attraction between the pleural surfaces has been described as the cause of the rising axillary line.

Kaunitz (1935), experimenting with balloons and lungs in containers of different shape, concluded that the characteristic curved line of a simple effusion depends to a large extent on the normal negative intrapleural pressure elevating the liquid through maintaining the lung in a distended condition, "while close proximity of the upper part of the lung and chest wall favours capillarity, so that the liquid is raised to a still higher level."

In our opinion, no satisfactory explanation on physical grounds has yet been proposed to support the concept of an upper curved limit to a pleural effusion.

On the contrary, it is easy to find evidence of a simple physical nature to explain an upper limit of horizontal shape.

Among the various forces which determine the shape of a free pleural effusion, two of the most important must be the effects of gravity and the resistance of the compressed lung. If the elasticity of the lung is uniform, the thickness of fluid in the pleura will become less from below upwards, owing to gravity. The upper limit of fluid is thus a thin horizontal edge. If there are no complicating lesions such as consolidation or adhesions, distribution of the fluid and its horizontal level will vary with the position of the patient.

The following evidence is submitted to support the contention that the upper limit of a free pleural effusion is horizontal.

Experimental Evidence

1. A coarse representation of the shape of a pleural effusion was made from a plaster cast. This was tapered in thickness from below upwards, but of uniform thickness in the horizontal planes. Radiographs of three different penetrations of the plaster model suggest that the density of the cast is greater towards the middle of its curvature though in fact the plaster material is of uniform consistence. The line of density in all three radiographs rises laterally until it reaches the true height of the plaster cast (Special Plate, Fig. 1). This graduation of density in the x-ray film is, of course, due to the variation in depth of plaster through which the x rays pass.

2. A plastic bucket was partly filled with water and an air-filled balloon immersed in it to simulate the physical conditions existing when aerated lung is partly surrounded by an effusion. Radiographs were taken of this combination (Text, Fig. 1). Though the fluid was known to be at the same level all round the balloon, it is not possible to see on the radiograph the upper limit of fluid in the central portion of the bucket. The fluid shadow is shown at a lower level extending upwards to the periphery until it coincides with the true fluid level. This rising line of density is shown in all positions of rotation of the bucket and contents. Here again the rising line of radiographic density is due to the greater layer of fluid through which the x rays pass towards the periphery.

It is suggested that in radiography of a pleural effusion a similar effect is obtained, the rising line of density towards the axilla being due to the greater depth of fluid in the antero-posterior dimension through which the x rays have to pass (Special Plate, Fig. 2), as originally proposed by Fleischner (1926). In the lateral view the x rays traverse the greatest depth of fluid at the front and back of the pleura, so that

TEXT FIG. 1.—Radiograph of water-filled bucket with air-balloon to demonstrate the rising line of density at the periphery though the thickness of water surrounding the air-balloon is uniform in the horizontal planes.

S. DAVIS, F. GARDNER, AND G. QVIST: SHAPE OF A PLEURAL EFFUSION

FIG. 1.—Plaster cast to demonstrate the shape of a pleural effusion. Radiographs of different penetration all show a rising line of density though the plaster cast is of uniform thickness in the horizontal planes.

FIG. 2.—Radiograph of pleural effusion with diagrammatic transverse section to demonstrate rising axillary line of density. 400 ml. of fluid was aspirated from radiotranslucent area at site of marker.

S. DAVIS, F. GARDNER, AND G. QVIST: SHAPE OF A PLEURAL EFFUSION

FIG. 3.—Lateral radiographs of pleural effusions showing (A) rising anterior line of density and (B) lines of density in greater and lesser fissures, due to "end-on" views of fluid.

C. D. CALNAN: CREAM FOR COVERING SKIN BLEMISHES

FIG. 1.—Showing extent of disfigurement.

FIG. 2.—Masking effect of covering cream.

the upper limit of density rises anteriorly behind the sternum and posteriorly in front of the spine (Special Plate, Fig. 3A).

A similar explanation applies to the apparent predilection of the fluid for the greater and lesser fissures as seen on a lateral chest radiograph (Special Plate, Fig. 3B). Provided the effusion is free and not encysted, the upper limit of the fluid in the fissures is no higher than that surrounding the rest of the lung, but it is radio-opaque because it is seen "end-on" and the rays have to traverse a greater thickness of fluid.

Clinical Evidence

1. It is a common experience in aspiration of pleural effusion that the volume of fluid withdrawn is often considerably greater than might be expected from the x-ray appearance.

2. Successful aspiration of a pleural effusion may be performed at a level above the radiographic shadow of fluid. This is demonstrated in Special Plate, Fig. 2, from a patient who had a chronic simple pleural effusion.

A simple comparison may be made by observing that in the experiment with the plastic bucket and air-filled balloon, if a needle were introduced through the wall of the bucket in the radiotranslucent area, fluid would obviously be withdrawn, though none is demonstrable in the x-ray film in that position (Text, Fig. 1).

Discussion

It is suggested that physical, experimental, and clinical data support the contention that the upper limit of a pleural effusion is horizontal, at a level indicated approximately by the top of the x-ray shadow of fluid

in the axilla in the antero-posterior film. It is probable that this is an underestimate and that the upper limit of fluid is actually higher than this point, because very often, when there is enough fluid to give a good oblique x-ray shadow, it will be noted that there is also a pleural cap of fluid at the apex. This is shown in Special Plate, Fig. 2, where fluid

TEXT FIG. 2.—Diagram to show the upper limit of a pleural effusion as demonstrated by radiography, percussion, and the authors' observations.

obviously reaches the apex of the pleura but there is still a radiotranslucent area of lung in the mid-zone, and, in fact, fluid was aspirated from this site.

It can be concluded from this evidence that clinical signs are more accurate than radiography in determining the extent of a pleural effusion: dullness to percussion may be present at the front and back of the chest over a radiotranslucent area of lung. If the upper limit of an effusion is horizontal, it might be expected that the upper limit of dullness to percussion would also be horizontal around the chest. It is possible

however, that physical conditions of partly aerated lung and the more acute curvature of the axilla could be responsible for the rising axillary line of impaired percussion described as the Ellis line. It is therefore suggested that, if the Ellis line is demonstrable, it does not indicate the true upper limit of fluid which is actually at a horizontal level corresponding, probably, to the highest point of the Ellis line (Text, Fig. 2).

Summary

Physical, experimental, and clinical evidence is presented to suggest that the upper limit of a free pleural effusion is horizontal. The level of this limit is indicated approximately by the highest point of dullness to percussion and by the apex of the axillary shadow in the antero-posterior radiograph.

It is suggested that radiological signs of fluid are misleading, and that clinical signs are more accurate in estimating the extent of a pleural effusion. Fluid may be aspirated from the pleura in a radiotranslucent area.

The Ellis line of dullness to percussion is not always demonstrable in cases of pleural effusion and it does not indicate the true extent of fluid.

We are greatly indebted to Miss J. A. Middleton, artist of the Medical Unit, and to Miss L. Hobson for the drawings, and to Mr. R. R. Phillips, Photographic Department of the Royal Free Hospital, for the photographs.

REFERENCES

- Damoiseau, H. (1843). Cited by Ellis, C. (1876).
Ellis, C. (1874). *Boston med. surg. J.*, 90, 13.
— (1876). *Ibid.*, 95, 689.
Fleischner, F. (1926). *Ergebnisse der medizinischen Strahlenforschung*, 2, 206. Thieme, Leipzig.
Kaunitz, J. (1935). *J. thorac. Surg.*, 4, 300.

A NEW CREAM FOR COVERING SKIN BLEMISHES

BY

C. D. CALNAN, F.R.C.P.

The Institute of Dermatology, St. John's Hospital for Diseases of the Skin, London, W.C.2

[WITH SPECIAL PLATE]

Whatever personal opinion one may hold as to the most desirable relationship between cosmetology and dermatology, few doctors will doubt that each can with advantage learn something from the particular skills of the other. Cosmeticians have made a particular study of the "acceptability" characteristics of the preparations which they apply to the skin. Dermatologists should therefore be able to learn some of the art of local treatment from cosmeticians.

A number of dermatological disorders are not, at the present time, amenable to treatment. Surgical repair may not be feasible or it may not produce a sufficiently satisfactory cosmetic result. This applies to certain congenital disorders producing skin blemishes which may be regarded as purely cosmetic disabilities.

A new type of covering cream* for the treatment of such lesions has recently been developed, and used on more than 50 patients. One of the first of these was a girl of 16 with an extensive port-wine stain. The result was so gratifying that one was encouraged to treat

* "Keromask," made by Innox (England) Limited, 436 Essex Road, London N.1.