Preliminary Communications

Refrigerated Femoral Venous-Venous Haemodialysis with Coil Preservation for Rehabilitation of Terminal Uraemic Patients

The rehabilitation of patients with terminal renal failure to a useful and economic existence has been achieved by periodic haemodialysis, and patients are now in their fourth year of treatment (Hegstrom *et al.*, 1962; Scribner, personal communication, 1963). The technique utilizes a low-flow pumpless refrigerated haemodialysis system with a permanent arteriovenous sialastic "teflon" shunt (Quinton *et al.*, 1962). The major disadvantages of this system have been the difficulty in maintaining the patency of the arteriovenous fistula prosthesis (Kolff *et al.*, 1962), inaccessibility of the prosthesis to the patient's own hands, and unpredictable blood-flow rates in a pumpless haemodialysis system with consequent difficulty in predictable ultrafiltration and removal of excess body water.

An alternative system of periodic haemodialysis for the rehabilitation of terminal uraemic patients has been developed at the Royal Free Hospital since September, 1961. The description and preliminary results of this technique will be reported elsewhere (Rae *et al.*, 1963; Shaldon *et al.*, 1963). These results demonstrated that the patient with absent renal function requires a minimum of 24 hours haemodialysis per week, with blood-flow rates of 200 ml. a minute through a twin-coil

kidney. The technique now utilizes a femoral venous-venous percutaneous catheterization system (Shaldon *et al.*, 1961) with a pumped, refrigerated, continuously heparinized haemodialysis using a twin-coil kidney and subsequent preservation of the dialysis circuit between dialyses. This has reduced the annual cost of haemodialysis to £500 per patient and reduced the blood requirement to 500 ml. a week.

The object of this report is to detail the most recent development, which has been the successful use of a permanent twin

femoral venous prosthesis (Fig. 1) that permits easy access to the patient's venous system. Two teflon catheters are inserted into the same femoral vein by a percutaneous technique (Shaldon *et al.*, 1961) and their tips situated in the mid inferior vena cava (Fig. 2). Patency of the catheters is maintained by an asbestos capillary heparin perfuser (Giovannetti *et al.*, 1963), which perfuses the catheters with 0.25 ml. per hour of a concentrated heparin solution (500 units/ml.). The perfuser is charged each day through a syringe and disconnected prior to haemodialysis. The present prosthesis has functioned perfectly for two months with 24 haemodialyses, and the patient now comes into hospital for two nights a week, and is able to connect himself to the artificial kidney and disconnect himself. The entire running of the dialysis system is done by a trained nurse without a doctor seeing the patient.

CASE HISTORY

A 30-year-old male draughtsman developed malignant hypertension and progressive renal failure. He was referred to the Royal Free Hospital Renal Unit three months after the onset of symptoms in a terminal uraemic state, vomiting, drowsy, and virtually blind from advanced papilloedema and with gross fundal haemorrhages and exudates with bilateral macular involvement. His blood-pressure was 260/160. Investigations revealed: blood urea, 370 mg./100 ml.; serum creatinine, 21 mg./100 ml.; serum sodium, 120 mEq/l.; serum potassium, 6.3 mEq/l.; serum bicarbonate, 12 mEq/l.; creatinine clearance, 0.5 ml./min.; P.C.V., 14%; chest x-ray picture normal; aortography, bilateral small renal arteries and contracted kidneys; E.C.G. normal.

In view of the recent onset of his symptoms and the absence of cardiac involvement it was decided to embark on a programme of chronic haemodialysis. The prognosis had been explained to the patient and he agreed to a trial of haemodialysis. Initially he was treated by short daily haemodialysis with progressive ultrafiltration. Between dialyses he received a fluid intake of 500 ml. with a diet containing 10 mEq of sodium and 30 g. of protein. Within one week his blood-pressure was stabilized at 120/80 and

FIG. 1.—Femoral-vein catheters *in situ* (after two months) with heparin perfuser unit attached. The white area at the site of the puncture is due to the antibiotic skin powder.

FIG. 2.—Plain x-ray film of the abdomen to show the position of the teflon catheter in the inferior vena cava.

his vision began to improve. (He was not receiving hypotensive therapy.) After 14 short haemodialyses over a period of three weeks he was started on a twice-weekly 14-16-hour overnight dialysis programme. He left hospital with his eyesight fully restored five weeks after starting the programme.

He is now completely asymptomatic, with a blood-pressure of 130/70, and ready to start work again. His blood urea rises to 100 mg./100 ml. pre-dialysis and falls to 25-30 mg./100 ml. post-dialysis. He is able to set up his own dialysis circuit, connect himself to the artificial kidney, and disconnect himself, and requires only a trained nurse to monitor the dialysis during his sleeping period. He enters the hospital at 4.30 p.m. on Mondays and Fridays and leaves at 9 a.m. on Tuesdays and Saturdays. He eats and drinks what he likes during dialysis, but any gain in weight is controlled by ultrafiltration. Between dialysis he remains on a diet with 10 mEq of sodium and 30 g. of protein, and 25 g. of "resonium" daily to control hyperkalaemia.

COMMENT

The ability to control uraemia and hypertension by a haemodialysis system which the patient is able to run himself has not previously been reported. The relatively low cost of this system—as the coil can be reused on as many as 8 to 10 occasions—and the low blood requirement make it an economic proposition for the treatment of uraemia and hypertension in carefully selected individuals.

The advantages of the femoral venous-venous prosthesis are the ease of insertion of the catheters without loss of the vessel and the accessibility of the catheters to the patient's hand (an essential prerequisite to self-dialysis). The absence of arterial catheterization renders the system safer if a catheter should fall out, as the patient has been instructed in the control of venous haemorrhage. The dangers of sepsis have not materialized because of very careful prophylaxis. The skin of the thigh has been cleaned daily with chlorhexidine and the puncture sites are sprayed with neomycin and bacitracin powder, resulting in clean

granulations without induration around the punctures. The danger of thrombosis to the vessels exists, but as there is a rapid flow of blood in the femoral vein and inferior vena cava and a state of regional heparinization is achieved (without significant prolongation of whole-blood clotting-times) this possibility is regarded as unlikely. However, the patient has been instructed to report immediately should swelling of the ankle of the catheterized limb occur.

The possibility of electronic monitoring replacing nursing monitoring must be considered. If this could be achieved the ultimate aim of home dialysis could result in the treatment of larger numbers of patients at present dying with terminal renal failure.

We would like to thank Dr. K. Ball for referring the patient; Nurses J. Barnett, P. Fowler, F. Giles, M. Higgins, B. Quennell, and S. V. Tooze for invaluable and skilled assistance; and Misses K. O'Connor, J. Ryder, and J. Hird for biochemical assistance. The teflon catheters and heparin perfusers will shortly be available from the Watson-Marlow Air Pump Co., Marlow, Bucks.

STANLEY SHALDON, M.D., M.R.C.P.
A. I. RAE, M.B., M.R.C.P.
S. M. ROSEN, M.B., M.R.C.P.
H. SILVA, M.D.
J. OAKLEY, S.R.N.

Renal Unit, Royal Free Hospital, London.

REFERENCES

- Giovannetti, S., Bigalli, A., Cioni, L., Della Santa, M., and Balestri, P. L. (1963). *Acta med. scand.*, **173**, 1.
Hegstrom, R. M., Murray, J. S., Pendas, J. P., Burnell, J. M., and Scribner, B. H. (1962). *Trans. Amer. Soc. art. Intern. Org.*, **8**, 266.
Koff, W. J., Nakamoto, S., and Scudder, J. P. (1962). *Ibid.*, **8**, 292.
Quinton, W. E., Dillard, D. H., Cole, J. J., and Scribner, B. H. (1962). *Ibid.*, **8**, 236.
Rae, A. I., Rosen, S. M., Silva, H., Pomeroy, J., Bienenstock, J., and Shaldon, S. (1963). *Proc. roy. Soc. Med.* In press.
Shaldon, S., Chianussi, L., and Higgs, B. (1961). *Lancet*, **2**, 857.
—, Rosen, S. M., and Silva, H. (1963). Second Symposium on Water and Electrolyte Metabolism, Amsterdam. In press.

Medical Memoranda

Spontaneous Rupture of Aortic Aneurysm into Inferior Vena Cava

Aneurysms of the abdominal aorta rarely rupture into the inferior vena cava. We have found no report of this occurrence in the British literature of the last 40 years, though 13 have appeared in other countries (Lehman, 1938; Dussaillant *et al.*, 1949; Viar and Lombardo, 1952; Cooley, 1955; Eiseman and Hughes, 1956; Shumacker and Jontz, 1956; Buzzi *et al.*, 1957; Zampi, 1957; Bulgrin and Jacobson, 1958; Greenstone *et al.*, 1959; Williams and Webb, 1959; Thistlethwaite *et al.*, 1960; Gerstle de Pasquet *et al.*, 1961). The increase in reports during recent years may reflect the greater possibility of successful treatment rather than a rising incidence. The true incidence is probably higher than is suggested by the number of reports, since the condition can be rapidly fatal and so escape diagnosis. Though our patient was too ill to withstand an operation, we are reporting his case to draw attention to a condition in which surgery, though hazardous, can now be life-saving.

CASE REPORT

A labourer aged 59 was admitted to Ashford Hospital complaining of severe and increasing pain in the lumbosacral region for a week. In the previous two days the pain had spread to the abdomen and genitals. He described it as a continuous, drawing pain. For six days he had eaten little and had no bowel motion. He thought there had been a swelling in his right groin for about 35 years. In 1946 he had noticed right-sided varicose veins, and these were tied in 1951. In 1953 he had had five days off work on account of low back pain.

He was a thin, elderly-looking man who was ill, restless, short of breath, and in great pain. His head and trunk moved perceptibly with each heart-beat. The pulse was regular at 130 a minute and collapsing in type. The systolic blood-pressure was 115 mm. Hg in both arms; the sounds faded gradually when the pressure in the cuff fell below 50 mm. but remained clearly audible at zero. There was a large mass in the abdomen at the level of the umbilicus, extending from the midline towards the right flank. Expansile pulsation was felt in it, and a continuous thrill spread out radially from it. A "machinery" murmur was audible over the whole abdomen, but was loudest over the mass. Large pulsating swellings were seen on both femoral arteries in the groins. There were many tortuous distended veins in both legs and thighs; the blood flow in them was upward and the pressure 12 cm. of saline above the sternal angle. The jugular venous pressure was estimated to be 5 cm. above the sternal angle. The cardiac impulse was