DERMATOMYOSITIS, WITH REPORT OF A CASE WHICH
PRESENTED A RARE MUSCLE ANOMALY BUT
ONCE DESCRIBED IN MAN.

BY WALTER RALPH STEINER, A. M., M. D., HARTFORD, CONN.
(Formerly Medical House Officer, The Johns Hopkins Hospital.)

Inflammatory changes in muscles have long been recognized as occurring independently or in connection with certain diseases. Hunter (1) is said to have described cases of myositis in 1784 and to have noted the insusceptibility of muscular tissue to inflammation and its consequences. I have been unable to find any such reference in his writings. During the past century cases have been published in many different countries and a considerable amount of literature is to be found on this affection.

A multiple muscle inflammation, however, presenting also other well known symptoms, was not described as such until 1887. In that year E. Wagner (2) of Leipsic, Unverricht (3) of the Polyclinic at Jena, and Hepp (4) of Kussmaul's clinic at Strasburg, almost simultaneously reported cases. Though Wagner's case was first published, it was in reality the last seen, for Hepp made his observation two months previously and Unverricht's case had been noted seven years before either of the above two. The pathological findings, with a meager history of this last case, were presented by Marchand (5) to the Schlesische Gesellschaft für vaterländ. Cultur on October 22, 1880, and later appeared in the society's transactions and in the Breslauer ärztliche Zeitschrift of that same year. The case has caused a good deal of confusion among writers on this disease. Herrick (6), Jacoby (7), and others have described it as another case of an earlier observation, while Lewy (8) thinks it is an additional one which Unverricht has later more fully reported. This error may be due to the fact that Unverricht in two of his articles

made the date of seeing the patient a year later than it should have been. Yet a careful examination will convince one that the same case is dealt with. Hepp's case was also reported by Henry Jackson (9) of Boston in 1887—a fact which seems to have been entirely overlooked.

Before these observations we find two others which, although representing cases rather meagerly described and otherwise diagnosed, should undoubtedly be considered as bearing on this disease. I refer to a case seen by E. Wagner (10) in 1863, and to a later one observed by Potain (11) in Paris. They were diagnosed at the time as instances of periostitis and an atypical form of glanders. Following the publication of the first three recognized instances Jacoby (12) and Löwenfeld (13) gave additional cases and discussed the previous papers on the subject. A little later Unverricht (14) presented the clinical history of another case, which differed from those previously reported in terminating favorably. The close association in this patient of dermatitis with a multiple muscle inflammation led Unverricht to coin the term *Dermatomyositis*, which has been frequently adopted to designate this affection. Boeck (15) and Strümpell (16) subsequently published other cases, the latter giving an excellent account of all that was known of the disease.

In the following years many German writers reported so-called cases which were subjected to a critical examination in 1896 by R. Pfeiffer (17), and the histories given by Schnell (18), Lewy (19) (three out of his four cases), Fraenkel (20), Buss (21), Herz (22), Albu (23), Schultze (24), and Schultzen (25) were, rightly we think, adjudged as atypical and belonging to another disease. The cases of v. Kornilow (26), Jollasse (27), and Thiele (28) can be set aside on the same grounds. Neubauer's case (29) more nearly resembles polymyositis haemorrhagica. During these years, however, Fuckel (30), Senator (31), Lewy (32), Köster (33), and Kell (34)¹ have given us undoubted examples of the disease, while Gowers

¹Two of Kell's three cases have not been added to our list as they were of a doubtful abortive type and somewhat atypical in character.

(35) has reported a case which probably should not be classed here, as it was associated with a polyneuritis. More recently Frohmann (36), Bonnet (37), Lepine (38), Janowsky and Wyssokowicz (39), Bacialli (40), Oppenheim (41), and Christen (42) have added to the number already recorded. Forcheimer's case (43) I have also placed in this list, although it is not as typical as the others. It is interesting to note that one of Oppenheim's cases, reported four years ago as a typical instance of this affection, now presents all the symptoms of scleroderma. During the past two years Vincent (44), Schüller (45) and Jessen (46) have likewise published histories of patients with polymyositis, but none of them had the special type of disease which would entitle them to admission to our group. From time to time different writers have collected the previously existing cases but, with the exception of Lorenz (47) and R. Pfeiffer (48), they appear to have done so without any discrimination. Lewy was able to find twenty-one cases in 1893, while Köster, three years later, added two more to that number. Both of these writers, however, included atypical cases and those of a closely allied affection—polymyositis haemorrhagica.

We have found, including our own, twenty-eight quite typical histories, which are classified according to countries as follows: Germany, 19; France, 2; Sweden, 2; Italy, 1; Cuba, 1; United States, 3; total 28.

Gowers' case is the only one as yet observed in England. In Italy two cases of polymyositis haemorrhagica have been described and one of dermatomyositis, while, though several writers have written on this affection, the only case yet published in Russia (49)² belongs to another type of myositis.

Dermatomyositis may be defined as an acute, subacute, or chronic disease of unknown origin, characterized by a gradual onset with vague and indefinite prodromata, followed by oedema, dermatitis, and a multiple muscle inflammation.

² I desire to thank Dr. Thayer for translating this case from the Russian for me.

ETIOLOGY.—On this point we are as much in the dark as were the three writers who first described the disease. Hepp and others have spoken strongly in favor of its being due to an infection and the presence of a splenic tumor, fever, and angina, seem to support this view. From Senator's time the following three theories as to its origin have been advanced:

I. It is due to a specific microorganism (vegetable parasite). This view is gradually losing ground. Bacteria have been frequently sought for in the inflammatory oedema and in the tissues, with negative results in every instance save one (Bacialli's case), and cases of this disease reported as being due to certain bacteria have been shown to belong to another class of muscle inflammation. Bauer (50) and Georgievski (51) have recently reported cases of the allied affection—polymyositis haemorrhagica—which were probably due to *Staphylococcus pyogenes aureus* and *Staphylococcus pyogenes albus*, obtained in pure cultures from the tissues. In Bacialli's case, however, the *Staphylococcus pyogenes albus* was said to be very much attenuated in virulence. His findings, as well as Georgievski's, can hardly be considered as conclusive in view of the universal presence of *Staphylococcus albus* in the deep layers of the skin (Thayer).

II. It is due to an animal parasite. This idea was first advanced by Unverricht, who thought the parasite might belong to the gregarinae. In support of this theory L. Pfeiffer (52) states that the muscle findings in this disease are similar to those seen in the muscles of horses and dogs infected with gregarinae, and he says that Virchow (53) has mentioned cutaneous rashes in hogs that were similar to the changes in the skin occasionally found in dermatomyositis. The only instance, however, in which these parasites were found in man was reported in 1891 by Klebs (54), who observed them in a case of muscular atrophy, where they were at first taken for muscle nuclei. Negative results have attended the search for them in seven cases, in two of which L. Pfeiffer also examined the sections. This unsuccessful quest Pfeiffer would explain as due to the fact that the methods of preserving the tissues and preparing the specimens mili-

tate against a ready recognition of the parasite. They may be present but be overlooked, or, although they can cause a muscle inflammation by their presence or their toxin, the muscles containing them may escape microscopic examination.

III. It is due to a toxin. This view was given prominence by Senator, to whom it was suggested by his second case, which began with symptoms of gastro-intestinal irritation after the patient had eaten stale crabs. Kell's case is also somewhat similar, the symptoms appearing a few hours after the ingestion of fish. No gastro-intestinal symptoms were here noted. In addition to these, Boeck's case has been put in this group, inasmuch as the disease was observed after the energetic rubbing of copaiba balsam into the skin. Boeck, however, does not consider this point to have any etiological significance.

Köster (55) suggests that the symptoms may be due to a primary implication of the vascular system. On this assumption Lepine has proposed the term *Angiomyositis*. It is interesting to note in this connection that Rosenblatt (56) has described a case resembling dermatomyositis, which showed thrombus formation in the vessels, degenerative changes in the vessel wall with fibrin and leucocytes in and about the vessel wall.

Cold and fatigue are said to play a very minor part, though occasionally, as in our case, they seem to be exciting causes.

Distribution.—Cases have been reported, as has been already noted, in the United States and in a number of European countries.

Race.—The Anglo-Germanic has furnished most of the instances, followed by the Latin and Scandinavian. Our patient is the only case in the Negro yet reported.

Season.—The time of year seems to have no connection with the disease. Ten were attacked in the winter, five in the spring, eight in the summer and five in the autumn.

Sex.—The affection has been observed in seventeen males and eleven females. Its distribution among the sexes is probably about equal.

Age.—The disease is usually met with in middle life, though cases at both extremes are found, *e. g.*, Löwenfeld's and Fuckel's cases.

MORBID ANATOMY.—Of the seventeen fatal cases autopsies have been performed in all but seven, and in four of these the excision of a piece of muscle was allowed. We have consequently a fair amount of data on which to base our conclusion as to the changes found in this disease.

With the exception of an enlarged and soft spleen, the pathological changes are limited to the muscles. Any or all may be attacked. It was early stated that the muscles of the eye, tongue, heart and diaphragm were exempt, but later investigation has shown that the eye muscles were implicated once (Strümpell's case); the tongue three times (Jacoby's, Strümpell's and Köster's cases); the heart apparently twice (Kell's and one of Oppenheim's cases), and the diaphragm five times (Strümpell's, Köster's, Senator's second case, Wagner's second case, and Janowsky's and Wyssokowicz's case). Batten (57) states that the masseters usually escape. They were implicated in Boeck's, Strümpell's and Unverricht's second case.

The skin covering the muscles is firm and hard on palpation and does not pit on pressure, though oedema is usually present, but may be slight, as in Jacoby's case. On section the subcutaneous tissues present a firm, tense oedema and are usually infiltrated with a yellowish serous fluid.

Macroscopically the muscles may exhibit extensive changes, as Unverricht has shown in an illustration of their gross appearance in his first case. The muscles are swollen, pale red, or pale yellow in color, or may reveal occasionally yellowish gray or diffuse reddish streaks. Hepp considers the muscles to resemble those of the dog. They are often strongly infiltrated with serum and quite moist. In consistency they vary, being hard and firm or soft and boggy. They may be quite friable.³ They are without lustre and of a dull opaque appearance. Haemorrhages are occasionally seen in them. Microscopically the changes are those of a parenchymatous and interstitial

³ In Hepp's case the left rectus muscle was found ruptured at the autopsy.

inflammation and may vary in extent, being either focal or diffuse. Again, the different muscle changes may be seen occurring in one and the same fiber. These are frequently separated from one another by the existing oedema or the presence of mononuclear and polymorphonuclear leucocytes. Small haemorrhages may also be seen between them. The fibers themselves are found in all stages of degeneration. They are swollen, coarsely or finely granular, hyaline, or waxy—occasionally fatty. The striae are normal, indistinct, or invisible. Longitudinal or cross cleavage of the fiber has been found, and vacuoles have been described in four instances. In many cases there is an increase in the number of muscle nuclei. Typical interstitial foci of small round cells are found in the perivascular connective tissue and, to a lesser extent, between the muscles. In the subacute and chronic cases the increase in connective tissue may be quite marked in both the perimysium externum and internum, and solitary muscle fibers in the process of degeneration may be seen surrounded on all sides by connective tissue. In five cases the blood-vessels were somewhat dilated and filled with blood.

Wagner has described new muscle formation as taking place in some of the fibers that had undergone waxy degeneration, while Senator has noted an abnormally large number of muscle spindles, two or three being found in each section. Senator, Pfeiffer and Lorenz consider the acute interstitial changes to be primary in this disease, and the degenerative muscle findings to occur chiefly in the second stage. They base this assertion on the reports of the different observers of the pathological processes here met with.

Illustrations of the microscopic changes are given by Wagner (in his second case), Jacoby, Strümpell, and Senator. For the detailed findings described by each writer see the table to be found at the end of this article.

In seven of the autopsies broncho-pneumonia was found as a terminal infection.

SYMPTOMS.—The disease generally attacks persons in the prime of life and in the best of health. Fuckel's case, however, was noted in a

girl after an attack of measles,⁴ and both of Wagner's cases had pulmonary tuberculosis, the second one having tuberculous ulcers of the intestines. Senator's second patient was a diabetic.

The onset is almost always gradual with the prodromal symptoms of malaise, weakness, anorexia, or headache. Vomiting was seen in two cases. These symptoms may be of several days' to three weeks' duration, or even longer, as in one case. Occasionally they are absent.

Pain.—Vague rheumatoid pains are next complained of, as well as a stiffness or rigidity in the extremities and back. These pains quickly take on a more definite character and become localized in the muscles. Different muscle groups are then successively attacked. Eventually the whole skeleton musculature may be implicated. Later in the disease the pains are more severe and prevent the patient from making the slightest movement.

Fever is soon noted. It is usually of moderate intensity, and remittent in character in the later stages of the disease. It rarely exceeds 104° F. Just before death it may rise several degrees.

Oedema.—With the fever oedema appears and may implicate the whole body and extremities, the latter presenting at times a most ungainly appearance. It is generally first seen on the face, especially about the eyelids. After the skin inflammation is noticed, it becomes more intense and may remain localized over the affected muscles or spread to surrounding parts. The wrists and ankle-joints are usually spared.

Dermatitis.—Dermatitis is an early symptom and varies greatly in character, being in different cases an erythema, a pseudo-erysipelas, an urticaria, a roseola, or an inflammation resembling erythema nodosum. It may spread continuously or remain limited to the parts where it is first observed. At times it occurs only later in the disease.

Profuse *perspiration* and an enlarged spleen usually accompany the other symptoms.

⁴ It is interesting to note that Jessen has but recently reported a case of polymyositis following measles (loc. cit.) It is not typical enough, however, to be classed as one of dermatomyositis.

Nervous System.—No disturbances of sensation are, as a rule, met with, and the nerves are not tender on palpation. The knee-jerks and the electrical reactions are usually either normal or diminished. Lewy got a partial reaction of degeneration in his case.

Stomatitis and *angina* are at times seen, either early or late, in the disease. In Lewy's case and in four of Oppenheim's ulceration of the mucous membranes was observed. On this account the latter has coined the word *dermato-mucoso-myositis* to more rightly name this disease.

The *urine* is usually normal but may contain albumin.

The course may be characterized by improvement and relapses in the subacute and chronic cases, and atrophy of the muscles may also be here observed. In four cases pigmentation of the skin was found after the dermatitis had disappeared.⁵

Complications.—As stated above, pulmonary tuberculosis, tuberculous ulcers of the intestines, and diabetes are complications that have been met with. Acute haemorrhagic nephritis was observed in Senator's second case, while Jacoby's patient had acute cirrhosis of the liver. In seven instances broncho-pneumonia was found as a terminal infection.

Prognosis.—As all the muscles in the body may be implicated, including those of respiration and deglutition, death may result from suffocation or broncho-pneumonia. Out of the total number of cases (twenty-eight in all) seventeen terminated fatally, so that the prognosis must always be grave. The outcome was fatal for the two patients in the two extremes of life.

Diagnosis.—As a rule no difficulty will be experienced in typical cases. Diseases presenting somewhat similar symptoms are: 1. Trichinosis; 2. Neuromyositis; 3. Infective myositis; and 4. Syphilitic myositis. In the first the initial gastro-intestinal disturbances and the discovery of trichinae in the stools and excised muscle; in 2 the chain of nervous phenomena and the absence of a dermatitis; in 3

⁵ For detailed account of the special symptoms, one is referred to Lorenz's excellent and exhaustive article or to Oppenheim's last paper.

the presence of a focus of infection with the bacteriological report on the muscle examination; and in 4 the history of the patient and the objective findings will generally sufficiently differentiate the affection. It is also well to bear in mind that one of Oppenheim's cases turned out, four years later, to be a typical example of scleroderma.

Treatment.—This should be chiefly directed to the relief of pain and to the keeping up of the patient's nutrition. Various analgesiacs have been tried for the former object but no particular drug employed has as yet been found to offer any especial advantage over the others. There is no specific treatment.

From the general observations upon dermatomyositis we now proceed to the detailed consideration of our case.

CASE.—John E., aet. 31 years (Medical History No. 9632), laborer, a dark complexioned negro, was admitted to the Johns Hopkins Hospital March 7, 1899, complaining of soreness in his legs, chest, and hands, swelling of his muscles and inability to use them.

Family History.—Unimportant.

Previous History.—The patient had measles and whooping-cough as a child. He also gave a somewhat vague history of malaria four years prior to admission to the hospital. He had had gonorrhoea nine or ten years ago. The attack lasted about seven days; there were no complications. He denied syphilis. He used tobacco sparingly and was a moderate beer drinker.

Work.—Had been working almost constantly since a boy in the stone quarries of Baltimore. His work had been that of a "trencher" and had obliged him to be exposed to all kinds of weather. He had often been drenched through by the snow or rain.

Food.—He had always had a special fondness for beef and pork. Had also eaten a good deal of sausage, bologna and especially blood-pudding. He would often make a meal off sausage which he would buy uncooked to carry with him to work and eat in that condition for his noon-day meal.

Present Illness.—About six weeks before admission he was obliged to stop work because his limbs were sore and swollen. He also suf-

ferred from pain in his chest muscles. The onset of his disease began sometime before this but he did not know the exact date. He first noticed that the sole of his right foot pained him, especially on walking. It seemed also somewhat swollen. From here the pain and swelling travelled up to the calf of that leg, then into the thigh, and finally to the hip. Soon after this—about one week, he thinks—his left leg was similarly affected. From here the pain and swelling went to his chest.

The patient stayed at home for about four weeks, either in bed or sitting up in an easy chair. At the end of this time he went to work again but found he could do nothing, for his arms were very sore and weak. Both arms were then affected like the chest and lower extremities. The pain and swelling began in the right hand and went to the forearm, and arm, successively. The left arm was next similarly affected. He described the pains as a "burning like the toothache," and said they were mostly present when he moved.

For about a year he had been somewhat troubled with shortness of breath. He had some nausea and several attacks of vomiting during the present illness but none at the onset.

Recently he had been voiding large quantities of urine, sometimes being obliged to get up as many as six times at night to urinate. There was no ardor urinæ, but often a sharp pain during micturition and at times a sudden stoppage of the flow.

He had not had any diarrhoea or abdominal pain.

On admission the temperature was 100.2° F., respirations 18, pulse 52.

Physical Examination.—Patient was a stout, well developed man and lay in bed in dorsal decubitus. His expression was dull and listless. Lips and mucous membranes were of good color; the conjunctivæ were muddy, but not jaundiced. Pupils were equal and reacted to light and accommodation. The eye movements were normal and no puffiness was noted about the lids. There was no oedema of the face or body and no implication of the muscles of mastication. No appearance of a dermatitis and no signs of a former luetic erup-

tion. No muscle atrophy detected. No swelling of the nerves. No sensitiveness to pressure of the nerve trunks, and no disturbances of sensation.

Thorax.—The pectoral muscles stood out rather prominently and seemed swollen. Some pain was complained of on palpation. The intercostals were apparently not implicated.

The lungs and heart were practically negative on examination. The pulse was 56 to the minute, regular in force and rhythm, and of good volume and tension. There was slight thickening of the vessel wall.

Abdomen negative. Spleen not palpable.

Extremities.—No oedema noted. Flexion, extension, adduction and abduction slight on account of the pain caused. The calf muscles were held very much contracted, and seemed to be quite swollen. The patient complained of much pain and tenderness in them as well as in the muscles of the thighs, forearms, and arms. In these latter regions the contractures and swellings were not as prominent, except in the triceps. Palpation was unsatisfactory on account of the pain elicited and consequently a rigid examination could not be made. The muscles apparently were hard and firm. When the patient lay quiet and motionless in his bed but slight pain was complained of. Marked *lineæ atrophicae* were seen on the flexor surfaces of both arms, especially about the elbow-joints. The grasps of both hands were feeble and the act was attended with pain. The dynamometer did not register in the right hand and only just registered in the left. The knee-jerks were slightly increased. When the patient endeavored to sit up in bed he held his back and legs very rigid. Great difficulty was experienced when he attempted to button his clothes. His gait was extremely unsteady and walking seemed very painful to him.

There was slight general glandular enlargement.

Blood Examination.—Red blood-corpuscles, 5,788,000; white blood-corpuscles, 5250; haemoglobin, 70 per cent.

The differential count was as follows:

Small mononuclears	18.00
Large mononuclears	9.75
Transitionals	1.25
Polymorphonuclears	64.25
Eosinophiles	6.75

The leucocytes were counted daily for about three weeks and averaged about 8000, the highest count being 12,800 on March 12. The red blood-corpuscles remained about the same as at the first examination but the haemoglobin decreased somewhat in percentage. Differential blood-counts were made from time to time and showed but little change. The last count, made two days before the patient's discharge, was as follows:

Small mononuclears	19.00
Transitionals	} 5.00
Large mononuclears	
Polymorphonuclears	69.00
Eosinophiles	7.00

Shortly after his admission, as the suspicion of his having trichinosis was entertained, a small piece of his left gastrocnemius muscle was excised under cocaine anaesthesia. The examination of this teased in salt solution was negative for trichinae. The individual muscle fibers showed marked degeneration. They were studded with minute granules which varied slightly in size and looked like fatty granules. Many of the fibers had lost their transverse striation. There appeared to be a distinct increase in the number of muscle nuclei. The connective tissue was increased in amount and in places showed well-marked cellular infiltration. The further examination of this muscle will be elsewhere described.

After the patient had been in the hospital about two weeks the pain and muscle tenderness subsided, so that he could be up and about the ward. His gait was, however, extremely unsteady and walking was slightly painful. It was only accomplished when he placed his legs

far apart and frequently he had to use a chair to guide himself with. No Romberg symptom was at any time observed.

On March 18 three small nodules were noted on his hands. They were about the size of a split pea and looked somewhat like acne pustules. They were situated: (1) At the junction of the first and second phalanges of the left thumb; (2) in the mid-metacarpal region of the right thumb; and (3) in the mid-metacarpal region of the right little finger. Pain was complained of in them, especially at night. To the touch they were quite hard and firm. One was excised for microscopic examination as will be later mentioned.

The electrical examination made by Dr. Thomas "was difficult on account of the pain which contractions of the muscles caused. A good contraction could be brought out with moderately strong faradic currents applied to the nerves and by direct stimulation of the muscles. When the muscle was stimulated by the faradic current it remained in a continuous tetanic contraction for several seconds. This was not observed when a moderately strong galvanic current was used. When either pole of the galvanic current was placed over the nerve (the ulnar at elbow) and the current gradually increased, pain was complained of and the muscles were thrown into tetanic contraction. This contraction was not confined to the muscles supplied by the nerve but seemed to be due to voluntary effort on account of the pain. In this case it was most marked in the triceps."

On April 8 he complained of a "feeling as if something was crawling down from the elbows of each arm to the fingers." This was especially noted at night. After about eleven days this symptom disappeared and the patient seemed to have better use of the muscles of his arms. The grasp in both hands then, though stronger, was still quite feeble. The pain in the muscles had entirely gone.

The electrical examination, made a few days later, showed no change except that the muscles were no longer thrown into a tetanic contraction by the faradic current. About this time Dr. Osler noticed a distinct myotonic contraction when the patient grasped a round

* Noted also in the cases of Hepp and Lewy, as well as in Wagner's second case.

object. The object was grasped but could not be dropped and his fingers fully extended. This was more observable in the right than in the left hand.

From then on till his discharge, on May 2, there was nothing special to record save that his strength was returning rapidly.

The urine on admission was pale amber in color, clear, 1023 in specific gravity, acid in reaction, negative for sugar but contained albumin to the amount of 0.15 per cent. There was no apparent sediment. Microscopically a few hyaline and granular casts were found. The urine gradually cleared up and at the last examination, made just before the patient's discharge, there was only a trace of albumin and no casts were seen.

Temperature on admission was 100.2° . It was normal the next day and so continued till March 12, when, without any apparent reason, it rose to 101.8° . It was normal again after three days and remained so till the patient's discharge (May 2, 1899).

His subsequent movements were lost track of for about a year, for, although he promised to come daily to the dispensary for electrical treatment, after one visit he disappeared. He was not known at the address he gave us.

He was admitted again to the Johns Hopkins Hospital on February 10, 1900, complaining of cough, soreness in his arms and weakness (Medical History No. 10,905).

Present Illness.—After his discharge from the hospital he had remained quite well and was working in a stone quarry, hammering stone, until nine weeks previous. The present attack occurred, he thought, because he got overheated on a warm day. Dizziness with impairment of vision were the first symptoms. These symptoms came on suddenly and obliged him to stop work. Since then he had remained at home and had complained of soreness and weakness of the trunk and extremities, especially of the right arm and of the left arm at the shoulder. He attributed the trouble in the arms to the fact that he used these muscles most and believed the muscles most used became sorest and weakest. Swelling of the muscles accompanied

this soreness and weakness. In addition to the above described symptoms he had suffered considerably from muscular cramps⁷—usually in his right arm. The forearm would be flexed on the arm during these attacks and the fingers partly bent and held close to each other laterally. The cramps generally lasted from ten to fifteen minutes but were sometimes of longer duration. He did not know how frequently they occurred, but they were never felt in both legs or both arms simultaneously. Pain, not of the cramp-like character, was also complained of in the muscles, especially of the right arm. It was worse on moving about, but was sometimes present when he was at rest.

Had had some attacks of vomiting, especially after eating meat or a good deal of food during his present illness. About ten days before entrance to the hospital he had a shaking chill.

Physical Examination.—The patient looked lethargic. Pain was complained of when the pectoral muscles were palpated. They felt very soft and flabby, even during strong contraction, as did the muscles elsewhere. Although the muscular development seemed enormous there was marked loss of muscular power. No oedema was noted; the spleen was not palpable. Myotonia was not present.

Measurements of Arms and Legs in Centimeters.

	Right.	Left.			Right.	Left
Forearm	31.5	31.5	Contracted {	Forearm	32.5	32
Arm	34.5	34		Arm	39	37.5
Calf	36.5	34		Calf	39	38
Mid-thigh	57.5	57		Mid-thigh	61	59

Blood Examination.—Red blood-corpuscles, 6,600,000; white blood-corpuscles, 4300; haemoglobin, 80 per cent.

Differential blood-count:

Small mononuclears	32.7
Large mononuclears	7.0
Transitionals	2.0
Polymorphonuclears	53.3
Eosinophiles	5.0

⁷ Hepp also has described this in the later stages of the disease.

Electrical Examination by Dr. Thomas showed that the facial nerve responded to a very slight strength of the faradic current on both sides. Direct stimulation of the muscle caused a tetanic contraction which persisted for several seconds and with stronger currents much pain was elicited. With the galvanic current an ordinary contraction was obtained if the current was weak, but a tetanic contraction if the current was of moderate strength. Direct stimulation of the platysma by the faradic current gave a tetanic contraction which lasted for several seconds after the stimulus was removed. The brachial plexus gave with the faradic a tetanic contraction of the muscles supplied by it. This persisted for several seconds after the current was broken. Direct stimulation (faradic) caused tetanus of the portion of the muscle stimulated and the contraction persisted. On stimulating with the galvanic current the right ulnar nerve at the elbow a sharp, quick contraction was obtained which subsided when the current was broken. Repeated, quick galvanic stimulation did not throw the muscles into tetanus, even with a fairly strong current. Direct stimulation of the right external popliteal nerve caused a contraction of the muscles, particularly of that portion under the electrode. This contraction persisted some time after the current was broken. With the galvanic current, stimulation of this nerve caused a sharp, quick contraction, which did not persist. Direct stimulation of the muscles caused a sharp, quick contraction.

Dr. Thomas also noted that when the patient grasped the hand of the observer but little force was used. When told to let go there was some delay. This was also noticed when he was told to grasp an object, squeeze it, and then let go.

On March 2, pain at the back of the neck was first complained of, and difficulty in turning his head was experienced. This soon vanished, however, and the soreness and weakness gradually subsided. The patient left the hospital in good condition March 21, 1900.

On admission the urine showed conditions similar to those found on his first entrance to the hospital. When he was discharged there was only a trace of albumin and no casts were demonstrated.

His temperature varied between normal and 99°. During his stay

a piece of muscle from the left biceps was excised for microscopic examination.

The patient this time also gave a wrong address and we have been unable to locate him.

MICROSCOPIC EXAMINATION OF EXCISED MUSCLE.—Portions of the muscle first removed were hardened in alcohol, bichloride of mercury, and Zenker's fluid. Celloidin and paraffin were used as imbedding agents. The sections were mostly stained with haematoxylin, eosin, safranin, congo red, picrocarmine, and van Gieson's fluid. The Gram-Weigert stain failed to show the presence of bacteria in a number of sections examined. No animal parasites were found. In every section some muscle fibers presented a peculiar anomaly. As seen in cross section this consisted in a collection of fibrillae cut transversely and encircled by a band of fibrillae cut longitudinally (Plate XXX, Fig. 1). Instead of one collection of fibrillae three or more bundles of them were sometimes so surrounded (Plate XXXI, Fig. 3). The muscle fiber showing this anomaly could be normal in structure but more frequently signs of degeneration were found, affecting first the centrally placed, transversely cut fibrillae. These would stain more intensely with eosin, but still reveal the structures that go to make up the Cohnheim fields. In subsequent stages this was difficult or impossible to make out. Finally, a mass staining intensely with eosin was seen, presenting the characteristics of waxy degeneration (Plate XXXI, Fig. 3). The surrounding or encircling longitudinally cut bundles were rarely concerned (Plate XXXI, Fig. 3), but if implicated the degeneration would begin with the innermost fibrillae and pass thence out to the periphery. These fibers were never attacked until the whole transversely cut bundle had undergone waxy degeneration. Eventually the whole muscle anomaly could occasionally be seen to have suffered this change. In almost every instance the outer muscle fiber appeared contracted and shrunken away from its sarcolemma sheath.

In longitudinal sections similar changes were seen. The first stage in the degeneration seemed to consist in the contraction of the inner or now longitudinal fibrillae, by which the cross striations were ren-

dered more prominent (Plate XXX, Fig. 2). Then cleavage of these fibrillae would be noted and a subsequent waxy condition of them with a loss, eventually, of their cross striation. The whole anomaly would occasionally show a marked curling or twisting, such as is met with in degenerating muscle fibers (Plate XXX, Fig. 2).

The sarcolemma was generally normal in appearance but sometimes showed the peculiar blebbing or vascular condition (Plate XXXI, Fig. 4) which Hoen (58) has described in some degenerated voluntary muscle fibers of the uvula. This blebbing was somewhat rare and its origin and development could not be traced so well as in Hoen's case. The appearances seemed to correspond to his description except that the granular material in the blebs was only seen in a few instances and it could not be determined whether this came from the disintegration of the fibrillae. The nuclei occupied the center of the bleb, and their frequent absence could probably be accounted for by the plane of the section being cut above or below that containing the nucleus. The more advanced the degeneration the larger the bleb, but no completely destroyed fiber was observed.

The muscle nuclei presented, at times, a normal appearance, but frequently they were increased in number, swollen, vesicular in shape and contained one to two nucleoli with chromatin granules. They were distributed over the whole extent of the fiber and were often found between individual fibrillae of fibers that had undergone waxy degeneration. Where this degeneration was extreme the nuclei also showed changes. They would be of no definite shape, swollen or decreased in size, would stain intensely in haematoxylin, and would assume bizarre positions.

As a rule, these muscle anomalies were somewhat below the average size of the muscle fibers. They were round or oval in shape, and could be found singly, when they were generally larger in size, or in groups of six or more (Plate XXXI, Fig. 3). They were scattered over all the sections examined and seemed to bear no special relation to the muscle spindles or normal muscle fibers.

The muscle not taking part in this anomalous condition showed interesting changes. Many of the degenerations which have been

described in muscles could be seen. The fibers were normal in size, or swollen and oedematous, or atrophied. First parenchymatous changes were noted, as a result of which the striations failed to come out distinctly in staining. Then the different gradations of waxy degeneration could be made out, and, finally, a complete homogeneous mass, with no intimate muscle structure discernible, could be seen. This was especially marked at the cut ends of the fibers and was probably here due to surgical trauma, as Erb (59) and Weber (60) have pointed out. Occasionally, by the teasing of a fiber, a gap was observed, due to the retraction of the contractile substance, which showed waxy degeneration. The sarcolemma sheath enclosed this gap and within were found very fine granules which took the eosin stain. Owing to the lack of simultaneous retraction there could be seen, in places, irregular cross bands of muscle, also waxy in character. At times this waxy condition had entirely disappeared and the thickened sarcolemma sheath alone remained, filled with rather fine, eosin-staining granules (Plate XXXI, Fig. 6), or fibers presenting this change in only a few fibrillae could be seen, the remaining part exhibiting waxy degeneration (Plate XXXI, Fig. 6).

Longitudinal cleavage was observed here as in the fibers of the muscle anomaly (Plate XXXII, Fig. 7). Cross cleavage was also made out, but rather infrequently. It could not be determined whether the cleavage took place in the membrane of Krause or the plane of Hensen (Plate XXXII, Fig. 8).

In rare instances the bursting of a sarcolemma sheath and the spreading out of the contained fibrillae was noticed (Plate XXXII, Fig. 7). The sheath at the point of rupture always showed Hoen's vesicular degeneration, which had evidently weakened it and might account for its rupture during life. The phenomenon, however, might be a purely artificial one.

Fatty degeneration of the muscle was not observed in any of the sections examined.

Vacuolic degeneration was rare. In all the muscles so affected a swelling of the fiber was seen and, at times, a slight loss of the cross striation. The vacuoles were generally round or oval in shape, of

varying sizes, and so situated that their long axes were parallel to the long axes of the fibers. Their borders were sharp and well defined in every case and their contents consisted of a coarsely or finely granular material containing occasionally two or three nuclei (Plate XXXI, Fig. 5). At times the vacuoles occupied the periphery of the fiber, and then, as a rule, contained a nucleus. Vacuoles occasionally were present in such numbers that the fiber, on cross section, appeared riddled with them. As Schaefer (61) has described, their appearance seemed to be heralded by a round or oval, opaque, non-staining area in a muscle fiber.

In one place there was a considerable extravasation of blood between the fibers. The sarcolemma sheath showed the vesicular degeneration, as above described, rather infrequently. Occasionally the sheath was considerably thickened, especially if its fiber had been completely destroyed.

The nuclei in an unchanged fiber were normal, but with commencing degeneration they would greatly increase in numbers and occupy the center as well as the periphery of the fiber. By a still further increase in number they sometimes almost hid the structure of the muscle. They were oval or vesicular in shape and contained one to two nucleoli. Their long axes were, as a rule, parallel to those of the fiber. At times the nuclei would undergo degeneration, become shrivelled up, stain intensely and lie at right angles to the long axes of the fiber. They disappeared apparently by karyolysis. In no place where there was this proliferation of nuclei were new muscle fibers to be found, but occasionally distinctly reparative processes were observed. The nuclei would lengthen out and become like a rod (Plate XXXIII, Fig. 10). In these cases two or three nucleoli would be seen in them and indentations would indicate a commencing division. Finally, one of these rods would form fourteen or more nuclei (Plate XXXIII, Fig. 11). The next step was the taking on of protoplasm by these nuclei and their formation into myoblasts or sarcoblasts. They received this protoplasm at the expense of the contractile substance of the muscle fiber and consequently might be in hollows of the fiber or, exhausting the whole contractile

substance, might simply be enclosed by the sarcolemma sheath (Plate XXXIII, Fig. 12). Up to this point amitosis was alone observed, but now the myoblasts sometimes showed exquisite examples of karyokinesis (Plate XXXIII, Fig. 13). The new nuclei so formed did not appear to make new cells, but probably in a further stage would form the sarcolemma nuclei of a new muscle cell.

The formation of spindle-cells from these myoblasts was seen, but no new muscle fibers were made out. This might be due to the fact that the process had not continued long enough. Muscle regeneration by the longitudinal cleavage of the old fibers and the subsequent budding of the same was also not observed, though longitudinal cleavage of a collection of fibrillae from an old fiber was occasionally met with (Plate XXXII, Fig. 8).

The myoblastic cells, at times, showed degenerative processes. Their nuclei would then stain intensely and finally disappear, leaving the cell protoplasm which would then take on an intense eosin stain. By the total destruction of all these cells the sarcolemma sheath might be so pressed together by the surrounding fibers as to make it difficult of recognition.

Muscle spindles were, at times, noticed in the sections, but there was no increase in their frequency or any pathological change discernible.

Nerves.—The nerves seemed normal in structure, save that in some instances there was an increased number of nuclei in the sheath of Schwann.

Connective Tissue.—The increase in small round cells in the perimysium externum and internum was comparatively infrequent. It could be seen especially in the perivascular connective-tissue space of the former. Much more marked in these situations was the increase in connective tissue, which was more noticeable in the perimysium externum. The tissue was oedematous and very loose in structure. Occasionally it was seen taking the place of a completely degenerated muscle fiber, or a collection of fibers (Plate XXXII, Fig. 8). It was especially rich in newly formed blood-vessels, and where it bordered on a degenerated muscle fiber it could be seen sending some shoots

across the fiber, if longitudinally cut, and, by further development of other blood-vessels and loose connective tissue, the position of the fiber might be completely occupied by these structures. The increase in fat cells was, at times, quite marked (Plate XXXII, Fig. 9). Occasionally the connective tissue underwent a waxy or granular change and an eosin-staining, granular mass might be alone discernible. In some places large multinucleated giant cells⁸ were noted in the connective tissue, near degenerated muscle fibers.

The small nodules, referred to in the patient's first history, on sectioning and staining, proved to be small subcutaneous abscesses. There was nothing specially remarkable about them. No bacteria were noted in them by the Gram-Weigert stain.

The muscle excised from the patient's left biceps on his second admission was hardened in alcohol, embedded in celloidin and stained with haematoxylin and eosin. I have been unable to see these sections as they were destroyed during a small-pox epidemic in the ward. Dr. Harris, who was in charge of the patient at that time, has written to me that the muscle anomaly was wanting in these sections, but a distinct myositis was present, which consisted, especially, in a great increase in connective tissue between the fibers.

It seems fair to conclude that the above case was one of dermatomyositis, although it is unfortunate that in the light of our further knowledge we have been unable to question the patient more minutely on some of the symptoms in this disease. The swelling spoken of as present at the onset was probably oedema, which is always associated with dermatomyositis. No oedema was noted on his admission to the hospital. He was not asked as to a skin eruption, but if it consisted of merely an erythema he might easily have overlooked it on account of his color. Further information as to fever and some other points is also lacking. I regret I did not pay any attention to the macroscopic condition of the portion of muscle first removed. Microscopically the findings⁹ are identical with those seen in this affection.

I have been unable to find a satisfactory explanation for the muscle

⁸ Seen also by Wagner in his second case.

⁹ Frohmann describes peculiar findings in his case, but they are too indefinite for us to judge whether or no he observed this anomaly.

anomaly. It has been described in two instances and two explanations have been given for its occurrence. Both, however, seem unlikely and rather fanciful. Bataillon (62) found it in frog larva. He considers it a degenerative process and says it begins by the tearing of certain of the outermost fibrillae of a muscle fiber. These torn fibrillae then circumscribe those centrally located, and are more or less obliquely placed. They run in all directions and are plainly separated from the central bundle. Often they show a more regular arrangement and by cross section may be seen to encircle, as a ring, with longitudinal fibrillae, the transversely cut fiber. The regularity of their cross and radial striations, as well as their staining properties, speaks strongly against this view. J. Schaffer (63), on the other hand, thinks that the muscle is not at all concerned, but that the encircling ring or band belongs to the sarcolemma, which, by a peculiar arrangement, simulates muscle striations. It is generally found in atrophied fibers of which the sarcolemma sheath has ruptured. After the rupture the sheath is retracted and drawn down in folds over the muscle fiber, below the site of rupture. The fiber above this retraction is without its sheath and the fibrillae are here widely spread apart. The successive infoldings of the retracted sarcolemma sheath causes the appearance of the circular striations. Our case, in Figs. 1 and 3, seems to refute this theory, as the ring or band can be plainly seen shrunk away from the surrounding sarcolemma sheath. (Plate XXX, Fig. 2, appears to show that this ring or band is muscle, cut in cross section.) The radial striations may possibly be due, Schaffer thinks, to the effect of the acids used in Flemming's solution—the hardening agent employed. Some of the tissue was hardened in Müller's fluid and similar pictures were seen, due, as Schaffer conjectures, to a contraction caused by the embedding process. The muscle in this case was from the gastrocnemius of a man. The description and plates correspond quite closely to the findings in our case, but some minor differences exist. In the lower types of vertebrates and invertebrates a bundle of transversely cut fibrillae is found, in cross sections of muscle, surrounded by a band of longitudinal fibrillae, which are in turn encircled by sarco-

lemma with a layer of undifferentiated sarcoplasm. Kölliker (64), Rollet (65), and others have described such pictures. Our anomaly, consequently, may be looked upon as a structure reverting to that seen in the lower types. Atavism, however, does not explain, but merely places the question in another phase.

In the other instance found in man it was also seen in the gastrocnemius muscle; so it may be a normal structure of that muscle and have some hitherto unknown function. It is our purpose to study further the histological structure of gastrocnemius muscles to see how frequently this so-called anomaly occurs and if it has any function of its own. It is interesting to note that it was not found in Kell's or one of Oppenheim's cases of dermatomyositis, though the gastrocnemius muscle was examined.

It is not inconceivable that what I have called above a muscle anomaly may be merely the result of pathological changes in the muscles. The blebbing of the sarcolemma, if of pathological occurrence, would doubtless weaken the muscle sheath and possibly a rupture might subsequently ensue. The outermost muscle fibrillae of the fiber, then no longer closely confined, would have a tendency to spread out. If consequently a cross section of this fiber were cut at a point where these fibrillae have begun to spread out a picture similar to the so-called anomaly (see Plate XXX, Fig. 1) would be obtained.

I have, also, been unable to explain why degeneration begins at the center of the anomaly and goes outward towards the periphery. In cross sections the longitudinal band seemed to press the transverse fibrillae very closely, and by this constriction the nutrition of the latter might be interfered with and the final result be degeneration of the centrally placed muscle bundle. I merely offer this as a possible partial explanation.

I am indebted to many friends who have looked over my sections and given helpful suggestions; especially do I desire to thank Dr. Welch for his kindly aid and valuable advice, and Dr. Osler for allowing me to report this case.

I am under obligations to Dr. Wolfe, of Hartford, Conn., for the photomicrographs which accompany this article.

TABLE I.—

Reporter.	Year.	Sex.	Age.	Occupation.	Onset.	Fever.	Parts Implicated.
1. E. Wagner.	1863	Woman.	43	Servant.	Swelling of both arms.	Yes.	Neck, thorax, abdomen and arms.
2. Potain.	1875	Man.	17	Farmhand.	Malaise, weakness, pains in extremities, and redness of eye-lids.	Yes.	Pharynx, arms and legs.
3. E. Wagner.	1887	Woman.	34	Cook.	Pain in back and sacrum.	Yes.	Pharynx, neck, thorax, diaphragm, arms and legs.
4. Unverricht and Marchand.	1887	Man.	24	Stone-mason.	Dragging pains in arms and legs.	Yes.	Pharynx, neck, thorax, b'k, abdomen, arms and legs.
5. Hepp and Jackson.	1887	Woman.	36		Malaise, weakness and depression.	Yes.	Pharynx, larynx, neck, thorax, back, abd'n, arms and legs.
6. Jacoby.	1888	Man.	38	Machinist.	Feeling of tension in right calf.	Yes.	Tongue, thorax, back, abdomen, arms and legs.
7. Löwenfeld.	1890	Man.	50	Postman.	Pain in left calf.	Yes.	Pharynx, neck, abdomen, arms and legs.
8. Unverricht.	1891	Woman.	39		Swelling of both legs and urticaria.	Yes.	Face, neck, thorax, back, abdomen, arms and legs.
9. Strömpell.	1891	Man.	70	Gardener.	Nausea, vomiting, headache, and general malaise.	Yes.	Face, eyes, tongue, pharynx, diaphragm, arms and legs.
10. Senator.	1891	Man.	50	Baker.	Pain in right leg.	Yes.	Thorax, arms and legs.
11. Senator.	1891	Man.	40	Restaurant-keeper.	Weakness, anorexia, sleeplessness, dragging pain in back and painful stiffness in limbs.	Yes.	Larynx, diaphragm, abdomen, arms and legs.
12. Boeck.	1892	Man.	21	Laborer.	Violent pains in loins and thighs.	Yes.	Face, neck, thorax, abdomen, arms and legs.
13. Fuckel.	1892	Girl.	8		Diarrhoea and slight fever.	Yes.	Arms and legs.
14. Lewy.	1893	Woman.	25		Malaise, cough and anorexia.	Yes.	Thorax, abdomen, arms and legs.
15. Köster.	1895	Woman.	36		Malaise and weakness.	No.	Face, tongue, pharynx, larynx, neck, thorax, back, abd'n, diaphragm, arms and legs.
16. Kell.	1896	Man.	23	Sailor.	Pain in right calf.	Yes.	Pharynx, larynx, thorax, arms and legs.
17. Frohmann.	1899	Man.	42	Inspector.	Malaise and pain in thighs.	?	Arms and legs.
18. Lepine and Bonnet.	1901	Man.	59	Carpenter.	Intermittent, dull pains in the head.	No.	Face, back and arms.
19. Janowsky and Wyssokowicz.	1901	Woman.	23	Servant.	Pain in arms.	Yes.	Pharynx, larynx, neck, diaphragm, abd'n, arms, legs and sphincter ani muscle.
20. Bacialli.	1902	Man.	39	Clerk.	Vague rheumatoid pains in upper part of body, especially the arms.	Yes.	Pharynx, neck, thorax, arms and legs.
21. Oppenheim.	1903	Man.	38		Malaise, anorexia, chilly sensations.	Yes.	Pharynx, arms and legs.
22. Oppenheim.	1903	Woman.	40		Erythema, fever, pain in neck and arms.	Yes.	Pharynx, neck, arms and legs.
23. Oppenheim.	1903	Woman.	28		Malaise, fever, profuse perspiration.	Yes.	Pharynx, tongue, arms and legs.
24. Oppenheim.	1903	Man.	48	Merchant.	Fever, roseola, pain in extremities.	Yes.	Pharynx, arms and legs.
25. Oppenheim.	1903	Man.	60	Gymnasium instructor.	Angina, pain in extremities, fever.	Yes.	Pharynx, larynx, arms and legs.
26. Christen.	1903	Boy.	10	Student.	Pemphigus-like eruption, vague rheumatoid pains.	Yes.	Face, abdomen, arms and legs.
27. Forcheimer.	1903	Woman.	40		Irritation of skin and mucous membrane, headache, pain in various parts of body.		Thorax and legs.
28. Steiner.	1904	Man.	31	Stone-quarryman.	Pain and swelling in right foot.	Yes.	Thorax, arms and legs.

CASES.

Skin Changes.	Atrophy.	Duration.	Result.	Complications.	Cause of Death.	Remarks.
Erythema.	No.	19 days.	Death.	Pulmonary tuber- culosis.		Autopsy.
Erythema.	No.	6 mos.	Death.	Broncho- pneumonia.	Broncho- pneumonia.	Autopsy.
Erysipelatous erup- tion.	No.	2 mos.	Death.	Pulmonary tuber- culosis.	Broncho- pneumonia.	Autopsy by Huber.
Urticaria.	No.	5 wks.	Death.	Broncho- pneumonia.	Broncho- pneumonia.	Autopsy by Marchand.
Roseola.	No.	11 wks.	Death.	Broncho- pneumonia.	Broncho- pneumonia.	Autopsy by von Recklinghausen.
Erysipelatous erup- tion.	Yes.	2½ yrs.	Death.	Acute cirrhosis of liver. Bronc'p'a.	Broncho- pneumonia.	{ No autopsy. Muscle excised during life.
Roseola. Skin pig- mentation.	Yes.	16 mos.	Death.	Bronchitis. Myo- carditis.		No autopsy.
Urticaria.	Yes.	6 mos.	Recovery.			
Erysipelatous erup- tion.	No.	6 wks.	Death.	Bronchitis.		Autopsy by Hauser.
Eruption like erythe- ma nodosum.	No.	15 days.	Death.	Pneumonia. Dia- betes.	Pneumonia.	{ No autopsy. Muscle excised at death.
Erysipelatous erup- tion.	No.	2 mos.	Recovery.	Bronchitis. Ne- phritis.		
.....	No.	3 mos.	Recovery.	Gonorrhoea.		
Erythema. Urtica- ria.	No.	8 days.	Death.	Bronchitis.		{ No autopsy. Muscle excised at death.
Roseola purpura.	Yes.	2½ mos.	Recovery.	Menorrhagia. Bronchitis.		
Erythema, eczema.	No.	9 mos.	Death.	Broncho- pneumonia.	Broncho- pneumonia.	Autopsy.
Roseola, urticaria.	No.	7 days.	Death.	Bronchitis.	Suffocation.	{ Autopsy. Muscle excised during life.
Erythema.	No.	7 mos.	Death.			Muscle excised during life.
None.	No.	9 mos.	Recovery.			
Erythema. Skin pig- mentation.	No.	5 mos.	Death.	Broncho-pneu- monia. Round duodenal ulcers.	Broncho- pneumonia.	Autopsy by Wyssokowicz.
Papular erythema.	No.	5 mos.	Death.		Suffocation.	Autopsy.
Erythema. Skin pig- mentation.	Yes.	8 mos.	Recovery.			
Erythema.	Yes.	?	?			
Papular eruption.	Yes.	3 mos.	Death.	Epistaxis. Ulcer'n of mucous mem'e.		{ No autopsy. Muscle excised during life.
Roseola. Desquama- tion.	Yes.	3 mos.	Recovery.	Ulceration of mu- cous membrane.		
Erythema.	Yes.	4 mos.	Death.	Myocarditis. Ulc'n of mucous mem'e.		No autopsy.
.....	No.	5 wks.	Recovery.	Epididymitis. Tu- berc's of knee sev'l yr's prev's to at'k.		
Pustular eruption, erythema, urticaria. Skin pigmentation. Desquamation.	Yes.	2½ mos.	Recovery.			
.....	No.		Recovery.			Muscle excised during life.

RÉSUMÉ OF PATHOLOGICAL FINDINGS.

E. WAGNER.—Muscles pale, greyish-red in color. On section the muscles were quite fragile and showed occasionally haemorrhages between the fibers.

Different grades of waxy degeneration to complete destruction of the contractile substance. An extreme grade of fatty metamorphosis. At times both of these changes are seen in the same fibrillae. No vacuoles, but an increase in muscle nuclei. Connective tissue unchanged. Numerous diffuse, very small foci of pus corpuscles. The nerves showed a waxy and fatty degeneration. Muscles of breast, abdomen, neck, upper arm and forearm examined and found implicated.

POTAIN.—Subcutaneous tissues infiltrated with serum.

Muscles pale and permeated with serum.

Many fibers exhibited granular degeneration. The striations were frequently indistinct. Spleen enlarged. Muscle from left forearm examined and found implicated.

E. WAGNER.—Subcutaneous tissues infiltrated with serum.

The muscles mostly of a uniform pale red color. In the flexors of the forearm spots and streaks of very pale appearance alternate with darker areas. The muscles are permeated with serum of a peculiar hard consistency, and are more fragile than normal.

Fibers swollen and showed parenchymatous or waxy degeneration in different stages, as well as fatty changes. The striae faint or invisible. Atrophy of single fibers with simultaneous nuclear increase, or vacuoles without nuclear increase or atrophy. Two or more pathological processes mostly found near each other. Diffuse and circumscribed serous infiltration between the muscles and in the connective tissue. Connective tissue showed proliferated nuclei and single multinucleated giant cells. Regenerative processes of the muscle and connective tissue seen. In places dilatation of capillaries and small haemorrhagic extravasations. Pectoralis major, intercostals, rectus abdominus, diaphragm, triceps brachii, iliacus, gluteus major and soleus examined and found implicated.

UNVERRICHT AND MARCHAND.—Firm, tense oedema of subcutaneous tissues, which are permeated by a yellowish serous fluid.

The muscles are generally of a dark reddish-brown color. They are soft, fragile and without lustre. On cross section pale, slightly transparent, light grey streaks alternate with dark red areas. When one predominates it gives its color to the muscles.

Fibers swollen and showed different stages of a hyaline or waxy degeneration and a finely granular cloudiness. Occasionally fatty degeneration was met with. Longitudinal and cross cleavage. Striae in places completely gone. No vacuoles and no increase in muscle nuclei. Interstitial tissue infiltrated with small round cells. Blood-vessels dilated and filled with blood. Occasionally large or small haemorrhagic extravasations noted. Spleen not enlarged. Extensors of arms (except triceps), extensors of thigh (except sartorius rectus and vasti) and extensors of toes much implicated. Sterno-

cleido-mastoid, pectoralis major and psoas muscles considerably changed. Flexors of thighs and legs as well as muscles of the back only slightly implicated. Abdominal muscles almost unchanged.

HEPP AND JACKSON.—Firm, tense oedema of subcutaneous tissues. Muscles pale and generally yellow in color, though occasional haemorrhagic areas are seen. They resemble the muscles of a dog and are markedly infiltrated with serum.

Different grades of hyaline or waxy degeneration to complete destruction of contractile substance. No fatty or granular degeneration observed. Striae faint or lacking. No vacuoles and no increase in muscle nuclei. Interstitial changes slight, but round-cell infiltration was found in the perivascular connective tissue or near degenerated fibers; in the latter places especially in the empty sarcolemma sheaths. Spleen enlarged. Rectus abdominis, triceps brachii, quadriceps femoris, pharynx and other muscles examined and found implicated.

JACOBY.—Coarsely granular and waxy degenerations in different stages. Fatty degeneration also noted. Longitudinal cleavage. Transverse striae generally lacking. Vacuoles and an increase in muscle nuclei. Increase in connective tissue, which is myxomatous or fibrous in structure. Perimysium externum, at times, exhibited fatty or waxy changes. Some nerves showed inflammatory changes, or an increase of nuclei in the sheath of Schwann. Left supinator longus and right gastrocnemius examined and found implicated.

STRÜMPPELL.—Muscles paler than normal and yellower in color.

Fibers swollen or atrophied and showed a hyaline or waxy degeneration or a finely granular cloudiness. No fatty degeneration noted. Longitudinal and cross cleavage. Transverse striae generally absent. Vacuoles and an increase in muscle nuclei. Occasionally a new formation of connective tissue between the fibers. Typical interstitial foci of small round cells, especially in the perivascular connective tissue. Capillaries dilated and filled with blood. Spleen enlarged. Eye, tongue, pharynx, deltoid muscles of arms, lower extremities and diaphragm examined and found implicated.

SENATOR.—Subcutaneous tissues infiltrated with a sero-purulent fluid.

Muscles paler than normal.

No muscle degeneration. No vacuoles but an increase in muscle nuclei. Connective tissue increased and oedematous. Interstitial changes most marked. Infiltration of round cells (mononuclear and polymorphonuclear leucocytes) seen everywhere, especially in the perivascular connective tissue. Blood-vessels dilated and filled with blood. Small masses of extravasated blood occasionally noted. Great increase in muscle spindles. Right biceps brachii examined and found implicated.

FUCKEL.—Subcutaneous tissues infiltrated with serum.

Muscles somewhat swollen, darker than normal and haemorrhagic. They are quite fragile and lack their normal lustre.

Fibers changed in places to cloudy granules.

Transverse striations partly destroyed.

Muscle from extensors of upper arms examined and found implicated.

KÖSTER.—The muscles were pale, not tense, but quite firm.

Some fibers showed finely or coarsely granular, or hyaline degeneration. No fatty change observed. Longitudinal cleavage. Transverse and longitudinal striae singly or both wanting at times. No vacuoles and no increase in muscle nuclei. Interstitial foci of small round cells especially in the perivascular connective tissue. Capillaries and smaller blood-vessels dilated and filled with blood. Some free red blood-corpuscles in the muscles. Spleen not enlarged. Biceps, pectorals, intercostals, peroneus, tongue, diaphragm, larynx and pharynx muscles examined and found implicated. Pharynx and intercostal muscles most changed.

KELL.—Many fibers swollen and some showed granular and hyaline degeneration. Striae partly or wholly absent. No vacuoles. Between the fibers many large cells with large nuclei (proliferating cells [?]). In places periarteritis.

Spleen enlarged. Supinator longus, pronator, intercostals, heart and gastrocnemius examined and found implicated.

FROHMANN.—Granular degeneration of many fibers and loss of cross striations. Waxy degeneration. Increase in nuclei. Peculiar puckering and folding of the sarcolemma. Slight interstitial changes of an inflammatory nature.

JANOWSKY AND WYSSOKOWICZ.—The muscles are pale and of a yellowish-grey color. They are permeated with serum and of a peculiar hard consistency.

Fibers atrophied and showed a granular and waxy degeneration. Marked increase in connective tissue. There was only a slight cellular infiltration. The nerves showed no inflammatory change. Spleen not enlarged. Muscles of back, upper and lower extremities, throat, diaphragm and sphincter examined and found implicated.

BACIALLI.—Muscles paler than normal.

Fibers swollen or atrophied. They showed different stages of hyaline and waxy degeneration with complete destruction of contractile substance. Longitudinal cleavage. Vacuoles and an increase in muscle nuclei. Marked increase in interstitial connective tissue with abundant infiltration of white blood-corpuscles and here and there small haemorrhagic extravasations. Spleen enlarged.

OPPENHEIM.—Piece of muscle excised from the gastrocnemius showed the characteristic changes of a severe interstitial and parenchymatous myositis.

BIBLIOGRAPHY.

1. Lorenz.—Die Muskelerkrankungen. Wien, 1898, 120.
2. Wagner. E.—Ein Fall von acuter Polymyositis. *Deutsches Archiv f. klin. Med.*, Leipz., 1886-7, xl, 241.
3. Unverricht.—Ueber eine eigenthümliche Form von acuten Muskelentzündung mit einem der Trichinose ahnelnden des Krankheitsbilde. *München. med. Wchnschr.*, 1887, xxxiv, 488. Also, *Cor.-bl. d. allg. ärztl. Ver. v. Thüringen*, Weimar, 1887, xvi, 207. Polymyositis acuta progressiva. *Ztschr. f. klin. Med.*, 1887, xii, 533.
4. Hepp.—Ueber Pseudo-trichinose, eine besondere Form von acuter parenchymatöser Polymyositis. *Berl. klin. Wchnschr.*, 1887, xxiv, 297 and 322.
5. Marchand.—*Jahresbericht der schlesischen Gesellschaft f. vaterländ. Cultur*, 1880, lviii, 65. Also, *Breslauer ärztl. Ztschr.*, 1880, ii, 246.
6. Herrick.—Polymyositis acuta, with report of a case presumably of syphilitic origin. *Am. J. Med. Sc.*, 1896, cxi, 414.
7. Jacoby.—Subacute progressive Polymyositis. *J. Nerv. and Ment. Dis.*, 1888, xv, 697. Also, Ueber primäre subacute progressive Muskelentzündung. *Med. Monatschr.*, 1889, i, 10. And Schmidt's *Jahrb.*, 1889, ccxxi, 248.
8. Lewy.—Zur Lehre von der primären acuten Polymyositis. *Berl. klin. Wchnschr.*, 1893, xxx, 475.
9. Jackson.—Myositis universalis acuta infectiosa, with a case. *Boston Med. and Surg. J.*, 1887, cxvi, 498.
10. Wagner, E.—Fall einer seltenen Muskelkrankheit. *Archiv d. Heilk.*, 1863, iv, 282.
11. Potain.—Morve chronique de forme anormale. *Bull. de la Soc. méd. des hôp. de Paris*, 1875, xii, 314. Also, *Union méd.*, Paris, 1876, xxi, 269.
12. Jacoby.—Op. cit.
13. Löwenfeld.—Ueber einen Fall von Polymyositis acuta. *München. med. Wchnschr.*, 1890, xxxvii, 531 and 549.
14. Unverricht.—Dermatomyositis acuta. *Deutsche med. Wchnschr.*, 1891, xvii, 41.
15. Boeck.—Et Tilfælde af Polymyositis acuta med Udgang i Helbredelse. *Norsk Mag. f. Lægevidensk.*, Christiania, 1891, 4 R., vii, 931. Also, *Centralbl. f. klin. Med.*, 1892, xiii, 405.
16. Strümpell.—Zur Kenntniss der primären acuten Polymyositis. *Deutsche Ztschr. f. Nervenhe.*, 1891, i, 479.
17. Pfeiffer, R.—Die Polymyositis resp. Dermatomyositis acuta. *Centralbl. f. allg. Path. u. path. Anat.*, 1896, vii, 81.

18. Schnell.—Ueber Polymyositis. Dissertation, Würzburg, 1892.
19. Lewy.—Op. cit. and Ibid., 420 and 449.
20. Fraenkel.—Ueber eigenartig verlaufene septicopyämische Erkrankungen. *Deutsche med. Wchnschr.*, 1894, xx, 193, 227 and 245.
21. Buss.—Ein Fall von acuter Dermatomyositis. *Deutsche med. Wchnschr.*, 1894, xx, 788.
22. Herz.—Ueber gutartige Fälle von Dermatomyositis. *Deutsche med. Wchnschr.*, 1894, xx, 790.
23. Albu.—Ueber die Autointoxicationen des Intestinal tractus. Bonn, 1895, 97.
24. Schultze.—Ein Fall von Dermatomyositis chronica mit Ausgang in Muskelatrophie bei einem dreijährigen Knaben. *Deutsche Ztschr. f. Nervenhe.*, 1895, vi, 245. Also, Ein Fall von Dermatomyositis chronica. Dissertation by George Müller, Bonn, 1894.
25. Schultzen.—Ueber Polymyositis, beziehungsweise Dermatomyositis acuta. Festschrift zur hundertjähr. Stiftungsfeier des med.-chir. Friedrich Wilhelm Institute, Berlin, 1895, 311.¹⁰
26. v. Kornilow.—Polymyositis primaria acuta. *Deutsche Ztschr. f. Nervenhe.*, 1896, ix, 119.
27. Jollasse.—Ueber acute primäre Polymyositis. *Mitt. a. d. Hamb. Staatskrankenanst.*, 1897, i, 326.
28. Thiele.—Zur Aetiologie der Polymyositis acuta. *Wien. klin. Rundschau*, 1899, xxxiv, 573.
29. Neubauer.—Ein Fall von akuter Dermatomyositis. *Centralbl. f. innere Med.*, 1899, xii, 289.
30. Fuckel.—Ein Fall von Dermatomyositis acuta. *Cor.-bl. d. allg. ärztl. Ver. v. Thüringen*, 1892, 74.
31. Senator.—Ueber acute Polymyositis und Neuromyositis. *Deutsche med. Wchnschr.*, 1893, xix, 933.
32. Lewy.—Op. cit.
33. Köster.—Om primär Dermatomyositis acuta et chronica. *Nord. med. Ark.*, 1896, N. F. vi, No. xviii, 1. Also, *Centralbl. f. innere Med.*, 1897, xviii, 606.
34. Kell.—Report of three cases of polymyositis acuta; with report of post-mortem finding in one case. *J. Am. Med. Asso.*, 1896, xxvi, 967.
35. Gowers.—A lecture on Polymyositis. *Brit. Med. J.*, 1899, i, 65.

¹⁰ I was unable to find this reference in the Surgeon General's Library. Lorenz speaks of the case as atypical as also does R. Pfeiffer, who gives a short resume of it. I have so classed it after reading what Pfeiffer says of it.

36. Frohmann.—Ueber primäre Polymyositis. *Deutsche med. Wchnschr.*, 1899, xxx, Vereins Beilage, No. 4, 24.
37. Bonnet.—Dermatomyosite. *Lyon Médical*, 1901, xxxiii, 10.
38. Lepine.—Polymyosite. *Revue de Méd.*, 1901, xxi, 426.
39. Janowsky and Wyssokowicz.—Ein Fall von Dermatomyositis. *Deutsches Archiv f. klin. Med.*, 1901, lxxi, 491.
40. Bacialli.—Contribuzione allo studio della patogenesi e delle alterazioni istologiche del tessuto muscolare nella polimiosite acuta primaria. *Il Policlinico*, 1902, ix, 16.
51. Oppenheim.—Ueber die Polymyositis. *Berl. klin. Wchnschr.*, 1903, xl, 381 and 416. Also, Zur Dermatomyositis. *Berl. klin. Wchnschr.*, 1899, xxxvi, 805.
42. Christen.—Ein Fall von Polymyositis acuta. *Cor.-bl. f. Schweiz. Aertzte*, 1903, xxiii, 822.
43. Forcheimer.—Dermato-myositis. *Boston Med. and Surg. J.*, 1903, cxlviii, 631.
44. Vincent.—Polymyosite infectieuse aiguë à récidives. *Bull. et mem. Soc. mèd. des hôp. de Paris*, 1903, 3 s., xx, 633.
45. Schüller.—Polymyositis im Kindesalter. *Jahrb. f. Kinderh.*, 1903, lviii, 193.
46. Jessen.—Polymyositis bei Masern. *Berl. klin. Wchnschr.*, 1904, xli, 847.
47. Lorenz.—Op. cit., and Ibid., 160.
48. Pfeiffer, R.—Op. cit.
49. Trinkler.—Kkazuist polymyositis acuta. *Chir. Vestnik.*, 1893, ix, 77.
50. Bauer.—Ein Fall von acuter hämorrhagischer Polymyositis. *Deutsches Archiv f. klin. Med.*, 1899, lxvi, 95.
51. Georgieoski.—A Case of Acute Haemorrhagic Polymyositis. *Bolnitsch. Gaz. Botkina*, 1900, xii, 696. See the reference in Thayer's article entitled, Notes on a Case of Acute Haemorrhagic Polymyositis. *Boston Med. and Surg. J.*, 1902, cxlvii, 313.
52. Pfeiffer, L.—Die Unverricht'sche Polymyositis gregarinosa oder Dermatomyocosis acuta. Die Protozoën als Krankheitserreger, Jena, 1891, 2 Aufl., 208.
53. Virchow.—Giebt es eine Psorospermien-Krankheit bei Schweinen? *Virchow's Archiv*, 1866, xxxvii, 255.
54. Klebs.—Eine neue Form der primären Muskelatrophie (Atrophia muscularis parasitaria symmetrica). Festschrift Rudolf Virchow zu seinem 71. Geburtstage, Berlin, 1891, 31.

55. Köster.—Zur Kenntniss der Dermatomyositis. *Deutsche Ztschr. f. Nervenhe.*, 1898, xii, 150.
56. Rosenblatt.—Ein seltener Fall von Erkrankung der kleinen Arterien der Muskeln und Nerven, der klinisch als Dermatomyositis imponirte. *Ztschr. f. klin. Med.*, 1897, xxiii, 547.
57. Batten.—Myositis. *Allbutt's System of Medicine*, 1899, vii, 461.
58. Hoen.—On a Form of Degeneration of Striated Muscle Met with in the Uvula. *The Journ. of Exp. Med.*, 1898, iii, 549.
59. Erb.—Bemerkungen über die sogenannte wachsartige Degeneration der quergestreiften Muskelfasern. *Virchow's Archiv*, 1868, xliii, 108.
60. Weber.—Ueber die Neubildung quergestreiften Muskelfasern, insbesondere die regenerative Neubildung derselben nach Verletzungen. *Virchow's Archiv*, 1867, xxxix, 216.
61. Schaefer.—Ueber die histologischen Veränderungen der quergestreiften Muskelgestreiften in der Peripherie von Geschwülsten. *Virchow's Archiv*, 1887, cx, 475.
62. Bataillon.—Recherches anatomiques et expérimentales sur la métamorphose des amphibiens anoures. *Ann. de l'université de Lyon*, 1891, ii, Fasc. 1.
63. Schaffer, J.—Beiträge zur Histologie und Histogenese der quergestreiften Muskelfasern des Menschen und einiger Wirbelthiere. *Sitzungsb. d. k. Akad. d. Wissensch., Math.-naturw. Cl.*, 1893, cii, 64.
64. Kölliker.—Handbuch der Gewebelehre des Menschen, Leipzig, 1889, i, 358.
65. Rollet.—Ueber die Flossen muskeln des Seepferdchens (*Hippocampus antiquorum*) und über Muskelstructure im Allgemeinen. *Archiv f. Mikr. Anat.*, 1888, xxxii, 233.

DESCRIPTION OF PLATES.

All of the illustrations are photo-micrographs. The objectives and oculars were the Zeiss apochromatic; the condenser was Bauch and Lomb's, with a numerical aperture of 1; the illumination was the electric arc; Zetnow's light filter with methylene blue was used. The plates were Cramer's isochromatic (medium) and the exposure was two and a half seconds.

PLATE XXX.

Fig. 1. $\times 656$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

The muscle anomaly is here well shown in cross section, in the central part of the field. The innermost fibrillae are cut transversely and are surrounded by a longitudinal band of fibrillae, which is somewhat shrunken away from the sarcolemma. The cross striation of the longitudinal band is quite prominent. The anomaly is about equal in size to the other fibers.

Fig. 2. $\times 570$ and reduced. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

The muscle anomaly represented cut longitudinally. The innermost fiber is contracted and somewhat curled on itself. Its striations are very prominent. To the right are waxy fibrillae, showing longitudinal cleavage and loss of some of the cross striations. The vesicular degeneration of the sarcolemma is to be noted in places.

PLATE XXXI.

Fig. 3. $\times 570$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

Six of these muscle anomalies, about equal in size to the normal fibers, are seen. Some of the fibrillary bundles are stained deeply and have a homogeneous appearance. The larger one in the lower row shows this waxy change especially well and the innermost border of the longitudinal muscle layer is beginning to be similarly affected. The upper right anomaly shows three fibrillary bundles cut in cross section and surrounded by longitudinal fibrillae. In some the muscle is considerably shrunken away from the sarcolemma.

Fig. 4. $\times 570$ and reduced $\frac{1}{2}$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

The blebbing or vesicular degeneration of the sarcolemma. In a few blebs nuclei are to be seen. There is a great increase in nuclei in the muscle fiber on the left.

Fig. 5. $\times 570$ and reduced $\frac{1}{2}$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

The swollen condition of a muscle fiber filled with vacuoles. The vacuoles are sharply defined, when well focused, and are generally fairly well filled with rather fine granules. A nucleus is to be seen in the innermost vacuole.

Fig. 6. $\times 570$ and reduced $\frac{1}{2}$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

The swollen and degenerated sarcolemma sheath is here represented as quite well filled with rather fine granules. The fiber on the right is swollen and shows waxy degeneration together with this granular condition.

PLATE XXXII.

Fig. 7. $\times 570$ and reduced. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

The bursting of the sarcolemma sheath and the spreading out of the fibrillae. Above and below this spot the muscle fiber is contracted. An increase in muscle nuclei is seen. This bursting of the sarcolemma sheath may be due to its weakening caused by the vesicular degeneration at this place, or it may be an artificial phenomenon.

Fig. 8. $\times 570$ and reduced. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

Increase of connective tissue at expense of the muscle which in places is broken up into bands or shows a waxy condition of the fibrillae. Cross and longitudinal cleavage is here seen. The connective tissue is loose and very oedematous. There is a marked increase in muscle nuclei.

Fig. 9. $\times 156$ and reduced $\frac{1}{2}$. Objective 16 mm. N. A. 30. Compensat. proj. ocular No. 6. Stain, van Gieson.

Increase of the perimysium externum. Many fat cells and new blood-vessels. In the upper part of the field a sarcolemma sheath containing myoblasts is to be noted. The muscle fibers show an increased number of nuclei.

PLATE XXXIII.

Fig. 10. $\times 570$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

Rod-like nuclei in a row of four are to be seen in the center of a muscle fiber.

Fig. 11. $\times 570$. Objective 6 mm. N. A. 95. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

By direct division of these rod-like nuclei long rows of nuclei can be found. In this figure a line of eleven can be seen.

Fig. 12. $\times 156$. Objective 16 mm. N. A. 30. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

These muscle nuclei are here situated in the hollow of a muscle fiber and in an empty sarcolemma sheath and are surrounded by protoplasm. These cells are known as myoblasts or sarcoblasts.

Fig. 13. $\times 1390$. Objective 2 mm. N. A. 1.42. Compensat. proj. ocular No. 6. Stain, haematoxylin-eosin.

One of the myoblasts seen in the last figure is shown under higher amplification undergoing karyokinesis.

FIG. 1.

FIG. 2.

FIG. 3.

FIG. 4.

FIG. 5.

FIG. 6.

FIG. 7.

FIG. 8.

FIG. 9.

FIG. 10.

FIG. 11.

FIG. 12.

FIG. 13.