ABC of clinical haematology

Macrocytic anaemias

Victor Hoffbrand, Drew Provan

Macrocytosis is a rise in the mean cell volume of the red cells above the normal range (in adults 80-95 fl (femtolitres)). It is detected with a blood count, in which the mean cell volume, as well as other red cell indices, is measured. The mean cell volume is lower in children than in adults, with a normal mean of 70 fl at age 1 year, rising by about 1 fl each year until adult volumes are reached at puberty.

The causes of macrocytosis fall into two groups: (a) deficiency of vitamin B₁₂ (cobalamin) or folate (or rarely abnormalities of their metabolism) in which the bone marrow is megaloblastic and (b) other causes, in which the bone marrow is usually normoblastic. In this article the two groups are considered separately, and then the reader is taken through the steps to diagnose the cause of macrocytosis and its management.

Deficiency of vitamin B₁₂ or folate

Vitamin B₁₂ deficiency

The body's requirement for vitamin B₁₂ is about 1 µg daily. This is amply supplied by a normal Western diet (vitamin B₁₂ content 10-30 µg daily) but not by a strict vegan diet, which excludes all animal produce (including milk, eggs, and cheese). Absorption of vitamin B₁₂ is through the ileum, facilitated by intrinsic factor, which is secreted by the parietal cells of the stomach. Absorption is limited to 2-3 µg daily.

In Britain, vitamin B₁₂ deficiency is usually due to pernicious anaemia, which now accounts for up to 80% of all cases of megaloblastic anaemia. The incidence of the disease is 1:10 000 in northern Europe, and the disease occurs in all races. The underlying mechanism is an autoimmune gastritis that results in achlorhydria and the absence of intrinsic factor. The incidence of pernicious anaemia peaks at age 60; the condition has a female:male incidence of 1.6:1.0 and is more common in those with early greying, blue eyes, and blood group A, and in those with a family history of the disease or of diseases that may be associated with it—for example, vitiligo, myxoedema, Hashimoto's disease, Addison's disease of adrenal, and hypoparathyroidism.

Other causes of vitamin B₁₂ deficiency are infrequent in Britain. Veganism is an unusual cause of severe deficiency as most vegetarians and vegans include some vitamin B₁₂ in their diet. Moreover, unlike in pernicious anaemia, the enterohepatic circulation for vitamin B₁₂ is intact in vegans, so vitamin B₁₂ stores are conserved. Gastric resection and intestinal causes of malabsorption of vitamin B₁₂—for example, ileal resection or the intestinal stagnant loop syndrome—are less common now that abdominal tuberculosis is infrequent and H₂-antagonists have been introduced for treating peptic ulceration, thus reducing the need for gastrectomy.

Folate deficiency

The daily requirement for folate is 100-200 µg, and a normal mixed diet contains about 200-300 µg. Natural folates are largely in the polyglutamate form, and these are absorbed through the upper small intestine after deconjugation and conversion to the monoglutamate 5-methyl tetrahydrofolate.

Megaloblastic bone marrow is exemplified by developing red blood cells that are larger than normal, with nuclei more immature than their cytoplasm. The underlying mechanism is defective DNA synthesis

Causes of megaloblastic anaemia

Diet

Vitamin B₁₂ deficiency—Veganism, poor quality diet
Folate deficiency—Poor quality diet, old age, poverty, synthetic diet without added folic acid, goats' milk

Malabsorption

Gastric causes of vitamin B₁₂ deficiency—Pernicious anaemia, congenital intrinsic factor deficiency, gastrectomy
Intestinal causes of vitamin B₁₂ deficiency—Stagnant loop, congenital selective malabsorption, ileal resection
Intestinal causes of folate deficiency—gluten induced enteropathy, tropical sprue

Increased cell turnover

Folate deficiency—Pregnancy, prematurity, chronic haemolytic anaemia (such as sickle cell anaemia), inflammatory and malignant diseases

Renal loss

Folate deficiency—Congestive cardiac failure, dialysis

Drugs

Folate deficiency—Anticonvulsants, sulphasalazine

Defects of vitamin B₁₂ metabolism—for example, transcobalamin II deficiency, nitrous oxide anaesthesia—or of folate metabolism (such as methotrexate treatment), or rare inherited defects of DNA synthesis may all cause megaloblastic anaemia

Other causes of macrocytosis*

- Alcohol
- Liver disease
- Hypothyroidism
- Reticulocytosis
- Aplastic anaemia
- Red cell aplasia
- Myelodysplasia
- Cytotoxic drugs
- Paraproteinaemia (such as myeloma)
- Pregnancy
- Neonatal period

*These are usually associated with a normoblastic marrow

Patient with vitiligo on neck and back.

Body stores are sufficient for only about four months. Folate deficiency may arise because of inadequate dietary intake, malabsorption (especially gluten induced enteropathy), or excessive use as proliferating cells degrade folate. Deficiency in pregnancy may be due partly to inadequate diet, partly to transfer of folate to the fetus, and partly to increased folate degradation.

Consequences of vitamin B₁₂ or folate deficiencies

Megaloblastic anaemia—Clinical features include pallor and jaundice. The onset is gradual, and a severely anaemic patient may present in congestive heart failure or only when an infection supervenes. The blood film shows oval macrocytes and hypersegmented neutrophil nuclei (with six lobes). In severe cases, the white cell count and platelet count also fall (pancytopenia). The bone marrow shows characteristic megaloblastic erythroblasts and giant metamyelocytes (early granulocyte precursors). Biochemically, there is an increase in plasma of unconjugated bilirubin and serum lactic dehydrogenase, with, in severe cases, an absence of haptoglobins and presence in urine of haemosiderin. These changes, including jaundice, are due to increased destruction of red cell precursors in the marrow (ineffective erythropoiesis).

Vitamin B₁₂ neuropathy—A minority of patients with vitamin B₁₂ deficiency develop a neuropathy due to symmetrical damage to the peripheral nerves and posterior and lateral columns of the spinal cord, the legs being more affected than the arms. Psychiatric abnormalities and visual disturbance may also occur. Men are more commonly affected than women. The neuropathy may occur in the absence of anaemia. Psychiatric changes and at most a mild peripheral neuropathy may be ascribed to folate deficiency.

Neural tube defects—Folic acid supplements in pregnancy have been shown to reduce the incidence of neural tube defects (spina bifida, encephalocele and anencephaly) in the fetus and may also reduce the incidence of cleft palate and hare lip. No clear relation exists between the incidence of these defects and folate deficiency in the mother, although the lower the maternal red cell folate (and serum vitamin B₁₂) concentrations even within the normal range, the more likely neural tube defects are to occur in the fetus. One underlying mechanism seems to be a genetic defect in folate metabolism.

Gonadal dysfunction—Deficiency of either vitamin B₁₂ or folate may cause sterility, which is reversible with appropriate vitamin supplementation.

Epithelial cell changes—Glossitis and other epithelial surfaces may show cytological abnormalities.

Cardiovascular disease—Raised serum homocysteine concentrations have been associated with arterial obstruction and venous thrombosis.

Other causes of macrocytosis

The most common cause of macrocytosis in Britain is alcohol. Fairly small quantities of alcohol—for example, two gin and tonics or half a bottle of wine a day—especially in women, may cause a rise of mean cell volume to > 100 fl, typically without anaemia or any detectable change in liver function.

The mechanism for the rise in mean cell volume is uncertain. In liver disease the volume may rise due to excessive lipid deposition on red cell membranes, and the rise is particularly pronounced in liver disease caused by alcohol. A modest rise in mean cell volume is found in severe thyroid deficiency.

Patient with coeliac disease: underweight and low stature.

Blood film in vitamin B₁₂ deficiency showing macrocytic red cells and hypersegmented neutrophil.

Glossitis due to vitamin B₁₂ deficiency.

In other causes of macrocytosis, other haematological abnormalities are usually present—in myelodysplasia (a frequent cause of macrocytosis in elderly people) there are usually quantitative or qualitative changes in the white cells and platelets in the blood. In aplastic anaemia, pancytopenia is present; pure red cell aplasia may also cause macrocytosis. Changes in plasma proteins—presence of a paraprotein (as in myeloma)—may cause a rise in mean cell volume without macrocytes being present in the blood film. Physiological causes of macrocytosis are pregnancy and the neonatal period. Drugs that affect DNA synthesis—for example, hydroxyurea and azathioprine—can cause macrocytosis with or without megaloblastic changes. Finally, a rare, benign familial type of macrocytosis has been described.

Diagnosis

Biochemical assays

The most widely used screening tests for the deficiencies are the serum B₁₂ and folate assays. A low serum concentration implies a deficiency, but a subnormal serum concentration may occur in the absence of pronounced body deficiency—for example, in pregnancy (vitamin B₁₂) and with very recent poor dietary intake (folate).

Red cell folate can also be used to screen for folate deficiency; a low concentration usually implies appreciable depletion of body folate, but the concentration also falls in severe vitamin B₁₂ deficiency, so it is more difficult to interpret the significance of a low red cell than serum folate concentration in patients with megaloblastic anaemia. Moreover, if the patient has received a recent blood transfusion the red cell folate concentration will partly reflect the folate concentration of the transfused red cells.

Specialist investigations

Assays of serum homocysteine (raised in vitamin B₁₂ or folate deficiency) or methylmalonic acid (raised in vitamin B₁₂ deficiency) are used in some specialised laboratories.

Autoantibodies

For patients with vitamin B₁₂ or folate deficiency it is important to establish the underlying cause. In pernicious anaemia, intrinsic factor antibodies are present in plasma in 50% and parietal cell antibodies in 90% of patients.

Other investigations

Radioactive vitamin B₁₂ absorption studies—for example, Schilling test—show impaired absorption of the vitamin in pernicious anaemia; this can be corrected by giving intrinsic factor. In patients with an intestinal lesion, however, absorption of vitamin B₁₂ cannot be corrected with intrinsic factor.

Endoscopy should be performed to confirm atrophic gastritis and exclude gastric carcinoma or gastric polyps, which are two to three times more common in patients with pernicious anaemia than in age and sex matched controls.

A bone marrow examination is usually performed to confirm megaloblastic anaemia. It is also required for the diagnosis of myelodysplasia, aplastic anaemia, myeloma, or other marrow disorders associated with macrocytosis.

If folate deficiency is diagnosed it is important to assess dietary folate intake and to exclude gluten induced enteropathy by endoscopy and duodenal biopsy. The deficiency is common in patients with diseases of increased cell turnover who also have a poor diet.

Investigations which may be needed in patients with macrocytosis

- Serum B₁₂ assay
- Serum and red cell folate assays
- Liver and thyroid function
- Reticulocyte count
- Serum protein electrophoresis
- For vitamin B₁₂ deficiency: serum parietal cell and intrinsic factor antibodies, radioactive vitamin B₁₂ absorption with and without intrinsic factor (Schilling test), possibly serum gastrin concentration
- Consider bone marrow examination for megaloblastic changes suggestive of vitamin B₁₂ or folate deficiency, or alternative diagnoses—for example, myelodysplasia, aplastic anaemia, myeloma
- Endoscopy—gastric biopsy (vitamin B₁₂ deficiency), duodenal biopsy (folate deficiency)

Results of absorption tests of radioactive vitamin B₁₂

	Dose of vitamin B ₁₂ given alone	Dose of vitamin B ₁₂ given with intrinsic factor
Vegan	Normal	Normal
Pernicious anaemia or gastrectomy	Low	Normal
Ileal resection	Low	Low
Intestinal blind-loop syndrome	Low	Low

*Corrected by antibiotics.

Bone marrow appearances in megaloblastic anaemia: developing red cells are larger than normal, with nuclei that are immature relative to their cytoplasm (nuclear:cytoplasmic asynchrony).

Treatment

Vitamin B₁₂ deficiency is treated initially by giving the patient six injections of hydroxocobalamin 1 mg at intervals of about three to four days, followed by four such injections a year for life. For patients undergoing total gastrectomy or ileal resection it is sensible to start the maintenance injections from the time of operation. For vegans, less frequent injections—for example, one or two a year—may be sufficient, and the patient should be advised to eat foods to which vitamin B₁₂ has been added, such as bread.

Folate deficiency is treated with folic acid, usually 5 mg daily orally for four months, which is continued only if the underlying cause cannot be corrected. As prophylaxis against folate deficiency in patients with a severe haemolytic anaemia—such as sickle cell anaemia—5 mg folic acid once weekly is probably sufficient. Vitamin B₁₂ deficiency must be excluded in all patients starting folic acid treatment at these doses as such treatment may correct the anaemia in vitamin B₁₂ deficiency but allow neurological disease to develop.

The illustration of the bone marrow is reproduced with permission from *Clinical Haematology* (AV Hoffbrand, J Pettit), 2nd ed, London: Mosby International, 1994.

Victor Hoffbrand is professor of haematology at the Royal Free Hospital, London.

Preventing folate deficiency in pregnancy

- As prophylaxis against folate deficiency in pregnancy, daily doses of folic acid 400 µg are usual
- Larger doses are not recommended as they could mask megaloblastic anaemia due to vitamin B₁₂ deficiency and thus allow B₁₂ neuropathy to develop
- As neural tube defects occur by the 28th day of pregnancy, it is advisable for a woman's daily folate intake to be increased by 400 µg/day at the time of conception
- Whether this can be achieved by increased consumption of foods with a high folate content—for example, liver, green vegetables, and cereals—or whether women have to take additional folic acid or eat foods deliberately fortified with added folate is the subject of current discussion
- The US Food and Drugs Administration announced in 1996 that specified grain products (including most enriched breads, flours, cornmeal, rice, noodles, and macaroni) will be required to be fortified with folic acid to levels ranging from 0.43 mg to 1.5 mg per pound (453 g) of product
- For mothers who have already had an infant with a neural tube defect, larger doses of folic acid—for example, 5 mg daily—are recommended before and during subsequent pregnancy

The ABC of clinical haematology is edited by Drew Provan, consultant haematologist and honorary senior lecturer at the Southampton University Hospitals NHS Trust, and Andrew Henson, clinical research fellow, university department of primary care, Royal South Hants Hospital, Southampton.

Lesson of the week

Microbial keratitis in intensive care

J F Kirwan, T Potamitis, H El-Kasaby, M W Hope-Ross, G A Sutton

Microbial keratitis is a severe complication of corneal exposure in unconscious patients. We report five cases of microbial keratitis in three patients who sustained visual loss as a result while unconscious in an intensive care unit. The devastating consequences of microbial keratitis continue to be seen despite preventive measures.

Case reports

Case 1

A 74 year old man underwent a coronary artery bypass graft. He subsequently had renal failure and the adult respiratory distress syndrome. He was admitted to intensive care, where he was artificially ventilated for two weeks. After a tracheostomy he contracted a respiratory tract infection with *Pseudomonas aeruginosa* for which no obvious source was found. Twelve days later he developed bilateral microbial keratitis secondary to corneal exposure. This rapidly deteriorated and he developed bilateral endophthalmitis and a corneal perforation in the left eye. He was treated with intravitreal and topical gentamicin, intravenous ceftazidime, and ciprofloxacin. The inflammation gradually resolved. The patient, however, was left with bilateral scarred, thinned, and vascularised corneas. Visual acuity was reduced to light perception only in each eye.

Over the following six months there was little improvement in vision. The main cause of visual loss in the right eye was thought to be the result of corneal opacity and cataract. He underwent a right combined corneal transplantation, cataract extraction, and insertion of an intraocular lens. The procedure was technically difficult because of the caseous nature of the recipient corneal tissue, which tended to disintegrate. Despite some persisting leakage, the graft remained clear with a visual acuity of 6/36.

Case 2

A 31 year old man was involved in a road traffic accident, sustaining multiple trauma, including a right frontotemporal fracture extending into the orbit. He was transferred to a neurosurgical centre and remained unconscious throughout, requiring ventilatory support. After 12 days the opinion of an ophthalmologist was requested because of a right proptosis. He had microbial keratitis, with extensive corneal exposure, and was treated with topical cefuroxime and gentamicin. The following day a temporary tarsorrhaphy was performed. The causative agent was *Acinetobacter calcoaceticus*; this was also the cause of a lower respiratory tract infection that became evident two days later. This agent was resistant to the topical chloramphenicol that had been used for prophylaxis.

Corneal exposure in unconscious patients may lead to devastating ocular consequences without meticulous care and early referral

Royal Eye Unit, Kingston Hospital, Kingston upon Thames, Surrey KT2 7QB

J F Kirwan, ophthalmology registrar

BMJ 1997;314:433-4

continued over

Birmingham and
Midland Eye
Hospital,
Birmingham
B3 2NS

T Potamitis,
registrar
H El-Kasaby,
senior registrar
M W Hope-Ross,
consultant
G A Sutton,
consultant

Correspondence to:
Mr J F Kirwan,
Department of
Ophthalmology,
St George's
Hospital, London
SW17 0QT.

The patient subsequently died of a pulmonary embolus.

Case 3

A 7 year old girl was admitted to hospital with a head injury after a road traffic accident. She sustained a skull fracture and a left third cranial nerve palsy and required assisted ventilation. After four days periorbital swelling and chemosis occurred. The opinion of an ophthalmologist was sought: the patient had corneal exposure, with bilateral microbial keratitis, from which *Pseudomonas aeruginosa* was cultured. This organism was not cultured from any other site. The infection was treated with topical gentamicin, carbenicillin, and later chloramphenicol. Both eyes healed over the next four weeks, leaving bilateral corneal scarring. Two months later visual acuities measured 6/60 in each eye. Over eight months best corrected visual acuity improved to 6/24 in the right eye and 6/6 in the left eye.

Discussion

The eyelids and the tear film are natural barriers to infection. The lids provide a physical barrier to trauma and desiccation, and they deter adherence of organisms to the ocular surface. Tears provide mechanical lubrication to wash away organisms, and they also contain antimicrobial substances. These include immunoglobulins and lysozyme, lactoferrin, ceruloplasmin, and complement components. The cornea is protected by an adherent glycocalyx and a mucin produced by goblet cells. The intact corneal epithelium also acts as a protective barrier.¹

Most corneal infections require predisposing factors such as trauma or impaired host defences, to develop. Most bacteria are unable to penetrate the intact corneal epithelium. If lid closure is ineffective then the tear film is unable to provide coverage and the epithelium becomes susceptible to desiccation. The effect of paralysing and sedating agents means that these normal physiological mechanisms are impaired. Inadequate lid closure may occasionally lead to corneal exposure in otherwise normal patients during sleep.² Normal lid closure is maintained during sleep by the tonic contraction of the orbicularis. Sedation leads to the loss of the blink response to corneal irritation, with impairment of tear film function. Orbital haemorrhage, lid trauma, conjunctival chemosis due to positive pressure ventilation, or a facial nerve palsy may lead to inadequate lid closure, exacerbated particularly by corneal anaesthesia with defective epithelial repair. A few case reports detail corneal complications developing in mechanically ventilated patients. The incidence of corneal exposure and secondary complications is probably low but remains unknown. A survey of patients in intensive care units, using fluorescein enhancement, indicated a very low incidence of corneal exposure (T Potamitis, unpublished data).

The organisms most commonly implicated in mild eye infections are *Staphylococcus aureus*, *Haemophilus influenzae*, and *Streptococcus* spp. An ocular infection rate of 7% has been reported in a paediatric intensive

care unit.³ Several reports have described ocular infections from respiratory tract pathogens.⁴⁻⁵ In two of our cases the patient had a simultaneous infection of the same pathogen in the respiratory tract and eye. Infection may occur during routine bronchial toilet.⁵ Pathogens found in intensive care are often difficult to eradicate, and antibiotic drug resistance is a continuing problem. In two patients the initial antibiotic treatment had to be changed once microbiological data were available.

A survey of intensive care units showed that in most of them eye care was carried out every two hours on ventilated patients and, in other units, review was undertaken at least every six hours. Most units used some method of maintaining eye closure. The most common method was the application of a polyacrylamide gel (Geliperm).⁶ Simple use of this measure alone is not enough to provide corneal protection, and although convenient, this method has no confirmed efficacy. A recent report on patients with microbial keratitis in an intensive care unit showed a reduction in the microbial culture rate and number of cases of microbial keratitis on instigation of a complete programme of maintained eye closure, prophylactic gentamicin, and regular application of lubricating ointment, combined with regular observation. (B T Parkin, A Turner, E P Moore, S D Cook, Oxford ophthalmological congress, 1996.) Monitoring of eye closure needs to be carefully performed as incomplete eye closure may be unrecognised, particularly medially. This may well be most effectively achieved by use of taping. Closure by tarsorrhaphy makes examination difficult, with the risk of a silent infection. Prophylactic use of antibiotic ointment may have something to offer, both by avoiding drying due to exposure, and by preventing secondary infection. Alternatively, lubricating ointment may be combined with a separate topical antibiotic. Widespread antibiotic use, however, encourages the spread of bacterial resistance.

Without meticulous care, corneal exposure and its sequelae, sometimes with devastating consequences, is very likely. We emphasise the need for maintenance of lid closure in at risk patients. Early referral in suspicious circumstances is essential. Fluorescein aided examination may enhance the care of such patients by detecting an epithelial defect before the onset of superimposed corneal infection. In cases with an epithelial defect, particularly with exacerbating risk factors, early lid closure with a lower lid traction suture may be appropriate, as lid closure is maintained, but examination of the eye is still possible.

- 1 Liesegang TJ. Bacterial and fungal keratitis. In: Kaufman HE, Barron BA, McDonald MB, Waltman SR, eds. *The cornea*. London: Churchill Livingstone, 1988: 217-70.
- 2 Katz J, Kaufman HE. Corneal exposure during sleep (nocturnal lagophthalmos). *Arch Ophthalmol* 1977;95:449-53.
- 3 Milliken J, Tait GA, Ford-Jones EL, Mindorff GM, Gold R, Mullins G. Nosocomial infections in a paediatric intensive care unit. *Crit Care Med* 1987;16:233-7.
- 4 Hilton E, Uliss A, Samuels S, Adams AA, Lesser ML, Lowy FD. Nosocomial eye infections in intensive-care units. *Lancet* 1983;i:1318-20.
- 5 Ommeslaag D, Colardyn F, De Laey J. Eye infections caused by respiratory pathogens in mechanically ventilated patients. *Crit Care Med* 1987;15:80-1.
- 6 Farrell M, Wray F. Eye care for ventilated patients. *Intensive Critical Care Nursing* 1993;9:137-41.

(Accepted 20 August 1996)