Recent advances

HIV infection—II

Claire Beiser

Despite recent impressive advances in antiretroviral therapy and the hope they bring for further delaying the onset of AIDS and even for cure of HIV infection, we are left with the need to prevent and treat the opportunistic complications that come with a diagnosis of AIDS. Worldwide, most people infected with HIV do not have access to antiretroviral drugs or to treatment and prophylaxis for many of the complications of HIV infection. In the developing world, therefore, prevention of HIV infection is receiving attention. This article summarises some recent advances in the aetiology, treatment, and prophylaxis of the opportunistic complications of HIV infection as well as progress in HIV prevention in the developing world.

Opportunistic infections

The spectrum of opportunistic infections depends on geography. *Pneumocystis carinii* pneumonia, for example, is common in the developed world, but not in Africa, where tuberculosis is by far the most common opportunistic problem.¹ *Penicillium marneffei* infection has recently emerged as a major opportunistic mycosis in northern Thailand,² and visceral leishmaniasis occurs in several parts of the Mediterranean and Latin America.³ The profile of opportunistic infections has changed since the beginning of the epidemic. In the developed world, survival has improved with the availability of antiretrovirals and prophylactic agents,⁴ and the opportunistic infections that occur with more advanced immunosuppression, such as cytomegalovirus retinitis and *Mycobacterium avium* complex infections have become more common.^{5,6} Furthermore, many opportunistic infections seem to be occurring at lower CD4 counts than earlier in the epidemic,⁷ highlighting again the successes of prophylactic measures and antiretroviral agents.

In 1995, the US Public Health Service published comprehensive guidelines for the prophylaxis of infections in people infected with HIV.⁸ Prophylaxis of *Pneumocystis carinii* pneumonia, tuberculosis, and toxoplasmosis have already been established as important measures that are highly cost effective. The US Public Health Service guidelines strongly recommend prophylaxis of these three infections as the standard of care. In the past two years, however, there have been important advances in the prophylaxis and treatment of two of the most common opportunistic problems, *Mycobacterium avium* complex infections and cytomeg-

Recent advances

- In the past few years impressive advances have been made in the prophylaxis and treatment of some of the most common and serious AIDS related opportunistic infections
- The aetiological agent of Kaposi's sarcoma has been discovered
- There has been a renewed understanding of the need for public health programmes to prevent tuberculosis
- Perhaps most importantly, tremendous successes have been shown with HIV prevention in the developing world: new public health programmes bring hope that far fewer new HIV infections will occur among the less fortunate people of the world

alovirus retinitis. These two infections, which tend to occur at very low CD4 counts, have traditionally been very difficult to treat.

Mycobacterium avium complex infection

Disseminated *Mycobacterium avium* complex infection is an important cause of morbidity and mortality in advanced HIV infection, occurring in as many as half of people who develop AIDS in North America.⁹ Significant progress has been made in the past year toward prophylaxis of this debilitating disease. The US Public Health Service guidelines recommended considering *Mycobacterium avium* complex prophylaxis for all patients with CD4 counts $<75 \times 10^6$ cells/l, and this has been shown to improve survival, leading many clinicians to regard it as a standard of care.

Rifabutin was the first agent shown to be effective at preventing *Mycobacterium avium* complex disease, reducing the incidence by 55%.¹⁰ Four recent trials of the macrolides clarithromycin and azithromycin, either alone or in combination with rifabutin, in patients with CD4 counts $<100 \times 10^6$ cells/l have shown considerable improvements over rifabutin alone in terms of prevention of *Mycobacterium avium* complex disease.¹¹⁻¹⁴ A placebo controlled trial of weekly azithromycin showed a 57% reduction in expected incidence of *Mycobacterium avium* complex bacteraemia,¹¹ and a similar study of clarithromycin showed a 69% reduction.¹² A study comparing rifabutin to weekly

Institute of Human Virology and Division of Infectious Diseases, University of Maryland Medical System, Adult HIV Program, Box 174, Baltimore, MD 21201-1595, USA
Claire Beiser, assistant professor

BMJ 1997;314:579-83

azithromycin and to the combination of rifabutin and azithromycin showed that azithromycin alone decreased the risk of *Mycobacterium avium* complex infection by 50% compared with rifabutin, and the combination lowered the risk by 82% compared with rifabutin.¹³ Preliminary data from a study comparing clarithromycin with rifabutin and with the combination of clarithromycin and rifabutin showed that clarithromycin was superior to rifabutin (40% fewer cases in the clarithromycin group) but the combination was no better than clarithromycin alone.¹⁴ Both clarithromycin and rifabutin improve survival^{12 15}; presumably azithromycin improves survival as well, but the placebo controlled trial of azithromycin was probably not large enough to show a survival benefit. Tolerability of the three regimens using single drugs is similar, but the combination regimens have considerably more side effects. Unfortunately, breakthrough infections on macrolides, particularly clarithromycin, often resulted in *Mycobacterium avium* complex isolates that were resistant to both macrolides—as high as 58% of isolates in the placebo controlled clarithromycin study. Resistance has not been a great problem with rifabutin. Still, the overall high efficacy of macrolides argue strongly for their routine use in preference to rifabutin. Another important consideration in the use of these agents is the potential for drug interactions, particularly with protease inhibitors.¹⁶ Research still needs to be done to enable identification of those who are at higher risk of *Mycobacterium avium* complex infections (as is possible with skin testing for tuberculosis) so that prophylaxis can be reserved for those who really need it.

Progress has also been made in the treatment of *Mycobacterium avium* complex infections. The results of several trials have indicated that the combination of clarithromycin and ethambutol, with or without rifabutin, are the most effective regimens.¹⁷⁻²⁰ Clofazimine should no longer be used because it has been associated with increased mortality, although the reasons for the increased mortality are unclear.²⁰

Cytomegalovirus retinitis

Like *Mycobacterium avium* complex infection, serious opportunistic cytomegalovirus disease is becoming more common, affecting as many as 45% of patients with HIV infection.²¹ Cytomegalovirus retinitis is the most common manifestation of cytomegalovirus disease, which is particularly dreaded by patients because it usually results in blindness and the treatments are cumbersome and expensive. Several new approaches represent significant advances in the treatment of cytomegalovirus retinitis, which usually occurs after the CD4 count drops below 50×10^6 cells/l.

Standard treatment for cytomegalovirus retinitis consists of high doses of either ganciclovir or foscarnet for at least two weeks (induction treatment) followed by lifelong infusions at lower daily doses (maintenance treatment) to prevent relapse. Even when treatment is prescribed appropriately and compliance is perfect, progression of disease is inevitable, occurring about two months after initial treatment with conventional agents.²² Furthermore, chronic indwelling catheters with their attendant risks of infection and other complications bring considerable morbidity and have a negative impact on lifestyle. Alternative approaches,

which include an oral formulation of ganciclovir and methods for delivering treatment locally to the eyes, have recently been developed and subjected to clinical trials. Oral ganciclovir as maintenance treatment was compared with intravenous ganciclovir in three randomised trials.²³⁻²⁵ Oral ganciclovir tended to result in shorter time to progression of retinitis, although this depended on the method of assessment of fundus lesions. A trial of two higher doses of oral ganciclovir did not show a significant improvement in efficacy.²⁶ In terms of overall efficacy, therefore, oral ganciclovir seems inferior to the intravenous form for maintenance treatment, but other factors, such as patients' convenience, also need to be considered.

Cidofovir, a drug recently approved by the US Food and Drug Administration, has a prolonged intracellular half life and thus can be given much less frequently (once weekly infusions for induction and biweekly infusions for maintenance) than either ganciclovir or foscarnet. Its main disadvantage is nephrotoxicity, which makes careful selection and follow up necessary. An unpublished trial comparing cidofovir to deferred treatment of peripheral cytomegalovirus retinitis had a median time to progression of 120 days in the group given cidofovir,²⁷ which compares favourably to ganciclovir or foscarnet.

Dramatic delays in progression have recently been shown with the use of sustained release intraocular implants containing ganciclovir (fig 1).^{29 30} A study comparing the implants to standard intravenous ganciclovir for maintenance treatment showed a median time to progression of 216 days with implants compared with 104 days with intravenous ganciclovir.³⁰ There are two problems with the implants, however. The incidence of retinal detachments after implants was 5% in one study and 18% in another. Furthermore, the implants did not prevent cytomegalovirus disease outside the eyes or in the contralateral eye. A combined approach using implants and oral ganciclovir may be attractive if the rate of detachments becomes acceptably low. A study of this combination is underway. Another approach to local treatment is intravitreal injections. Cidofovir has been shown to control cytomegalovirus retinitis when given as intravitreal injections at intervals of five to six weeks,³¹ but this regimen needs to be compared with other regimens in larger trials.

New options for the treatment of relapsing retinitis have become available recently. Intravenous cidofovir was shown to delay progression considerably in

Fig 1 Intravitreal ganciclovir device in the vitreal cavity. The device is firmly sutured to the incision and is immobile. Reproduced with permission from Sanborn *et al*²⁸

patients who had progressive cytomegalovirus retinitis despite systemic treatment with ganciclovir or foscarnet, or both.³² In a trial of combination ganciclovir and foscarnet compared with either ganciclovir or foscarnet alone for patients with relapsed retinitis, the combination was significantly more effective at delaying progression but not in increasing survival.³³

Given the morbidity of cytomegalovirus retinitis, oral ganciclovir has been studied as a prophylactic regimen. Patients with CD4 counts $<50 \times 10^6$ cells/l or a history of an AIDS defining opportunistic infection and CD4 counts $<100 \times 10^6$ cells/l were randomised to placebo or oral ganciclovir.³⁴ The oral ganciclovir group had a 49% reduction in the incidence of confirmed cytomegalovirus disease. A separate unpublished trial found no overall prophylactic benefit of oral ganciclovir, for reasons that are unclear.³⁵ The entry criteria and the follow up procedures were somewhat different. Also, because of the results of the first study, patients were allowed to switch to open label ganciclovir, potentially reducing the final perceived prophylactic effect of the drug. Because of the unclear benefit combined with the extraordinarily high cost of oral ganciclovir (annual wholesale cost of \$14 000) it is not cost effective, and the US Public Health Service guidelines do not recommend oral ganciclovir as prophylaxis in most patients. An area of active investigation is the use of laboratory techniques, such as quantitative polymerase chain reaction of cytomegalovirus DNA in the blood, to identify people at highest risk of developing cytomegalovirus disease and target those who might benefit from prophylaxis.³⁶ Such techniques might also be helpful in diagnosing cytomegalovirus disease and monitoring the success of treatment.

Tuberculosis

Globally, the most common opportunistic infection and the leading cause of death related to HIV infection is tuberculosis. The two main approaches to tuberculosis control are casefinding combined with treatment of active cases, and prevention of disease in infected people. A recent trial conducted in Zaire compared a six month therapeutic regimen (daily rifampin, isoniazid, pyrazinamide, plus ethambutol for two months followed by twice weekly rifampin plus isoniazid for four months) with a 12 month regimen.³⁷ The rate of relapse was higher after the shorter regimen, but survival was comparable. Other similar trials are underway. A shorter course of treatment, therefore, may be an acceptable, more economic alternative in settings poor in resources. Prophylaxis of active disease with isoniazid for patients infected with HIV and tuberculosis is effective, but widespread implementation of such an approach in developing countries may not be feasible. Trials of prophylaxis of tuberculosis are currently under way in some developing countries.

The United States saw a resurgence of tuberculosis in the late 1980s and early 1990s that highlighted the devastating consequences of cutbacks in the infrastructure of tuberculosis control programmes. Several outbreaks of multidrug resistant tuberculosis occurred, primarily affecting people infected with HIV. In some areas, like New York city, as many as 19% of the cases of tuberculosis were attributable to multidrug resistant tuberculosis.³⁸ Renewed attention to tuberculosis control has begun to deal with the problem. Between

Fig 2 Number of patients with tuberculosis in New York city (solid line) and number receiving directly observed treatment at the end of each year (bars). Reproduced with permission from Frieden *et al*³⁸

1992 and 1994, the number of cases in New York city, which was hit particularly hard, have declined by 21%.³⁹ This improvement was attributed primarily to directly observed treatment (fig 2). Directly observed treatment significantly reduces the rate of primary drug resistance, acquired drug resistance, and relapse.⁴⁰ Furthermore, multidrug resistant tuberculosis can be treated effectively. The most important element to successful treatment is giving at least two drugs to which the organism is susceptible *in vitro*.⁴¹⁻⁴³ In recent years, thanks to several molecular techniques, we have learned that a much higher proportion of cases of tuberculosis (as high as 53% in urban homeless people⁴⁴) are due to recently acquired infection rather than reactivation of latent infection, and that HIV infection increases the risk for disease due to recently acquired infection.⁴⁵

Aetiology of Kaposi's sarcoma

Kaposi's sarcoma (KS) remains an important complication of HIV infection worldwide, although not all groups are affected to the same degree. Recent studies are providing ever more convincing evidence that the cause of Kaposi's sarcoma is a new herpesvirus—Kaposi's sarcoma associated herpesvirus. Viral sequences have been detected in the vast majority of Kaposi's sarcoma tissues tested from patients positive⁴⁶ and negative for HIV,^{47 48} including specimens from Africa. These sequences were detected in only a small percentage of other tissues. Furthermore, the virus was detectable in peripheral blood mononuclear cells of half of patients with Kaposi's sarcoma but in none of the controls tested in one study,⁴⁹ and its detection was correlated with lower CD4 counts. In this same study, people infected with HIV who did not have Kaposi's sarcoma but had detectable Kaposi's sarcoma associated herpesvirus in peripheral blood cells had a significantly higher chance of developing Kaposi's sarcoma than similar patients without detectable virus in blood. This virus is probably sexually transmitted and infects a large proportion of healthy adults: its DNA sequences are commonly found in genital tissues of healthy adults, particularly in semen.⁵⁰ A serological test for antibodies to the virus was recently developed.^{51 52} Antibodies to the virus were correlated with Kaposi's

Fig 3 Reduction of HIV infection over two years by improved management of patients with sexually transmitted diseases in Tanzania. Reproduced with permission from Grosskurth *et al*⁵⁵

sarcoma in people infected with HIV, and in most patients with Kaposi's sarcoma and AIDS these antibodies develop before Kaposi's sarcoma develops.

HIV prevention in developing countries

Prophylaxis of opportunistic infections is generally not feasible in developing countries, and the cost of antiretroviral drugs is prohibitive. Prevention of HIV infection, therefore, has received greater attention. Several recent breakthroughs in HIV prevention in the developing world bring new hope for curtailing the pandemic even in settings poor in resources. In Thailand, for example, a government sponsored programme was begun in the early 1990s to promote safer sexual practices in commercial sex establishments; it resulted in a decrease of nearly 50% in HIV seroprevalence among young Thai men.⁵³ This decline in seroprevalence was accompanied by a considerable increase in reported condom use. This "100% Condom Program" involved the distribution of condoms to brothels and other sex establishments, a media campaign to promote condom use, and an enforcement programme to monitor and encourage use of condoms in brothels. The programme is estimated to have prevented approximately two million HIV infections.⁵⁴

Another approach in the developing world is improved treatment of sexually transmitted diseases, which have been linked with HIV transmission. A randomised community trial in rural Tanzania showed a reduction in the incidence of HIV infection as a result

of improvements in management of patients with sexually transmitted diseases (fig 3).⁵⁵ The intervention, which was fairly modest and replicable in other settings poor in resources, involved training healthcare workers in treating sexually transmitted diseases, ensuring the supply of effective antibiotics for the diseases, and promoting healthcare seeking behaviour. A large trial is underway in Uganda of intensive control versus mass treatment of sexually transmitted diseases. Preliminary results show significant reductions in disease rates.⁵⁶

- Lucas SS, Hounnou A, Peacock C, Beaumel A, Djomand G, N'Gbichi J-M, *et al*. The mortality and pathology of HIV infection in a West African city. *AIDS* 1993;7:1569-79.
- Supparatpinyo K, Khamwan, Baosoung V, Nelson KE, Sirisanthana T. Disseminated *Penicillium marneffei* infection in Southeast Asia. *Lancet* 1994;344:110-3.
- Montalban C, Martinez-Fernandez R, Calleja JL de D, Garcia-Diaz J, Rubio R, Dronda F, *et al*. Visceral leishmaniasis (kala-azar) as an opportunistic infection in patients infected with the human immunodeficiency virus in Spain. *Rev Infect Dis* 1989;11:655-60.
- Enger C, Graham N, Peng Y, Chmiel JS, Kingsley LA, Detels R, *et al*. Survival from early, intermediate, and late stages of HIV infection. *JAMA* 1996;275:1329-34.
- Jones JL, Hanson DL, Ward JW, Kaplan JE. Incidence trends in AIDS-related opportunistic illnesses in injecting drug users and men who have sex with men. XI International Conference on AIDS, Vancouver, 1996. (Abstract WeC2418.)
- Dore G, Kaldor J, Hoy J, Mallal S, Li Y, Mijch M, *et al*. Trends in AIDS-related illness in Australia: the Australian AIDS Cohort. XI International Conference on AIDS, Vancouver, 1996. (Abstract TuB2236.)
- Moore RD, Chaisson RE. Natural history of opportunistic disease in an HIV-infected urban clinical cohort. *Am Intern Med* 1996;124:633-42.
- USPHS/IDSA Prevention of Opportunistic Infections Working Group. USPHS/IDSA guidelines for the prevention of opportunistic infections in persons infected with human immunodeficiency virus: disease-specific recommendations. *Clin Infect Dis* 1995;21(suppl1):S32-43.
- Nightingale SD, Byrd LT, Southern PM, Jockusch JD, Cal SX, Wynne BA. Incidence of *Mycobacterium avium*-intracellular complex bacteremia in human immunodeficiency virus-positive patients. *J Infect Dis* 1992;165:1082-5.
- Nightingale SD, Cameron WD, Gordin FM, Sullam PM, Cohn DL, Chaisson RE, *et al*. Two controlled trials of rifabutin prophylaxis against *Mycobacterium avium* complex infection in AIDS. *N Engl J Med* 1993;329:828-33.
- Oldfield EC, Dickinson G, Chung R, Wallace MR, Craig DB, Fessel WJ, *et al*. Once weekly azithromycin for the prevention of *Mycobacterium avium* complex (MAC) infection in AIDS patients. Third Conference on Retroviruses and Opportunistic Infections, Washington, DC, 1996. (Abstract 203.)
- Pierce M, Crampton S, Henry D, Heifets L, LaMarca A, Montecalva M, *et al*. A randomized trial of clarithromycin as prophylaxis against disseminated *Mycobacterium avium* complex infection in patients with advanced acquired immunodeficiency syndrome. *N Engl J Med* 1996;335:384-91.
- Havir DV, Dubé MP, Sattler FR, Forthal DN, Kemper CA, Dunne MW, *et al*. Prophylaxis against disseminated *Mycobacterium avium* complex with weekly azithromycin, daily rifabutin, or both. *N Engl J Med* 1996;335:392-8.
- Cohn DL, Benson CA, Williams P, Nevin T, Korvick J, Hafner R, *et al*. A prospective, randomized, double-blind, comparative study of the safety and efficacy of clarithromycin vs. rifabutin vs. the combination for the prevention of *Mycobacterium avium* complex (MAC) bacteremia or disseminated MAC disease in HIV-infected patients with CD4 counts < 100 cells/mm³. XI International Conference on AIDS, Vancouver, 1996. (Abstract WeB421.)
- Moore RD, Chaisson RE. Survival analysis of two controlled trials of rifabutin prophylaxis against *Mycobacterium avium* complex in AIDS. *AIDS* 1995;9:1337-42.
- Piscitelli SL, Flexner C, Minor JR, Polis MA, Masur H. Drug interactions in patients infected with human immunodeficiency virus. *Clin Infect Dis* 1996;23:685-93.
- Shafan SD, Singer J, Zarowny DP, Phillips P, Salit I, Walmsley S, *et al*. A comparison of two regimens for the treatment of *Mycobacterium avium* complex bacteremia in AIDS: rifabutin, ethambutol and clarithromycin versus rifampin, ethambutol, clofazimine and ciprofloxacin. *N Engl J Med* 1996;335:377-83.
- May T, Vincent V, Brel F, Beuscart CI, Perronne C, Doco-Lecompte T, *et al*. The French randomized clinical trial of combination therapy with clarithromycin for MAC bacteremia in AIDS patients: final results. XI International Conference on AIDS, Vancouver, 1996. (Abstract WeB422.)
- Dubé M, Sattler F, Torriani F, See D, Havir D, Kemper C, *et al*. Prevention of relapse of MAC bacteremia in AIDS: a randomized study of clarithromycin plus clofazimine, with or without ethambutol. Third Conference on Retroviruses and Opportunistic Infections, Washington, DC, 1996. (Abstract 206.)
- Chaisson RE, Keiser P, Pierce M, Fessel WJ, Ruskin J, Lahart C, Meek K. Controlled trial of clarithromycin/ethambutol with or without clofazimine for *Mycobacterium avium* complex bacteremia in AIDS. Third Con-

- ference on Retroviruses and Opportunistic Infections, Washington, DC, 1996.
- 21 Hoover DR, Saah AJ, Bacellar H, Phair J, Detels R, Anderson R, *et al*. Clinical manifestations of AIDS in the era of pneumocystis prophylaxis. Multicenter AIDS Cohort Study. *N Engl J Med* 1993;329:1922-6.
 - 22 Mortality in patients with the acquired immunodeficiency syndrome treated with either foscarnet or ganciclovir for cytomegalovirus retinitis. Studies of Ocular Complications of AIDS Research Group, in collaboration with the AIDS Clinical Trials Group. *N Engl J Med* 1992;326:213-20.
 - 23 Intravenous versus oral ganciclovir: European/Australian comparative study of efficacy and safety in the prevention of cytomegalovirus retinitis recurrence in patients with AIDS. The Oral Ganciclovir European and Australian Cooperative Study Group. *AIDS* 1995;9:471-7.
 - 24 Drew WL, Ives D, Lalezari JP, Crumpacker C, Follansbee SE, Spector SA, *et al*. For the Syntex Cooperative Oral Ganciclovir Study Group. Oral ganciclovir as maintenance treatment for cytomegalovirus retinitis in patients with AIDS. *N Engl J Med* 1995;333:615-20.
 - 25 Squires KE, Stempien MJ, Shadman A, Buhles WC. Oral ganciclovir versus intravenous ganciclovir maintenance therapy for cytomegalovirus retinitis in patients with AIDS: preliminary results of a phase III study. First National Conference on Human Retroviruses and Related Infections, Washington, DC, 1993. (Abstract 540.)
 - 26 Lalezari J, Friedberg D, Bisset J, Giordano M, Hardy D, Robinson C. A comparison of the safety and efficacy of 3 g, 4.5 g, and 6 g doses of oral ganciclovir versus IV ganciclovir for maintenance treatment of CMV retinitis. XI International Conference on AIDS, Vancouver, 1996. (Abstract ThB305.)
 - 27 Kupperman B, Lalezari J, Stagg R, Kramer F, Holland G, Ives D, *et al*. A phase II/III randomized study of immediate versus deferred intravenous cidofovir for AIDS patients with peripheral CMV retinitis. International Symposium on Experimental and Clinical Ocular Pharmacology and Pharmaceutics, Geneva, 1995.
 - 28 Sanborn GE, Anand R, Torti RE, Nightingale SD, Cal SX, Yates B, *et al*. Sustained-release ganciclovir therapy for treatment of cytomegalovirus retinitis. Use of an intravitreal device. *Arch Ophthalmol* 1992;110:188-95.
 - 29 Martin DF, Parks DJ, Mellow SD, Ferris FL, Walton RC, Remaley NA, *et al*. Treatment of cytomegalovirus retinitis with an intraocular sustained-release ganciclovir implant. A randomized controlled clinical trial. *Arch Ophthalmol* 1994;112:1531-9.
 - 30 Chiron ganciclovir implant study group. A randomized controlled multicenter clinical trial of a sustained-release intraocular ganciclovir implant in AIDS patients with CMV retinitis. 35th Interscience Conference on Antimicrobial Agents and Chemotherapy, San Francisco, 1995. (Abstract I215).
 - 31 Rahhal FM, Arevalo JF, Chavez de la Paz E, Munguia D, Azen SP, Freeman WR. Treatment of cytomegalovirus retinitis with intravitreal cidofovir in patients with AIDS. *Ann Intern Med* 1996;125:98-103.
 - 32 Lalezari JP, Kemper C, Stagg R, Holland G, Ives D, Kramer F, *et al*. A randomized, controlled study of the safety and efficacy of intravenous cidofovir for the treatment of relapsing cytomegalovirus retinitis in patients with AIDS. XI International Conference on AIDS, Vancouver, 1996. (Abstract ThB304.)
 - 33 Studies of Ocular Complication of AIDS Research Group in Collaboration with the AIDS Clinical Trials Group. Combination foscarnet and ganciclovir therapy vs. monotherapy for the treatment of relapsed cytomegalovirus retinitis in patients with AIDS. The Cytomegalovirus Retreatment Trial. *Arch Ophthalmol* 1996;114:23-33.
 - 34 Spector SA, McKinley GF, Lalezari JP, Samo T, Acdruczk R, Follansbee S, *et al*. Oral ganciclovir for the prevention of cytomegalovirus disease in persons with AIDS. *N Engl J Med* 1996;334:1491-7.
 - 35 Brosgart C, Craig C, Hillman D, Louis TA, Alston B, Fisher E, *et al*. Final results from a randomized, placebo-controlled trial of the safety and efficacy of oral ganciclovir for prophylaxis of CMV retinal and gastrointestinal mucosal disease. XI International Conference on AIDS, Vancouver, 1996. (Abstract ThB301.)
 - 36 Advances in the management of AIDS-related cytomegalovirus retinitis. NIH Conference; H Masur, moderator. *Ann Intern Med* 1996;125:126-36.
 - 37 Perriens JH, St. Louis ME, Mukadi YB, Brown C, Prignot J, Pouthier F, *et al*. Pulmonary tuberculosis in HIV-infected patients in Zaire. A controlled trial of treatment for either 6 or 12 months. *N Engl J Med* 1995;332:779-84.
 - 38 Frieden TR, Sterling T, Pablos-Mendez A, Kilburn JO, Cauthen GN, Dooley SW. The emergence of drug-resistant tuberculosis in New York City. *N Engl J Med* 1993;328:521-6.
 - 39 Frieden TR, Fujiwara PI, Washko RM, Hamburg MA. Tuberculosis in New York City—turning the tide. *N Engl J Med* 1995;333:229-33.
 - 40 Weis SE, Slocum PC, Blais FX, King B, Nunn M, Matney B, *et al*. The effect of directly observed therapy on the rates of drug resistance and relapse in tuberculosis. *N Engl J Med* 1994;330:1179-84.
 - 41 Goble M, Iseman MD, Madsen LA, Waite D, Ackerson L, Horsburgh R. Treatment of 171 patients with pulmonary tuberculosis resistant to isoniazid and rifampin. *N Engl J Med* 1993;328:527-32.
 - 42 Turett GS, Telzak EE, Torian LV, Blum S, Alland D, Weisfuse I, *et al*. Improved outcomes for patients with multidrug-resistant tuberculosis. *Clin Infect Dis* 1995;21:1238-44.
 - 43 Salomon N, Perlman DC, Friedmann P, Buchstein S, Kreiswirth B, Milder D. Predictors and outcome of multidrug-resistant tuberculosis. *Clin Infect Dis* 1995;21:1245-52.
 - 44 Barnes PF, El-Haji H, Preston-Martin S, Cave MD, Jones BE, Otaya M, *et al*. Transmission of tuberculosis among the urban homeless. *JAMA* 1996;275:305-7.
 - 45 Alland D, Kalkut GE, Moss AR, McAdam RA, Hahn JA, Bosworth W, *et al*. Transmission of tuberculosis in New York city. An analysis by DNA fingerprinting and conventional epidemiologic methods. *N Engl J Med* 1994;330:1710-6.
 - 46 Moore PS, Chang Y. Detection of herpesvirus-like DNA sequences in Kaposi's sarcoma in patients with and those without HIV infection. *N Engl J Med* 1995;332:1181-5.
 - 47 Chang Y, Cesarman E, Pessin MS, Le F, Culpepper J, Knowles DM, *et al*. Identification of herpesvirus-like DNA sequences in AIDS-associated Kaposi's sarcoma. *Science* 1994;266:1865-9.
 - 48 Chang Y, Ziegler J, Wabinga H, Katangole-Mbidde E, Boshoff C, Schulz T, *et al*. Kaposi's sarcoma-associated herpesvirus and Kaposi's sarcoma in Africa. *Arch Intern Med* 1996;156:202-4.
 - 49 Whitby D, Howard MR, Tenant-Flowers M, Brink NS, Copas A, Boshoff C, *et al*. Detection of Kaposi sarcoma associated herpesvirus in peripheral blood of HIV-infected individuals and progression to Kaposi's sarcoma. *Lancet* 1995;346:799-802.
 - 50 Monini P, de Lellis L, Fabris M, Rigolin F, Cassai E. Kaposi's sarcoma-associated herpesvirus DNA sequences in prostate tissue and human semen. *N Engl J Med* 1996;334:1168-72.
 - 51 Miller G, Rigsby MO, Heston L, Grogan E, Sun R, Metroka C, *et al*. Antibodies to butyrate-inducible antigens of Kaposi's sarcoma-associated herpesvirus in patients with HIV-1 infection. *N Engl J Med* 1996;334:1292-7.
 - 52 Gao S-J, Kingsley L, Hoover DR, Spira TJ, Rinaldo CR, Saah A, *et al*. Seroconversion to antibodies against Kaposi's sarcoma-associated herpesvirus-related latent nuclear antigens before the development of Kaposi's sarcoma. *N Engl J Med* 1996;335:233-41.
 - 53 Nelson KE, Celentano DD, Eiumtrakol S, Hoover DR, Beyrer C, Suprasert S, *et al*. Changes in sexual behavior and decline in HIV infection among young men in Thailand. *N Engl J Med* 1996;335:297-303.
 - 54 Robinson NJ, Silarug N, Surasiengsunk S, Auvert B, Hanenberg R. Two million HIV infections prevented in Thailand: estimate of the impact of increased condom use. XI International Conference on AIDS, Vancouver, 1996. (Abstract MoC904.)
 - 55 Grosskurth H, Mosha F, Todd J, Mwiruabi E, Klokke A, Sendoro, *et al*. Impact of improved treatment of sexually transmitted diseases on HIV infection in rural Tanzania: randomized controlled trial. *Lancet* 1995;346:530-6.
 - 56 Wawer MJ, Sewankambo NK, Gray RH, Serwadda D, Paxton L, Quinn TC, *et al*. Community-based trial of mass STD treatment for HIV control, Rakai, Uganda: preliminary data on STD declines. XI International Conference on AIDS, Vancouver, 1996. (Abstract MoC443.)

Grand Round—University Hospital of Wales, Cardiff

Pyrexia of unknown origin

The difficulty of establishing a diagnosis

Pyrexia of unknown origin is a common problem in medical practice and encompasses a broad spectrum of diagnostic possibilities. The clinical phenomenon of pyrexia of unknown origin was defined in 1961 as an illness persisting for three weeks or longer, with episodes of fever greater than 38.3°C and with no diagnosis reached after one week of hospital investigation.¹ This case highlights the difficulty that

can be encountered in establishing a diagnosis in a patient presenting with pyrexia of unknown origin.

Case history

A 34 year old man developed a mild sore throat on 29 October 1995. This became more severe during the following week, when he consulted his general

Department of Medicine,
University Hospital of Wales, Heath Park, Cardiff CF4 4XW

BMJ 1997;314:583-6

continued over

Case presented by:
 Richard H Evans,
*registrar in medicine
 and infectious diseases*
 Chairman:
 R E Mansel,
professor of surgery
 Discussion group:
 L K Borysiewicz,
professor of medicine
 B D Williams,
*professor of
 rheumatology*

practitioner and was given a seven day course of penicillin V. The patient discontinued the antibiotics after two days because of nausea. During the subsequent fortnight he developed myalgia and arthralgia, followed by drenching night sweats and rigors.

After four weeks he was unable to get out of bed without help because of the severity of the myalgia and arthralgia, and he had lost over 10 kg. He was admitted to his district general hospital in November 1995. He appeared pale and unwell with a fever of 40°C. He complained of weakness and intense myalgia of his shoulder and pelvic girdle muscles, arthralgia of his left knee and right elbow, and left sided pleuritic chest pain. He had no medical history of note and no history of recent foreign travel. Physical examination was unremarkable: there was no palpable lymphadenopathy; no organomegaly; and no stigmata of infective endocarditis.

Laboratory data on admission showed a blood haemoglobin concentration of 81 g/l with normochromic normocytic indices, a raised leucocyte count of $21.85 \times 10^9/l$ with a neutrophil count of $19.65 \times 10^9/l$ and a raised platelet count of $648 \times 10^9/l$. His erythrocyte sedimentation rate and C reactive protein were both raised, at 124 mm in the first hour and 276 mg/l (normal 0-6 mg/l) respectively. Renal function was normal, but liver function tests were raised: aspartate aminotransferase 58 IU/l (5-45 IU/l); alkaline phosphatase 276 IU/l (30-115 IU/l); γ -glutamyl-transferase 174 IU/l (5-48 IU/l); bilirubin 6 μ mol/l (1-17 μ mol/l); albumin 28 g/l (35-50 g/l). Anti-streptolysin O titres were normal. The chest radiograph was normal.

His fevers, myalgia, and arthralgia had not settled after one week in hospital, and he was started empirically on intravenous benzylpenicillin. Four days later he developed a mildly pruritic maculopapular rash over his trunk, arms, and groin. The rash was attributed to the penicillin—which was stopped—but the rash was still present one week later and seemed to be getting worse.

Serological tests failed to show any evidence of a bacterial or viral infection. Blood and urine cultures were negative. Antinuclear antibody, rheumatoid factor, anti-double stranded DNA antibodies, and anti-neutrophil cytoplasmic antibody concentrations were negative. Serum angiotensin converting enzyme concentrations were normal. A bone marrow aspirate showed a hypercellular marrow consistent with reactive change but without any evidence of lymphomatous or other malignant infiltration. The trephine biopsy was normal. A Ziehl-Neelsen stain of the bone marrow was negative for acid-alcohol fast bacilli, and a Mantoux test was negative. He developed a left knee effusion, an aspirate of which showed numerous polymorphs but no organisms or crystals, and culture was negative. Plain radiographs of the affected joints were normal, as were an ultrasound scan of his abdomen, a computed tomogram of his thorax, abdomen, and pelvis, and an isotope lung scan. A transthoracic echocardiogram showed normal heart function with no evidence of infective endocarditis. An indium-111 labelled white cell scan failed to show any focus of infection or inflammation.

Seven weeks after the initial onset of his illness he was transferred to the infectious diseases unit at this hospital for further assessment. He had lost over 22 kg

Differential diagnosis of fever and arthritis²

- Arthritis associated with infectious agents
 - Direct (bacterial, fungal)
 - Indirect (bacterial: acute rheumatic fever, reactive arthritis; viral: hepatitis B)
- Crystal induced conditions
 - Monosodium urate (gout)
 - Calcium pyrophosphate (pseudogout)
- Diffuse connective tissue diseases
 - Systemic lupus erythematosus
 - Mixed connective tissue disease
 - Necrotising vasculitis and other vasculopathies
 - Adult onset Still's disease
 - Rheumatoid arthritis
 - Polymyalgia rheumatica
- Sarcoidosis
- Arthritis associated with chronic inflammatory bowel disease
- Haematological malignancies
- Whipple's disease
- Familial Mediterranean fever

and continued to complain of generalised myalgia and arthralgia of his wrists, left knee, and right shoulder. The left knee effusion was still clinically apparent, and the movement of his right shoulder was restricted. There was evidence of synovitis of both carpometacarpal joints with restriction to extension. The maculopapular rash over the trunk, upper limbs, and groin became more florid during fevers, which were occurring regularly at 10 pm each night and were preceded by intense rigors and sweats. Physical examination was otherwise normal. In addition to the previously documented laboratory abnormalities, he was found to have a greatly raised serum ferritin concentration of 12 000 μ g/l (normal 15-300 μ g/l). Transoesophageal echocardiography, performed to exclude endocarditis, was normal.

Diagnosis

The differential diagnosis of fever and arthritis is wide (box), and most causes had been excluded by the exhaustive investigations. We thought that the pattern of his illness was most suggestive of acute onset connective tissue disease. In this clinical context, and with the benefit of the widespread negative investigations, we were able to secure the diagnosis of adult Still's disease.

Treatment

The patient was treated with pulses of 1 g intravenous methylprednisolone on three consecutive days. His symptoms improved dramatically within 12 hours of starting treatment: his fever settled and never returned (fig 1); the myalgias and joint pains disappeared, and he was able to get out of bed without help. He was subsequently started on oral prednisolone 30 mg and azathioprine 150 mg.

Four weeks after discharge from hospital the patient remained asymptomatic. His erythrocyte sedimentation rate had fallen to 50 mm in the first hour and the serum ferritin concentration to 400 μ g/l. His only complaint was of some persistent restriction in wrist extension.

Comment

The diagnostic spectrum of pyrexia of unknown origin has changed since its original definition over 30 years ago. The incidence of rheumatic fever has declined; infective endocarditis is diagnosed more readily by the routine use of blood cultures and echocardiography; advances in imaging techniques now allow earlier diagnosis of solid tumours; and the general health of the population has improved. As the number of cases of pyrexia of unknown origin due to infectious diseases has declined there has been a corresponding increase in the number of cases due to multisystem diseases—a significant proportion of these patients have adult Still's disease.³⁻⁵

Still's disease, the systemic form of juvenile chronic arthritis, was first described in 1897,⁶ but Still's disease in the adult was not recognised until 1971.⁷ Adult Still's disease characteristically affects young adults, with 75% of patients first presenting between the ages of 16 and 35 years, although several cases have been reported with disease onset after the age of 60 years.^{8,9} Still's disease affects men and women about equally. Although data on incidence and prevalence vary, the disease is not thought to occur in more than 1 case per 100 000 population.¹⁰

The cause of adult Still's disease is unclear. There are no consistent associations with HLA antigens,^{2,11,12} and although numerous infectious agents have been implicated as causative agents¹³⁻¹⁷ the precise role of micro-organisms has yet to be established. It is likely that immune, infectious, and environmental factors may have a role in a genetically predisposed individual.

The diagnosis of adult Still's disease is based entirely on clinical findings (box). Since no reliable serological markers or other diagnostic tests exist, recognition of the classic features of adult Still's disease remains the key to the diagnosis. Fever, the most constant feature of the disease, is high and spiking—the spike typically occurring in the early evening and falling to normal at least once during a 24 hour period. Most patients have pronounced myalgias, often accompanied by polyarthralgias around large joints, commonly the knees, fingers, and wrists. The rash, often said to be the most useful feature for diagnosis, is salmon-pink, macular or maculopapular, and evanescent. It is often pruritic and is frequently misinterpreted as a drug allergy.¹⁹ A sore throat is also said to be characteristic of the disease as no pathogens are isolated, anti-streptolysin O titres remain normal, and because a sore throat is rarely a feature of other

Principal diagnostic criteria for adult Still's disease¹⁸

Major criteria

Fever of 39°C

Arthralgia

Typical rash

Neutrophil leucocytosis

Minor criteria

Sore throat

Lymphadenopathy or splenomegaly, or both

Liver dysfunction

Negative rheumatoid factor, and negative antinuclear antibody

Exclusions

Infections

Malignancies

Rheumatic diseases

Suggested criteria for the diagnosis of adult Still's disease, requires five or more criteria, including two or more major criteria

connective tissue diseases.²⁰ Other common features of the condition include lymphadenopathy, splenomegaly, pericarditis, and pleuritis.^{2,18,21} The early clinical presentation of this condition, however, is extremely variable, making the diagnosis difficult. The classic triad of fever, arthritis, and rash occurs in fewer than 50% of cases at the time of presentation.¹⁸

Laboratory investigations serve to exclude other diagnostic possibilities. Almost all patients with adult Still's disease have a high erythrocyte sedimentation rate and a neutrophil leucocytosis. Many patients have a normochromic normocytic anaemia and abnormal liver function tests. Tests for serum antinuclear antibody concentration and IgM rheumatoid factor are typically negative.^{2,18,21}

Recent studies have shown a correlation between disease activity in adult Still's disease and extremely high concentrations of serum ferritin.²¹⁻²³ Raised serum ferritin concentrations are frequently observed as part of the host response to systemic inflammation. Noticeably raised concentrations are seen in patients with haematological and germ cell malignancies, with acute liver necrosis, and in haemochromatosis, but concentrations rarely exceed 3000 µg/l. In contrast, serum ferritin concentrations above 10 000 µg/l have been noted in patients with adult Still's disease in the active phase of their illness and may therefore be useful not only in the diagnosis of the disease but also in monitoring disease activity.

Aspirin and other non-steroidal anti-inflammatory agents have been used successfully in the treatment of adult Still's disease, though a significant proportion of patients may need either systemic or oral corticosteroid treatment to control their disease. Pulsed intravenous methylprednisolone has been shown to be effective in treating evolving adult Still's disease and, when combined with low dose oral prednisolone, is also effective in preventing short term relapse.^{2,24,25}

Discussion

REM: Were all of these investigations strictly necessary to obtain the diagnosis?

LKB: Yes. The diagnosis is based on the clinical findings alone and relies heavily on exclusion of underlying

Fig 1 Temperature, measured at 8 am and 10 pm, showing high spiking fevers in evening, returning to normal in morning. The fevers settled immediately after first pulse of methylprednisolone

infection or malignancy. Giving high doses of intravenous corticosteroids may prove catastrophic if the diagnosis is not secure. The patient's presentation was incomplete at first, and the investigations performed at the district general hospital were therefore entirely appropriate. Only with the benefit of these negative investigations could we be confident of the diagnosis.

LKB: Is the mechanism of production of these very high serum ferritin concentrations known?

RHE: Moderately raised concentrations probably arise from iron dependent regulation of ferritin synthesis. The greatly raised concentrations in adult Still's disease are likely to be due to a different mechanism: pronounced increases in ferritin synthesis have been observed in human hepatoma cells in response to interleukin 1 β , suggesting direct activation of ferritin mRNA translation by cytokines²²; abnormalities in the glycosylation of ferritin have been shown in some patients with adult Still's disease, a process that may lead to impaired secretion and accumulation in the endoplasmic reticulum.²³ It remains unclear, however, which one (or more) of these mechanisms is responsible for the pronounced hyperferritinaemia seen in adult Still's disease.

LKB: Have any other treatment options been tested?

BDW: The response of patients with adult Still's disease to treatment is difficult to evaluate, partly because the relative rarity of the disease does not lend itself to randomised controlled trials and partly because of the remittent nature of this disorder. High doses of salicylates or other non-steroidal agents have traditionally been used as first line treatment but are effective in only about 20% of patients.¹⁹⁻²⁶ Some of the largest studies have shown that between 50% and 100% of patients with adult Still's disease need corticosteroids to control systemic disease.²⁰⁻²⁷⁻²⁹ Unfortunately, even in those patients who respond to corticosteroids, joint destruction may progress when the systemic symptoms are adequately controlled, and the disease frequently relapses when the dose is reduced. Other agents have been used, but the evidence in favour of these agents is based either on small studies or on single case reports. Several immunosuppressive agents have been used with some success as steroid sparing agents, and remittive agents have been used to control joint symptoms. An initial trial with a non-steroidal agent may be appropriate, but corticosteroid treatment is needed to control the severe systemic symptoms in the most patients. Steroid sparing or remitting agents should be added to the treatment regimen as the steroids are withdrawn.

REM: What is the prognosis for patients with adult Still's disease?

BDW: The early reports of adult Still's disease suggested that it was a relatively benign condition. Recent studies have suggested a more ominous prognosis with recurrence of disease activity being characteristic, and an appreciable number of patients progressing to a chronic disease. Estimates on the proportion of patients who progress to this chronic disease course differ,²⁰⁻²⁷⁻²⁹ but polyarticular onset, root joint involvement at presentation, or the presence of a typical Still's rash have been identified as prognostic markers associated with a chronic disease course and a prolonged time to remission.¹²⁻¹⁹⁻²⁹ Despite the frequently chronic

nature of this illness, disease flares in adult Still's disease are usually milder than the initial episode,¹⁹ and assessments of functional outcome have been quite favourable: about 80% of patients have a functional class of I or II, according to the American Rheumatism Association's classification.¹²⁻²⁹

We are grateful to Dr O M Gibby (consultant physician, Royal Gwent Hospital, Newport) for referring this patient to our unit, and for the patient's permission to publish his case.

The *BMJ* welcomes grand rounds from other hospitals.

- Petersdorf RG, Beeson PB. Fever of unexplained origin: report on 100 cases. *Medicine* 1961;40:1-30.
- Van de Putte LB, Wouters JG. Adult onset Still's disease. In: Sturrock RD, ed. *Rheumatic manifestations of haematological disease. Clinical rheumatology international practice and research*. Vol 5(2). London: Baillière Tindall, 1991:263-75.
- Petersdorf RG. Fever of unknown origin—an old friend revisited. *Arch Intern Med* 1992;152:21-2.
- Larson EB, Featherstone HJ, Petersdorf RG. Fever of undetermined origin: diagnosis and follow-up of 105 cases 1970-1980. *Medicine* 1982;61:269-92.
- Kashiwagi H. Fever of unknown origin: a changing diagnostic spectrum. *Intern Med* 1994;2:65-6.
- Still GF. On a form of chronic joint disease in children. *Med Chir Trans* 1897;80:1-13 (reprinted *Am J Dis Child* 1978;132:195-200).
- Bywaters EG. Still's disease in the adult. *Ann Rheum Dis* 1971;30:121-33.
- Steffe LA, Cooke CL. Still's disease in a 70-year-old woman. *JAMA* 1983;249:2062-3.
- Wouters J, Van Rijswijk M, Van de Putte L. Adult onset Still's disease in the elderly: a report of two cases. *J Rheumatol* 1985;12:791-3.
- Magadur-Joly G, Billaud E, Barrier JH, Pennec YL, Masson C, Renou P, et al. Epidemiology of adult Still's disease: estimate of the incidence by a retrospective study in west France. *Ann Rheum Dis* 1995;54:587-90.
- Miller ML, Aaron S, Jackson J, Fraser P, Cairns L, Hoch S, et al. HLA gene frequencies in children and adults with systemic onset juvenile rheumatoid arthritis. *Arthritis Rheum* 1985;28:146-50.
- Wouters JM, Reekers P, van de Putte LB. Adult-onset Still's disease: disease course and HLA associations. *Arthritis Rheum* 1986;29:415-8.
- Huang SH, De Coteau WE. Adult-onset Still's disease: an unusual presentation of rubella infection. *Can Med Assoc J* 1980;122:1275-6.
- Gordon SC, Lauter CB. Mumps arthritis: unusual presentation as adult Still's disease. *Ann Intern Med* 1982;97:45-7.
- Colebunders R, Stevens WJ, Vanagt E, Snoeck J. Adult Still's disease caused by Yersinia enterocolitica infection. *Arch Intern Med* 1984;144:1880-2.
- Hurst NP, Martynoga AG, Nuki G, Sewell JR, Mitchell A, Hughes GR. Coccidial B infection and arthritis. *BMJ* 1983;286:605.
- Wouters JM, van der Veen J, van de Putte LB, de Rooy DJ. Adult onset Still's disease and viral infections. *Ann Rheum Dis* 1988;47:764-7.
- Yamaguchi M, Ohta A, Tsunematsu T, Kasukawa R, Mizushima Y, Kashiwagi H, et al. Preliminary criteria for classification of adult Still's disease. *J Rheumatol* 1992;19:424-30.
- Pouchot J, Sampalis JS, Beaudet F, Carrette S, Décary F, Salusinsky-Sternbach M, et al. Adult Still's disease: manifestations, disease course, and outcome in 62 patients. *Medicine* 1991;70:118-36.
- Larson EB. Adult Still's disease—evolution of a clinical syndrome and diagnosis, treatment, and follow-up of 17 patients. *Medicine* 1984;63:82-91.
- Ohta A, Yamaguchi M, Kaneoka H, Nagayoshi T, Hiida M. Adult Still's disease: review of 228 cases from the literature. *J Rheumatol* 1987;14:1139-46.
- Schwarz-Eywill M, Heilig B, Bauer H, Breitbart A, Pezzutto A. Evolution of serum ferritin as a marker for adult Still's disease activity. *Ann Rheum Dis* 1992;51:683-5.
- Van Reeth C, Le Moel G, Lasne Y, Revenant M-C, Agnery J, Kahn M-F, et al. Serum ferritin and iso-ferritins are tools for diagnosis of active adult Still's disease. *J Rheumatol* 1994;21:890-5.
- Bisagni-Faure A, Job-Deslandre C, Menkes C. Intravenous methylprednisolone pulse therapy in Still's disease. *J Rheumatol* 1992;19:1487-8.
- Khraishi M, Fam A. Treatment of fulminant adult Still's disease with intravenous pulse methylprednisolone therapy. *J Rheumatol* 1991;18:1088-90.
- Aydintug AO, D'Cruz D, Cervera R, Khamashta MA, Hughes GRV. Low dose methotrexate treatment in adult Still's disease. *J Rheumatol* 1992;19:431-5.
- Wouters JM, van de Putte LB. Adult onset Still's disease; clinical and laboratory features, treatment and progress of 45 cases. *Q J Med* 1986;61:1055-65.
- Bujak JS, Aptekar RG, Decker JL, Wolff SM. Juvenile rheumatoid arthritis presenting in the adult as fever of unknown origin. *Medicine* 1973;52:431-44.
- Cush JJ, Medsger TA, Christy WC, Herbert DC, Cooperstein LA. Adult onset Still's disease: clinical course and outcome. *Arthritis Rheum* 1987;30:186-94.

(Accepted 6 December 1996)