Legal challenge to rationing of haemophilia treatment is lost

Clare Dyer, legal correspondent, *BMJ*

A High Court judge in London last week refused to give the families of four boys with haemophilia the go ahead to challenge a decision by their local health authorities to withdraw treatment with the genetically engineered blood clotting agent recombinant factor VIII.

Duncan Ouseley, QC for the boys, aged between 5 years and 9 years, told Mr Justice Jowitt that South Lancashire, East Lancashire, and Bury and Rochdale Health Authorities had all made "flawed" decisions and failed to consider each individual case on its merits. But the judge said that health authorities had the right to adopt certain policies, and the parents would have an arguable case only if they could establish that their children had special

needs that the authority had failed to consider. The application was "premature" because they had not yet applied for their children to be considered as exceptional.

Mr Ouseley told the judge that the children had been treated with factor VIII derived from human blood until October 1996, when they were switched to the newer recombinant factor VIII, prescribed by their doctors. But the three health authorities had decided that the human blood-derived product was quite safe and the advantages of the newer treatment did not justify the extra cost. The court heard that the genetically engineered product cost each authority an extra £200 000 (\$320 000).

Thousands of patients with

haemophilia were infected with hepatitis C and HIV from human blood-derived factor VIII. This is now heat treated to destroy all viruses, but the parents fear that it could be contaminated by unknown viruses. After the judgment, Ann Alexander, solicitor for one of the families, said: "The judge refused us leave to proceed to judicial review but made it quite clear that he accepts that families can go along to the health authorities and express individual grounds which would justify having this treatment." But the judge warned that it was wrong for parents to think they could "rush off and be treated to their satisfaction as special cases." They should not build up false expectations because it could be

difficult to show that their child was a special case.

Recent guidelines from the UK Haemophilia Centre Directors Organisation favour recombinant factor VIII over the human blood-derived clotting factor (p 749). They state that new patients and those not infected with viruses should be regarded as priority groups for receiving recombinant factor VIII if it has to be phased in on grounds of cost. In Scotland special funds have been set aside for the treatment, but in England and Wales its use is patchy.

Philip Hunt, director of the National Association of Health Authorities and Trusts, said: "This clearly shows the enormous pressures the NHS is under to make decisions about priorities." □

Military medical services are at crisis point

John Warden,
parliamentary correspondent, *BMJ*

The House of Commons Defence Select Committee last week expressed fears for the future of Britain's military medical services.

Its report states that the medical services are not capable of supporting Britain's global military commitment: "A choice would have to be made between sending troops without proper medical support—or only sending the limited number of troops who could be supported."

The prime minister, John Major, responded by maintaining that the defence medical services can provide the necessary support for the international deployment of British troops. He described the problems as

"teething troubles" that the Ministry of Defence recognised and was determined to overcome.

The report echoes the select committee's warning following defence cuts two years ago. Britain's defence medical services have been about halved since 1990, in both beds and staff. At the end of last year there were only 56 surgeons and 48 anaesthetists compared with the 77 and 91 needed respectively. Three military hospitals have been closed and replaced by units within NHS hospitals.

The committee views the staff shortages as so serious that it is not clear whether the defence medical services will recover. Fearing that the military ethos of medicine has been destroyed, the committee is incredulous that the Ministry of Defence has allowed a major and potentially critical staff shortage to develop.

The committee also found that morale at all levels of the defence medical services is lower

Morale is low throughout the defence medical services

than it had ever encountered in the armed forces. It urges the government to implement in full the recommendations of the coming report of the armed forces' pay review body, as well as offering financial incentives to staunch the haemorrhage of doctors and nurses from the defence medical services.

Also last week the Ministry of Defence admitted a "fundamental failure of working practices" that resulted in ministers and parliament being misled over the extent to which organophosphate insecticides were used in the Gulf war. Officials as well as staff in the surgeon-general's department now face disciplinary action. □

In brief

Human haemoglobin from plants: French researchers have inserted the genes for producing human haemoglobin into tobacco plants (*Nature* 1997;386:29-30). The extracted and purified protein product from the transgenic tobacco plants is fully functional. Such plants may in the future provide a cheap and uncontaminated source of various human proteins, the authors say.

Widow allowed to export husband's sperm: The Human Fertilisation and Embryology Authority has decided to allow Diane Blood to export her late husband's sperm to Belgium for her treatment. This follows the judgment by the Court of Appeal on 6 February confirming that treatment in Britain without the donor's written consent would be illegal but that this case was exceptional and no other cases could recur.

Warning over drug interactions and grapefruit juice: The Committee on Safety of Medicines in Britain has warned that cyclosporin, terfenadine and most calcium channel blockers should not be taken with grapefruit juice because of the potential for adverse reactions. Grapefruit juice contains a psoralen that inhibits the metabolism of such drugs.

AIDS deaths down in United States: For the first time the number of deaths among people with AIDS in the United States has declined. A report from the Centers for Disease Control and Prevention shows that deaths declined by 13% during the first six months of 1996, compared with the same period in 1995.

Immunoglobulin benefits patients with multiple sclerosis: A placebo controlled trial of 150 people with relapsing remitting multiple sclerosis found that monthly intravenous immunoglobulin was effective and well tolerated. Among those receiving immunoglobulin, 31% had improved, 53% were stable, and 16% showed worse clinical disability, compared with 14%, 63%, and 23% respectively in the placebo group. (*Lancet* 1997;349:589-93).

US to encourage more black people to join research trials

Norra Macready, *California*

American researchers and government officials met last week to examine ways to encourage more people from ethnic minority groups to take part in research trials. The meeting took place in Tuskegee, Alabama, the site of one of the most infamous examples of racism and paternalism in American medical history, in a bid to emphasise that such abuses will never be repeated.

In 1932 government researchers began studying 400 black men with syphilis in Tuskegee. The purported aim was to study the natural course of syphilis in black men, so the subjects were offered no treatment, even after penicillin became available in 1943. The experiment continued until 1972, when public outrage forced the government to end the study. As a result today, many black Americans are still suspicious of the healthcare system.

"Our intention was to emphasise that African-Americans don't have to be concerned about a Tuskegee-type situation again," said one of the meeting's coordinators, Dr Mack Roach, associate professor of radiology and medical oncology, University of California, San Francisco.

Many black Americans are suspicious of the healthcare system

He said that the medical community has established safeguards such as investigational review boards and human subject experimentation committees. He added: "No African-American physicians participated in the Tuskegee experiment; today there are more African-American physicians in the loop."

Dr Roach said that it is important for African-Americans to enter clinical trials because mortality from cancer is 50% higher in black people than in white people. However, when black and white patients receive similar investigations and treatments their survival rates are also similar.

"There are differences in the incidence of certain diseases. For

example, black men have the highest incidence of prostate cancer," explained Dr Oscar Streeter, assistant professor of radiation oncology and chief physician of radiation oncology, University of Southern California. "But when they are diagnosed and treated promptly, their mortality goes down and they do as well as white men." Overcoming cultural barriers, raising awareness in local communities, and the efforts of black doctors were all mentioned as ways of getting more black people to participate in clinical trials. Dr Streeter said. "We must reassure black Americans that a trial does not automatically mean an experiment. It can also mean care." □

Heroin users wanted for Dutch medical experiment

Tony Sheldon, *Utrecht*

Experienced heroin users are being sought by the Dutch Ministry of Health to test the practical and technical aspects of smoking heroin as part of a medical experiment.

The health minister, Professor Els Borst-Eilers, announced in a recent letter to parliament that up to 10 heroin users are required to smoke a maximum of five tablets in tests from March to June. The tests are part of a preliminary investigation linked to a proposal to prescribe heroin for up to 750 long term addicts that will be decided by

parliament later this year.

Dutch addicts tend to smoke rather than inject heroin, and there is little research in this area. A tablet is being developed that when heated should produce the correct quantity of smokeable heroin without other dangerous materials. This must now be tested along with the means of smoking that allows the maximum amount of heroin to be inhaled without releasing it into the room where it can be breathed in by medical staff. Both the usual method of smoking heroin—using aluminium foil

and a lighter—and the use of a heated block will be tested. The tests must also ensure that a whole tablet is used at one time and that the heroin user is motivated to follow the trial conditions, which include returning to the same room several times a day to smoke the heroin.

Some MPs see the tests as being in line with the policy of limiting the damage caused to public health by heroin use; others say that it gives a signal that anything goes in the Netherlands.

According to the European Monitoring Centre for Drugs and Drug Addiction, the Netherlands has the lowest number of drug related deaths in Europe—2.4 deaths per million in 1995, compared with 9.5 per million in France. □

Genetics body reviews cloning law

John Warden, *parliamentary correspondent, BMJ*

The UK government's Human Genetics Advisory Commission is to examine the adequacy of the existing law on cloning of human embryos after a lamb was cloned in Scotland (1 March, p 623).

The commission discussed at its inaugural meeting last week the new research on the cloning of sheep and acknowledged the public concern surrounding the possibility of human cloning. It decided to discuss with the Human Fertilisation and Embryology Authority the adequacy of the existing law, which, in Britain, forbids the cloning of human embryos by nuclear transfer. Cloning by splitting embryos may be carried out

only with a licence from the embryology authority, which has decided against granting any.

Head of the commission, Sir Colin Campbell, said that if the commission finds the law to be inadequate it would ask parliament to act. He said that whereas scientists could invent techniques, the commission would try to control and guide science rather than let it impose solutions. A consultative meeting with scientists is planned to review progress at the frontiers of genetic research that could impact on human genetics.

The commission also decided to give immediate priority to the implications of genetic test-

ing for insurance and has set up a group to explore the issues (1 March, p 625). Its future agenda will include questions of privacy, genetic services provided by the NHS, and the international dimension.

In the Commons a cross party motion by MPs called for a select committee to be set up to discuss the ethical issues of cloning. Its sponsor, Conservative MP Bill Cash, said: "We really cannot have these matters decided by scientific advisory committees." Ironically, the new commission was a reluctant concession by the government under pressure from the Commons science and technology committee.

In Brussels the president of the European Commission, Jacques Santer, announced a study of the moral and ethical issues of cloning. □

Communication increases donor transplant rate

Naomi Craft, *London*

The number of transplant donors in Britain can be increased substantially if critical care teams are helped to communicate with bereaved families, according to new findings presented to delegates at the *Transplantation in the 21st Century* conference, held at the Royal Society of Medicine, London, last week.

In the first study of its kind in Britain, staff from 40 intensive care and neurosurgical units in the north west of England attended one day courses on the management of emotions such as anger, fear, sadness, and uncontrollable grief. The cost of training was estimated at £110 (\$176) per person. In the hospitals where staff had received training, the donor rate increased by nearly 20%. There was no change in the donor rate from hospitals where staff were unaware of the trial.

The study forms part of the European donor hospital education programme, which originated in the Netherlands in 1991 and has already been introduced in over 30 countries worldwide. Originally set up by the Eurotransplant Foundation, the programme is a tool that aims to improve rates of organ donation by focusing on communication skills.

European data published in 1996 identified five situations in which inadequate efforts were being made to secure the donation of organs: donor detection, donor referral, organ maintenance, care of the family, and organ retrieval. "As many as 45% of potential donors are not detected and the major factor is failure to communicate with bereaved families," said Celia Wight from the Eurotransplant Foundation.

Mr Robert Johnson, president of the British Transplant Society said that Britain should be aiming to increase its number of donors from 15 per million population per year to nearer the Spanish rates of 27 per million population per year. The Department of Health has yet to decide the programme's future in Britain. □

Inhaled steroids increase risk of glaucoma

Zosia Kmietowicz, *London*

Inhaling steroids carries similar risks for developing glaucoma as taking them orally, Canadian researchers have established. They have shown that people with asthma who take high doses of inhaled steroids for more than three months are 44% more likely to develop glaucoma than non-users.

Investigators from the Royal Victoria Hospital in Montreal, Canada, studied the relation between the use of inhaled and nasal steroids in 48 118 patients in the Quebec universal health insurance programme for elderly people.

They found that both high dose administration and prolonged continuous use had to be present to raise the risk of glaucoma (*JAMA* 1997;277:722-7). High dose is defined as 1500 µg or more of the inhaled steroid flunisolide or 1600 µg of any other inhaled steroid. The use of low and medium doses of inhaled steroids or high doses of nasal steroids was not linked to an increased risk of glaucoma.

Steroids are generally believed to raise the pressure in the eye by increasing the resistance to outflow of the watery fluid in the front chamber of the eye. The authors recommend: "The use of these drugs should be routinely

questioned in newly diagnosed cases of ocular hypertension and open angle glaucoma. If patients receive high doses of steroids for several months, ocular pressure should be monitored."

Mr John Salmon, consultant ophthalmologist at the Oxford Eye Hospital at the Radcliffe Infirmary in Oxford, said that it was routine policy to check whether patients newly diagnosed as having glaucoma were asthmatic—primarily because the standard treatment is β blockers, which are contraindicated in these patients. "Most ophthalmologists accept that whichever route one uses, steroids will be

absorbed systemically even in small doses, so patients using them should be considered at risk of glaucoma," he said.

However, the risk of developing glaucoma as a result of steroid use is quite small, and in people with asthma the advantage of the treatment will usually outweigh the potential risk of glaucoma, added Mr Salmon.

A substantial number of asthmatic people currently take high dose steroids long term, according to Dr Sebastian Johnston, senior lecturer in medicine at Southampton General Hospital. But more evidence would be needed before deciding whether regular screening for increased intraocular pressure was necessary. "These patients are under constant supervision to see whether the dose of steroid can be reduced," he said. □

High doses of inhaled steroids have been linked with glaucoma

Doctors can reduce the harmful effects of poverty

Richard Smith, *BMJ*

Strong evidence exists to show that action taken by doctors, other health workers, and politicians can reduce the harmful effects of poverty on health, said Professor Trevor Sheldon, director of the NHS Centre for Reviews and Dissemination, at a conference in London last week.

A randomised controlled trial undertaken in Gary, Indiana, that guaranteed a minimum income to pregnant women in low income families by using a negative income tax showed a significant increase in birth weight. Other trials of nutrition support sponsored by the state have also shown significant improvements in child health.

Children constitute one of the groups worst affected by poverty, said Professor Sheldon. The number of children living in poverty has increased in Britain from 10% in the late 1970s to 32% now. The rise in unemployment and changes in social and fiscal policies are the main

causes of increased poverty in Britain.

Professor Sheldon said: "It is possible that some of the gains still to be made in the health of the nation are not going to be achieved by improving health services, or even behaving more healthily, but rather by improving the living standards of the poor." He advocated a statutory minimum wage, good quality and affordable child care, a large increase in child benefits, and a substantial increase in income support and housing benefit for families with children. "However, the most effective instrument for achieving a greater measure of income equality remains the progressive income tax."

In addition, there is much that health workers and authorities can do. The centre for reviews and dissemination will soon be publishing a systematic review of 122 studies that looked at specific ways of reducing health inequalities. The box

Evidence based interventions that reduce inequalities in health

- Targeting effective healthcare promotion and services at the groups with the greatest health needs
- Providing social, financial, and psychological support during pregnancy and childbirth
- Providing smoking cessation programmes to pregnant women
- Providing folic acid supplements before and around the time of conception
- Providing personal support for breast feeding
- Providing free school milk
- Targeting effective interventions to reduce accidents at children in deprived communities
- Providing sex education and available, accessible, and acceptable services to reduce teenage pregnancy
- Improving oral hygiene, reducing sugar intake, and promoting use of fluoride among people in deprived communities

shows some of the interventions that have firm evidence to support them.

The conference was organised by the Royal Colleges of General Practitioners, Nursing, and Physicians, the Faculty of Public Health Medicine, Action in International Medicine, and the *BMJ*. The plan is to form an intercollegiate task force to take

forward the work to reduce the harmful effects of poverty. Dr Sandy Macara, chairman of the council of the BMA, said: "This is a long term job, and we must be committed to working together as effectively to counter the effects of poverty on health as we have worked together to combat the harmful effects of tobacco." □

Loophole exposed in testing of child medicines

John Warden,
parliamentary correspondent, BMJ

Numerous gaps in the health care of children and young people in Britain—notably in the testing of child medicines—have been identified by the House of Commons Health Committee.

In the first two reports the MPs state that the special needs of children are given insufficient priority by policymakers and health service professionals. A main finding is that medicines for children are not controlled in the same way as for adults. Some medicines given to children are not licensed for human administration at all, and many are administered by a route, formulation, or dosage not approved by the Medicines Control Agency. The committee was astonished to find that such a situation existed

and regard it as unacceptable.

"It is surely wrong that clinicians dealing with children should be put in the position of having to guess what an appropriate dosage or route of administration might be," the report states. And although there is no evidence of children having been harmed through lack of "child specific" testing, the report says that this cannot be known for certain.

One example cited is vitamin K, given at birth to almost all babies, although there is no licence for it to be given orally, which is now the usual method. Theophyllines are commonly administered to premature infants, though they are not recommended for children aged under 3 years.

The Department of Health explained that the anomaly was due to pharmaceutical companies applying for licences for medicinal products before they have been fully tested in children. Because of the lack of data, licences are limited to adult use.

The MPs advise the Department of Health to implement

the conclusions of a recent, cross professional working party under Sir David Hull, which recommended a regimen of age specific licensing, clinical trials in children, expert surveillance, and fuller information for parents and health professionals.

The health committee, whose year long inquiry is still in progress, is also reporting on children's health care in the community, child and adolescent mental health care, and hospital services for children

and young people. It is the most comprehensive study of the subject for 20 years. The recurring theme is for child care services to be more child centred, less fragmented, and based on accurate information. □

The Specific Health Needs of Children and Young People and Health Services for Children and Young People in the Community: Home and School are published by the Stationery Office, price £11.50 and £13.60 respectively.

Medicines given to children are not controlled adequately

MARK CLARK/SPFL

Red Cross ensures health of hostages in Peru

Josh Hamilton, *New York*

For nearly 80 days 72 hostages have been caught up in a life or death political contest between the Peruvian government and the leftist Tupac Amaru Revolutionary Movement. The survival and good health of the diplomats and government officials from various countries is largely due to efforts by the International Committee of the Red Cross.

The Tupac Amaru Revolutionary Movement seized the Japanese ambassador's residence in Lima, Peru, on 17 December. Since then the Red Cross has been providing medical assistance and helped to prevent the break down of negotiations between the parties.

According to officials of both the Red Cross and the Peruvian government the hostages are all "fine," and none is medically in

jeopardy. Even those with previous medical conditions—such as the Bolivian ambassador, who has diabetes, and the Peruvian foreign minister, who has asthma—are being properly treated, with doctors providing care to the hostages on a daily basis.

According to Daniel Augsberger of the Red Cross in New York, proper care entails more than simply dispensing medicines. In a long term hostage situation, he explained, it is important to provide psychological support as well, to prevent psychosomatic illness. Such psychosomatic symptoms are quite real and can be serious in a hostage situation.

One way the Red Cross is providing the hostages with psychological support is by keeping the possibility of a negotiated solution alive. The guerrillas and

RICARDO MAZALAN/PHOTO

Doctors are providing daily medical care to the hostages in Peru

the government continue to press mutually unacceptable demands: the rebels want jailed comrades freed; the government refuses to be coerced into freeing avowed terrorists. Nevertheless, the Red Cross has repeatedly managed to use its credibility with both sides, shuffling between the parties and coaxing them back to the bargaining

table when talks break down.

Additionally, Mr Augsberger says, the Red Cross has used its "long term relationship" with the Peruvian security forces to argue for patience and help to forestall a more aggressive posture that might otherwise develop, given Peru's history of responding to terrorism and rebellion with draconian measures. □

Turkish doctors collude in torture

Paul Mansour, *Merseyside*

Doctors in Turkey are colluding in the torture of detainees, says a new report by Amnesty International. While many are put under intense pressure to cover up evidence of torture by issuing false medical or postmortem reports, others seem to collaborate willingly with the security forces, the human rights organisation claims.

Doctors have been prosecuted for "humiliating the Turkish State" by issuing medical reports that document torture injuries. The report describes how health workers have been affected by the 13 year armed conflict between the security forces and the separatist Kurdish Workers' Party in the south east of the country, a conflict that has resulted in widespread human rights' violations by both sides.

It cites numerous examples of doctors and nurses being at the sharp end of human rights abuses, including Dr İlhan Diken, who was tortured and sentenced to over three years' imprisonment for providing medical care to a member of the

Kurdish Workers' Party.

The report claims that such problems are not confined to the troubled south east. It quotes statements by the European Committee for the Prevention of Torture that torture is widespread and routine in police stations throughout Turkey. It says "inadequate" medical care in Turkish prisons causes continuing unrest, including regular hunger strikes.

Dr Hüseyin Gazi Yaman, a GP, had his jaw and pelvis fractured by the security forces for insisting on his right to belong to the Turkish Health Workers' Union. And in what the report describes as "a coordinated campaign" by the authorities against the Turkish Human Rights Foundation—which provides care for victims of torture—doctors working for the foundation are being prosecuted for refusing demands by the authorities to disclose details of torture victims.

The report praises the Turkish Medical Association for its defence of human rights and promotion of medical ethics. □

Turkey: Human Rights and the Health Professions is available from Amnesty International, 99-119 Rosebery Avenue, London WC1R 4RE.

HIV trial stopped early after good results

Alison Boulton, *London*

A United States National Institutes of Health trial of patients with advanced HIV disease has been stopped early because a combination of three drugs showed significant benefits.

The US National Institute of Allergy and Infectious Diseases study known as ACTG 320 examined the safety and efficacy of adding a protease inhibitor (indinavir) to two nucleoside analogue reverse transcriptase inhibitors—zidovudine and lamivudine.

The study included 1156 patients infected with HIV who had a CD4 T cell count below 200 per ml of blood and had not received protease inhibitors before. They received either indinavir (2400 mg/day) plus zidovudine (600 mg/day) and lamivudine (300 mg/day) or zidovudine and lamivudine alone for an average of 38 weeks. Patients who were intolerant to zidovudine were given stavudine instead.

The decision by an indepen-

dent data and safety monitoring board to recommend no further enrolment and immediate closure of the trial was taken last month after a review of the preliminary results. This showed that among those receiving triple combination treatment AIDS related illnesses, including opportunistic infections and cancers, and deaths fell by nearly half. There were 63 instances of disease progression in those receiving reverse transcriptase inhibitors alone compared with 33 in the triple combination arm. There were eight deaths in the triple treatment arm compared with 18 in the double treatment arm. No difference in safety or toxicity was found between the two groups.

"While these results further support the late use of protease inhibitors in advanced HIV disease, what we still need to assess is their efficacy in early use," said Dr Brian Gazzard, clinical research director at the Chelsea and Westminster Hospital in London.

Dr David Hawkins, consultant physician at the same hospital, said: "It remains impossible to predict how durable that survival benefit will prove, although the short to medium term results are encouraging." □

No sign of slowing down

Sir Richard Doll features in a photographic exhibition of 20th century British doctors which opens this week. He tells Hilary Bower about his lifetime of research

Sir Richard has published 436 pieces of research

At 84 Sir Richard Doll has no intention of slowing down in his pursuit of high quality research, which he has helped develop from data gathering into today's statistically rigorous epidemiology. Fêted as the world's leading cancer epidemiologist, his most famous finding—that smoking causes cancer—will be 50 years old in the year 2000. By then Sir Richard hopes he will have added to his body of work more research on the causes of childhood cancer, the health effects of electromagnetic fields, and the lifesaving links between diet and heart disease. But what he most wants for the millennium is simple. "I want to see it," he says.

At the last count Sir Richard has published 436 works—eight for every year of his working life. He has won 22 awards and held some of the most prestigious posts of clinical research. Technically retired, he is an honorary member of the Imperial Cancer

Research Fund's Cancer Studies Unit in Oxford, though his only concession to the honorary status is starting at 10 am, allowing time for the daily indulgence of breakfast in bed. One of those rare people who not only finds statistics exhilarating but can inspire others with them too, Sir

■ *"The whole promotion of tobacco ought to be banned—it is quite immoral, and it is unreasonable of governments to allow it to happen"*

Richard dropped mathematics for medicine at university after discovering that the only mathematics based job on the horizon in the 1930s was astronomy.

However, he lost little time in applying one love to the other. "I've always attempted to utilise statistical techniques in medicine. I wrote an article for the St Thomas's Hospital journal in 1935, when I was a medical student, showing the value of the χ^2

test for medical research. In fact I wasn't very popular because I showed that a favourite treatment of one of the surgeons for undescended testes was based on absolutely nothing of statistical importance."

His real introduction to epidemiology came, however, in 1948 when Professor Austin Bradford Hill invited him to join him an investigation into lung cancer. "I was fascinated by the enormous increase in mortality from lung cancer. At first I thought it was more likely to have something to do with motor cars. But I used to go around the wards checking the notes after discharge to see whether the diagnosis was confirmed or not and what immediately struck me was that if a person was a non-smoker the diagnosis was practically never confirmed, but if he or she was a smoker then it was almost always lung cancer."

Sir Richard is caustic in his criticism of government inaction over tobacco promotion after 50 years of proof. "We're lagging behind terribly. I have no doubt that the whole promotion of tobacco ought to be banned—it is quite immoral and it is unreasonable of governments to allow it to happen."

While the smoking findings are undoubtably his biggest achievement, Sir Richard says that the study that has given him most pleasure as a statistician was one he designed to find out whether, in the shadow of Hiroshima, there was a dose response with ionising radiation and cancer. "It was quite a difficult study to design and carry out, but it was the first paper to demonstrate that the effect of ionising radiation is proportional to the dose. Before then it was

thought cancer was only produced with big doses sufficient to cause macroscopic damage to tissue and that it was the repair process that caused cancer."

Always driven by a fascination for the effects of changes in society, Sir Richard says today's greatest health conundrum is the influence of diet, particularly on heart disease. "It's the biggest factor we can control, and it does look like the thing to do is to eat

more green vegetables and fruit. But what we want to find out is what are the chemicals within these things that do good."

But as chairman of the Advisory Group on Non-ionising Radiation he is also excited by the puzzle of electromagnetic fields. "It's the conflict between the biological and epidemiological evidence and the difficulty of getting firm epidemiological evidence that makes it so interesting. It would have a considerable impact on society if it were shown to produce a hazard—or if it didn't—because then we could stop worrying about it."

Unlike many who eulogise the old days, Sir Richard says two improvements—the recognition of the need for huge controlled trials and the resulting increase in collaboration—have changed the atmosphere of medical research completely. "Research is still competitive, particularly between countries, but in epidemiology the best research workers have recognised the need for collaboration. Meta-analysis is fine in concept but it can't really be done properly on published data. To get clear conclusions people need to put their data together in comparable form, like the latest collaboration on the effects of oral contraception on breast cancer."

But he thinks scientists these days are too eager to rush into print. "In 1950 we didn't say: 'Look, this is an association which looks fishy,' we said: 'The evidence is that smoking causes cancer of the lung'—and we were sure because we didn't publish until we'd checked the London results on patients in Leeds, Bristol, Cambridge, and Newcastle."

But the most concerning facet of research today is that money is channelled into such tightly focused work. "It is distressing that there is practically no money available now for people to try out their own mad ideas, and it's these that often provide the new clue that solves the problem," he says. He wants "x" per cent of the research budget let loose for researchers to follow up their pet theories, or, he warns, the millennium might end up short on new ideas. □

The photograph of Sir Richard Doll (above) features in an exhibition, organised by the *BMJ*, *With Head and Heart and Hand: Photographs of 20th Century British Doctors*, at the National Portrait Gallery in London. Admission is free.