Chronic venous ulcer

Niren Angle, John J Bergan

Venous ulceration is the most severe and debilitating outcome of chronic venous insufficiency in the leg. It is a common problem in clinical practice, with an estimated prevalence of 1-1.3%.^{1,2} Such ulceration is part of the complex of chronic venous insufficiency and is associated with distal vein hypertension. Chronic venous insufficiency and its accompanying venous hypertension have been termed the post-thrombotic syndrome, but it is now known that primary valvar incompetence, not just prior thrombosis, is an important cause of venous leg ulcer.³ Dysfunction of the superficial or perforating veins of the legs, alone or in combination, may be the only finding in leg ulceration.⁴ Nevertheless, a risk factor for venous ulceration remains a previous episode of deep vein thrombosis, which may or may not destroy venous valve function.⁵ It is clear, however, that a venous ulcer in the leg occurs mainly from reflux and not usually from persistence of the original obstructive process.^{6,7}

Methods

We relied largely on our own files to prepare this review. We accumulated these by weekly searches of *Current Contents*, which provides tables of contents of almost all medical publications published worldwide. Our files are arranged into categories, and the category of venous ulcer crosses many of these designations. Searches for evidence based information about venous disorders are often disappointing, but we used at least level 2 information whenever possible, tempered by our own experience. Diagnosis and treatment of venous disorders commands greater interest, and level 1 information is starting to appear, including treatment of venous thromboembolic disorders. This heralds better care of patients with venous problems. (Level 1 information means findings from large prospective randomised studies with definite conclusions that should be considered when managing any patient with the specified problem; level 2 information means findings from smaller prospective randomised studies with less definite results that may be relevant to some patients.)

Causes of severe chronic venous insufficiency

Gravitational reflux

Venous insufficiency can be classified as primary or secondary. Primary venous insufficiency occurs when no obvious cause of valvar dysfunction is identified.

Summary points

- Venous leg ulcers are common, with a pathophysiology is similar to that of varicose veins
- Ulceration is thought to result from the distal adherence, trapping, and activation of leucocytes, which causes tissue destruction
- Non-invasive testing while providing compression treatment clarifies the pathophysiology
- Most patients will benefit from simple surgery
- Appropriate intervention decreases costs by limiting recurrence of venous ulcer

Many causes have been postulated, including loss of elasticity of the vein wall.⁸ Secondary venous insufficiency is the proper term when there is an identifiable condition that has caused development of valvar dysfunction (such as previous deep vein thrombosis).

Whatever the cause, gravitational reflux from valvar dysfunction results in early refilling of the venous pool in the leg after muscle contraction (see box). Arnoldi elegantly described the normal physiology of the muscle pump and the effect of valve reflux on venous pressure in the leg after exercise.⁹ When calf muscles contract there is a rise in pressure in both superficial and deep vein systems. On relaxation, pressure in both sets of veins is below that found in the resting muscle. This is due to the protective effects of competent valves preventing downward flow in the venous system. During muscular contraction, pressure in the muscular compartment and the deep vein system rises above that in the superficial vein system. During muscular diastole, pressure in the deep vein system falls below that of the superficial vein system.¹⁰ The time to refill the veins of the calf after a series of muscular contractions is shortened in patients with valvar insufficiency because of venous reflux. This causes a progressive and sustained increase in calf vein pressure.¹¹

This pressure, termed ambulatory venous hypertension, has been the subject of much interest. It is due to the weight of the blood column from the right atrium transmitted through a valvar abdominal and pelvic veins to dysfunctional limb veins. The peripheral valve dysfunction may be primary or secondary, but the haemodynamic effect of incompetent valves is the same—distal vein hypertension. It is thought that this leads to capillary dilatation and increased leakage of plasma, plasma proteins, and red blood cells.

University of California, San Diego San Diego, CA, USA

Niren Angle,
resident in surgery

John J Bergan,
professor of surgery

Correspondence to:
Professor Bergan,
North Coast
Surgeons Medical
Group, 9850
Genesee Avenue,
Suite 560, La Jolla,
CA 92037, USA.

BMJ 1997;314:1019-23

Causes and consequences of chronic venous insufficiency

Causes:

- Gravitational (hydrostatic) reflux
- Muscular compartmental reflux (hydrodynamic) through perforating veins
- Leucocyte trapping and activation

These result in:

- Pigmentation
- Dermatosclerosis
- Ulceration

Compartmental pressure

Varicosities of the superficial vein system, lipodermatosclerosis, cutaneous hyperpigmentation, and leg ulcers are often associated with deep to superficial outflow through incompetent perforating veins. As mentioned above, this can be due to venous dilation without previous thrombosis.

The consequence of incompetent perforating veins is exposure of superficial veins, venules, and reticular veins to the high systolic pressures developed in calf muscles during muscular contraction. The check valve function of perforating veins is no longer operative.¹² Chronic exposure of the superficial system to high pulsatile pressures generated by muscular contraction dilates and elongates veins and venules of the skin and subcutaneous tissues.¹⁰ The repeated pulses of high pressure during walking has been termed the "hydraulic ram" by Arnoldi.⁹ This produces a localised venous hypertension, which in turn causes filtration oedema. This continues until tissue pressures rise to restore pressure equilibrium. Since tissues are relatively compliant, the limb can become considerably oedematous before such equilibrium is reached.¹³

Leucocyte trapping and activation

Microcirculation in tissues surrounding venous ulcers has been the subject of much study. Among the most important observations is that there is a greater accumulation of leucocytes in the dependent limbs of patients with chronic venous insufficiency than in those of healthy subjects and limbs with varicose veins only.^{14 15} The most plausible theory about venous ulceration implicates trapping of leucocytes within the microcirculation, with their subsequent activation.¹⁶ Endothelial cells, presumably activated by hypoxia or other cause, stimulate adherence of leucocytes. The activated leucocytes then release oxygen free radicals and other toxic products such as elastase, tumour necrosis factor, and collagenase, which injure the surrounding tissue. This process is potentiated by blockage of capillary blood flow by adherent neutrophils and thrombosis induced by platelet activation.¹⁷

Increased production of collagenase and elastase in leucocytes from patients with chronic venous insufficiency compared with controls adds weight to the theory that neutrophil trapping and activation are the final pathway to ulceration.¹⁸ This theory is provocative and attractive because it suggests the possibility of drug treatment of venous leg ulcer in the future.

Thus, to summarise, the effects of gravitational reflux added to muscular compartment pressure are

transmitted through perforating veins in a pulsatile manner. This favours adherence of leucocytes and their subsequent activation, causing localised tissue destruction. This produces the venous leg ulcer.

Evaluation

Physical findings

Proper treatment of venous leg ulcer depends on accurate diagnosis. The term leg ulcer is therefore too general and requires qualification by a statement of cause supported by objective evidence. With that in mind, it should be remembered that the leg ulcer is accompanied by chronic venous hypertension. Therefore, a history of chronic venous insufficiency, with its attendant signs and symptoms of oedema and hyperpigmentation, is expected (see box). Virtually all patients with a venous ulcer exhibit perimalleolar skin changes of lipodermatosclerosis.¹⁹ This term describes the pigmented, indurated skin in a perimalleolar and supramalleolar distribution. Oedema of the leg is also almost always present. Thus, absence of chronic oedema in the affected leg or absence of such skin changes would suggest a diagnosis other than venous insufficiency. There are many other causes of leg ulceration.

Continuous wave Doppler ultrasonography

Doppler ultrasound scanning has become part of the physical examination in clinical vascular practice. It is useful in assessing arterial and venous patency and to ascertain the presence of venous reflux. Venous leg ulcer is most commonly accompanied by venous reflux, and, for this to be detected, the patient must be upright while the following sites are investigated: the greater saphenous vein at the junction and along its course, and the lesser saphenous vein in the popliteal fossa and distally. Perforating veins should be sought. The patency and competency of the femoral vein in the groin, the popliteal vein, the posterior tibial vein behind the medial malleolus, and the anterior tibial vein can also be ascertained.²⁰ Spontaneous flow cannot be detected with a 8.0-10.0 MHz probe when the patient is upright. Instead, augmented flow in the various venous segments can be elicited by manual compression of the sole of the foot or calf. Releasing the compression will elicit reverse flow through incompetent valves. Venous reflux can also be elicited by a deep inspiration, a Valsalva manoeuvre, or proximal compression.

Proximal obstruction to flow can be detected in a supine patient by observing the venous flow during spontaneous breathing. Continuous flow unaffected by movement of the abdominal wall may be noted when the affected leg is compared with the other leg. This may give useful information about the degree of proximal obstruction.

Signs of venous hypertension

- Perimalleolar oedema
- Pigmentation
- Lipodermatosclerosis
- Eczema or dermatitis
- Ulceration

Duplex Doppler ultrasonography

Ultrasonic B mode scanning combined with pulsed Doppler flow examination allows for both anatomical and functional assessment of the venous system. Flow velocity, flow rate, and the diameter of the veins can be measured. Deep vein visualisation may reveal occlusion, recanalised lumina, or irregularities in vein walls. Incompetent perforating veins can also be visualised. Real time colour duplex scanning makes orientation of venous flow much easier and provides information about reflux in specific veins. Phlebography is reserved for special situations.

Treatment

Chronic deep vein insufficiency with ulceration due to blood stasis is not yet curable. Palliation in the form of healing ulcers and preventing their recurrence has focused on surgically eliminating or counteracting the transmission of increased venous pressure to the skin (see box).

Compression

Limb compression is the fundamental conservative treatment for chronic venous ulcers. The most commonly used methods are four layer bandaging, the Unna boot, elastic stocking, and, in the United States, the CircAid nonelastic compression legging.

Elastic stockings

First developed in the late 19th century, elastic compression stockings were not fully effective because they provided uniform or even reverse gradient compression.²¹ Conrad Jobst developed graded elastic compression stockings in the late 1940s.²² Although their superiority over uniform compression stockings has not been established, considerable experimental evidence has led to general acceptance of this hypothesis. The beneficial effects of gradient compression stockings are probably due to local alteration of microvascular haemodynamics and regional Starling forces, relieving oedema rather than any direct effect on deep vein haemodynamics. Thus, stockings do not cure venous hypertension but do protect the skin from its effects.

The initial treatment may include complete bed rest and elevation of the leg to relieve the oedema. After resolution of the oedema and local wound cleansing, patients can be fitted with elastic compression stockings with 30 to 40 Torr gradient to allow ambulatory care.

The presence of occlusive arterial disease is a relative rather than an absolute contraindication to the use of external compression. However, the pressure applied must be low. Doppler ultrasound measurement of the systolic pressure at the ankle will provide an objective estimate of arterial insufficiency.²³ An ankle brachial index of less than 0.90 should alert the doctor to the presence of arterial insufficiency.

Non-elastic compression

Severe forms of oedema require non-elastic compression. The most commonly used device is the Unna gel paste gauze boot. One current formulation consists of calamine, zinc oxide, glycerin, sorbitol, gelatin, and magnesium aluminum silicate. This dressing is

Treatments for venous ulceration

Compression

- Elastic compression stockings (gradient)
- Gel paste gauze boots (Unna)
- CircAid
- External pneumatic compression

Drug treatments (questionable value)

- Systemic agents (minimally effective)
 - Zinc
 - Fibrinolytic agents
 - Pentoxifylline
- Topical agents (not recommended)
 - Antibiotics
 - Free radical scavengers
 - Hydrocolloid dressing

Surgery (alone or combined)

- Skin grafting
- Free flap grafting
- Superficial vein stripping
- Perforating vein interruption
- Valve plasty
- Thrombolysis, dilation, stenting

designed to provide both compression and topical treatment. A rolled gauze bandage impregnated with the gel paste is applied from the metatarsals of the foot to just below the knee. Next, a four inch wide, continuous gauze dressing is applied to absorb moisture from the first layer. The outer layer is an elastic wrap with graded compression. The bandage stiffens after drying and, being non-elastic, prevents further oedema.

Existing oedema decreases through proximal flow in lymphatics and veins. One advantage of the Unna boot is that the dressing is applied and removed by staff and requires minimal compliance by patients. Several controlled studies suggest that this treatment improves healing times, although the studies all have important qualifiers that weaken their conclusions.

The American CircAid achieves the advantages of the Unna boot yet allows dressings to be changed. It consists of pliable, inelastic, adjustable compression bands that wrap around the leg and are held in place by Velcro fasteners.

Local wound treatment

Antibiotics have been advocated in treating venous ulceration, but in the absence of local infection they seem to have little effect on healing.²⁴ Topical antibiotics also do not help wound healing. Although povidone-iodine, acetic acid, and sodium hypochlorite have been shown to be toxic to cultured fibroblasts, they are helpful and remain in use. Unlike many other ointments and creams, they are not allergenic.²⁵

Hyperbaric oxygen

Hyperbaric oxygen has been used in treating a variety of conditions, but there is no convincing evidence that it has any beneficial effect on healing of a chronic venous ulcer.²⁶ It should not be considered as treatment.

Surgery

Surgical treatment of venous ulcer may be directed toward eliminating the cause of the venous hypertension or treating the ulcer itself by skin grafting. No single operation is applicable to every type of venous

ulceration, and surgery is but one treatment of many that can be used.

In principle, correcting superficial vein insufficiency or perforating vein outflow should be done early. Deep vein reconstruction by bypass, stenting, or valve plasty should be done only rarely and as a last resort. Incompetent perforating veins should be suspected in every case and searched for by Doppler ultrasonography. Their interruption by endoscopic techniques has been shown to accelerate healing of ulcers.²⁷

Superficial stripping and excision of varices

While conservative compression treatment is being pursued, diagnostic procedures should be done to clarify the patterns of venous reflux and to detect segments of venous occlusion. Reflux is usually the only abnormality uncovered by diagnostic testing.²⁷ It is important to know that in advanced chronic venous stasis, superficial reflux may be the only finding. Hanrahan *et al* studied 95 cases of venous ulceration with duplex ultrasonography: they found that 16% of the affected legs had only superficial incompetence while 19% had superficial incompetence combined with perforating vein incompetence—thus, 35% of cases were potentially curable by simple surgery.²⁸ Our own study of 58 cases of severe chronic venous insufficiency showed that 17% of the affected legs exhibited only superficial reflux and that superficial reflux was a major contributor to chronic venous dysfunction in another 17 limbs.²⁹ Reflux proceeding distally down the greater saphenous vein and re-entering perforating veins in the proper direction may overload the deep vein system and cause elongation and dilatation of the deep vessels so that their valves become incompetent. This problem is treated relatively easily by stripping the superficial veins and excising the varices.²⁹

Subfascial perforating vein interruption

Robert Linton in the United States³⁰ and Cockett and Dodd in Britain emphasised the need to control perforating veins in the effective treatment of chronic venous insufficiency. Unfortunately, morbidity from poor healing of the extensive incisions limited the application of their operations, although recurrence rates were mainly only 15-20%. The advent of endoscopic surgical techniques has made interruption of perforating veins much less morbid. Clinically important perforating veins from the medial malleolus to the knee as well as the gastrocnemius point and the soleal point are accessible to endoscopic interruption.³¹ This eliminates the problems from extensive incisions and accelerates the healing of the leg ulcer without creating a major problem of infection (see fig 1).

Excision and skin grafting

Success in covering venous ulcers can be achieved by split thickness skin grafting.³² This may be performed at the same time as division of the perforating veins or may precede or follow as a separate procedure. Grafting is not necessary for small ulcers but may decrease healing times for large ulcers. The wound bed must be debrided gently yet effectively—all non-viable, infected tissue must be excised down to healthy tissue, on which the skin graft may be laid down for temporary closure. For definitive coverage, wide excision of all surround-

ing pigmented tissue down to and including the fascia must precede coverage with a graft.

Excision and free flap coverage

Excision of the ulcer bed as described above will result in a large soft tissue defect. When such a defect is anticipated, coverage with a free flap may be desirable.³³ A free flap transfer also speeds healing and provides a local venous arborisation containing hundreds of competent valves. Bone defects can also be reconstructed using the iliac crest for a graft with attached muscle.³⁴ Comparing postoperative recovery time with local wound care shows that the procedure is cost effective under the right circumstances.

Prophylaxis for deep vein thrombosis

Since deep vein thrombosis is an important cause of subsequent chronic venous insufficiency, prevention of recurrence and further attendant damage to valves must be included in surgical intervention. Subcutaneous heparin, pneumatic compression devices, and low molecular weight heparin have been shown to reduce the incidence of postoperative deep vein thrombosis. Outpatient surgery allows early mobilisation of patients, and simultaneous prophylactic measures against deep vein thrombosis may be pursued with low molecular weight heparin.

Aftercare and follow up

Patients seen with early signs and symptoms of chronic venous insufficiency must have the cause ascertained. Preventing the development or recurrence of an ulcer depends on correcting the causes of the chronic venous insufficiency. In order to prevent further deterioration, especially if surgical correction is not possible, patients must be instructed in skin care, avoidance of prolonged standing, and the permanent use of compression stockings. After the ulcer has healed, the patient must be enrolled in a follow up programme supervised by qualified staff. This must be vigilant and regular and involve the patient so that compliance is improved. Long term results revolve around continuity of good care, ulcer care clinics, home health nursing, and regular evaluation by a doctor. These are vital to the amelioration of this common and morbid condition.

Fig 1
Preoperative photograph of leg with two year old, non-healing venous ulceration: markings show refluxing varicose veins and perforating veins identified by duplex ultrasonography (circles)

- 1 Callam MJ, Ruckley CV, Harper DR, Dale JJ. Chronic ulceration of the leg: extent of the problem and provision of care. *BMJ* 1985;290:1855-6.
- 2 Widmer LK. *Peripheral venous disorders. Prevalence and sociomedical importance. Basel Study III*. Bern: Hans Huber, 1978:43-50.
- 3 Lees TA, Lambert D. Patterns of venous reflux in limbs with skin changes associated with chronic venous insufficiency. *Br J Surg* 1993;80:725-8.
- 4 Nelzen O, Bergqvist D, Lindhagen A. Leg ulcer etiology: a cross-sectional population study. *J Vasc Surg* 1991;14:557-64.
- 5 Partsch H. Investigations on the pathogenesis of venous leg ulcers. *Acta Chir Scand* 1988;544(suppl):25-9.
- 6 Labropoulos N, Leon M, Nicolaides AN, Giannoukas AD, Volteas N, Chan P. Superficial venous insufficiency: correlation of anatomic extent of reflux with clinical symptoms and signs. *J Vasc Surg* 1994;20:953-58.
- 7 Raju S, Frederick R. Venous obstruction: an analysis of 137 cases with hemodynamic, venographic, and clinical correlations. *J Vasc Surg* 1991;14:305-13.
- 8 Travers JP, Brookes CE, Evans J, Baker DM, Kent C, Makin GS, *et al*. Assessment of wall structure and composition of varicose veins with reference to collagen, elastin, and smooth muscle content. *Eur J Vasc Endovasc Surg* 1996;11:230-7.
- 9 Arnoldi CC. Venous pressure in the leg of healthy human subjects at rest and during muscular exercise in the nearly erect position. *Acta Chir Scand* 1965;130:570-83.
- 10 Arnoldi CC. Venous pressure in patients with valvular incompetence of the veins of the lower limb. *Acta Chir Scand* 1966;132:427-30.
- 11 Payne SPK, London NJM, Newland CJ, Thrush AJ, Barrie WW, Bell PRF. Ambulatory venous pressure correlation with skin condition and role in

- identifying surgically correctable disease. *Eur J Vasc Endovasc Surg* 1996;11:195-200.
- 12 Bergan JJ. New developments in surgery of the venous system. *J Cardiovasc Surg* 1993;1:624.
 - 13 Guyton AC. Interstitial fluid pressure II. Pressure volume curves of the interstitial space. *Circ Res* 1965;16:452.
 - 14 Moyes C, Cederholm-Williams SA, Michel CC. Haemoconcentration and accumulation of white cells in the feet during venous stasis. *Int J Microcirc Clin Exp* 1987;5:311-20.
 - 15 Thomas PR, Nash GB, Dormandy JA. White cell accumulation in dependent legs of patients with venous ulceration: a possible mechanism for trophic changes in the skin. *BMJ* 1988;290:1693-5.
 - 16 Sarin S, Andaz A, Shields DA, Scurr JH, Coleridge Smith PD. Neutrophil activation in venous disease. *J Vasc Surg* 1993;17:444.
 - 17 Bauersachs J, Fleming I, Busse R. Pathophysiology of chronic venous insufficiency. *Phlebology* 1996;11:16-22.
 - 18 Whiston RJ, Hallet MB, Lane IF, Harding KG. Lower limb neutrophil oxygen radical production is increased in venous hypertension. *J Vasc Surg* 1993;17:445.
 - 19 Mayberry JC, Moneta GL, Taylor LM, Porter JM. Nonoperative treatment of venous stasis ulcer. In: Bergan JJ, Yao JST, eds. *Venous disorders*. Philadelphia: W B Saunders, 1989: 383.
 - 20 Goldman MP, Bergan JJ, eds. *Ambulatory treatment of venous disease: an illustrative guide*. St Louis, C V Mosby, 1995: 13.
 - 21 Johnson G Jr. The role of elastic support in venous problems. In: Bergan JJ, Yao JST, eds. *Surgery of the veins*. Orlando: Grune and Stratton, 1985: 541-50.
 - 22 Conrad Jobst and the development of pressure gradient therapy for venous disease. In: Bergan JJ, Yao JST, eds. *Surgery of the veins*. Orlando: Grune and Stratton, 1985: 529-40.
 - 23 Callam MJ, Harper DR, Dale JJ, Ruckley CV. Arterial disease in chronic leg ulceration: an underestimated hazard? Lothian and Forth Valley leg ulcer study. *BMJ* 1987;294:929-31.
 - 24 Gilchrist B, Reed C. The bacteriology of chronic venous ulcers treated with occlusive hydrocolloid dressings. *Br J Dermatol* 1989;121:337.
 - 25 Lineaweaver W, Howard R, Souey D, George RG. Topical antimicrobial toxicity. *Arch Surg* 1985;120:267.
 - 26 Hyperbaric oxygen therapy: review article. *N Engl J Med* 1996;334: 1642-6.
 - 27 Glociczki P, Bergan JJ, Menawat S, Hobson RW, Kistner RL, Lawrence PF, et al. Safety, feasibility, and early efficacy of subfascial endoscopic perforator surgery (SEPS): a preliminary report from the North American Registry. *J Vasc Surg* (in press).
 - 28 Hanrahan LM, Araki CT, Rodriguez AA, Kechejian GJ, LaMorte WW, Menzoian JO. Distribution of valvular incompetence in patients with venous stasis ulceration. *J Vasc Surg* 1991;13:805.
 - 29 Walsh JC, Bergan JJ, Moulton SC, Beeman S. Proximal reflux adversely affects distal venous function. *Vasc Surg* 1996;30:89-96.
 - 30 Linton R. Communicating veins of the lower leg and the operative technique for their ligation. *Ann Surg* 1938;107:582.
 - 31 Bergan JJ, Murray J, Greason K. Subfascial endoscopic perforator vein surgery: a preliminary report. *Ann Vasc Surg* 1996;10:211-9.
 - 32 Wood MK, Davies DM. Use of split-skin grafting in the treatment of chronic leg ulcers. *Ann R Coll Surg Engl* 1995;77:222-3.
 - 33 Dunn RM, Fuden GM, Walton RL, Anderson FA Jr, Malhotra R. Free flap valvular transplantation for refractory venous ulceration. *J Vasc Surg* 1994;19:525-31.
 - 34 Davies DM. Microvascular surgery. *BMJ* 1985;290:910.

Grand Rounds—Hammersmith Hospital

Hazards of running a marathon

Creatine kinase MB can be raised without myocardial infarction

The notion that participating in strenuous physical exertion could be hazardous to health dates from 490 BC, when Pheidippides allegedly dropped dead after running 39 km to Athens and delivering news of victory over the Persians on the battlefield at Marathon.¹

Case history

A 61 year old physician volunteered to run in the 1993 London marathon to raise money for a medical charity. He trained for the previous four months, running an average of 35 km a week, and in the marathon he completed the 42.2 km in about 4.5 hours. His race was uneventful apart from a brief spell of dizziness towards the end, which was relieved by carbohydrate and which he attributed to hypoglycaemia. Earlier in the race he had passed a middle aged runner who was being resuscitated unsuccessfully after sustaining a cardiac arrest.

Apart from myalgia in his legs he felt well after the race (fig 1) and dined out with friends and relatives that evening. He went to work as usual the next day, but in view of the persistence and severity of his myalgia he had blood taken for measurement of serum creatine kinase activity. The results showed a total creatine kinase activity of 1800 IU/l (normal <200 IU/l) and a creatine kinase MB of 38 ng/ml (normal <6 ng/ml). The chemical pathology registrar on duty regarded these values as pathognomonic of myocardial infarction. However, in view of his general wellbeing and lack of cardiorespiratory symptoms the physician discounted this diagnosis. His myalgia gradually subsided, and he was jogging again within a week.

Fig 1 61 Year old physician after running the London marathon—apart from myalgia in his legs he felt well

Because of his secretary's success in raising sponsorship money the physician won a free trip to participate in the New York marathon later that year. He again completed the race uneventfully but afterwards learnt that two other runners had sustained fatal cardiac arrests.

See Personal view by Kjeldby

Clinical Sciences Centre, Royal Postgraduate Medical School, Hammersmith Hospital, London W12 0NN

Case presented by:

G R Thompson, professor of clinical lipidology

Chairman: J Scott, professor of medicine

Discussion group:

M J Walport, professor of rheumatology

C M Oakley, professor of clinical cardiology

J M B Hughes, professor of thoracic medicine

C J Bulpitt, professor of geriatric medicine

Series edited by: Dr S D Taylor-Robinson.

BMJ 1997;314:1023-5

After publication in a newspaper of an item describing his experiences, the physician received a letter from a runner who, several years previously, had been admitted to hospital after falling and sustaining concussion during a 16 km race. "The enzyme test suggested that I had indeed had a heart attack. But I did not seem to have any of the regular symptoms—breathlessness and the like—and I did not feel unwell. The doctor in charge told me 'marathon running is not for you, old boy.' I celebrated that by winning the National Veterans over 55 marathon in 2 hours 54 minutes ... only 14 months after my 'heart attack.'" The runner subsequently ran several more marathons including the London marathon, which he completed in less than three hours at the age of 60 years.

Comment

Pseudomyocardial infarction after a marathon is well documented. Siegel *et al* studied runners in the 1979 and 1980 Boston marathons and found that creatine kinase MB values rose from 6.9 IU/l before to 105 IU/l 24 hours after the event.² However, technetium-99m pyrophosphate scans performed 24-72 hours after the race were all normal. Subsequently these workers demonstrated normal thallium scans immediately after a marathon and concluded that the increase in creatine kinase MB activity was of skeletal origin.³ This was borne out by muscle biopsies that showed an increase in creatine kinase MB activity during training⁴; the MB isoenzyme accounted for 8.9% of total creatine kinase activity in the skeletal muscle of trained marathon runners versus only 3.3% in controls.⁵ Additional evidence of a non-cardiac origin for the increase in serum creatine kinase MB activity after a marathon was the lack of any rise in cardiac troponin-I or C reactive protein concentrations.⁶ Therefore a rise in creatine kinase MB activity after strenuous exercise in the absence of other indices of myocardial damage should be regarded as indicative of nothing more sinister than transient rhabdomyolysis of skeletal muscle. Electrocardiograms can also be misleading as a high proportion of runners evince electrocardiographic abnormalities both before and after a marathon, which may be of neurogenic origin.⁷

Sudden death during a marathon can be due to hypertrophic cardiomyopathy,⁸ especially in younger athletes, but is more often due to myocardial infarction consequent on coronary artery disease.⁹ Runners in the latter category are usually middle aged and may show evidence of severe atherosclerosis at postmortem examination.¹⁰ It has long been debated, however, whether the incidence of sudden death from this cause in runners during a marathon is greater than would be expected in sedentary members of the population of similar age and sex.

Up to and including 1993 there were two deaths from coronary heart disease in 14 London marathons. On the basis of an estimated total of 275 000 runners who competed during this period (85% of whom were male with a mean age of 39 years) and with extrapolation from an average time of 3.75 hours, the derived annual death rate from coronary heart disease during the London marathon can be estimated to be 2% (this value is derived by multiplying the death rate per 100 000 male runners per hour by 24×365).

Fig 2 New York marathon crossing the Verrazano Narrows Bridge—sudden death during a marathon is usually due to myocardial infarction consequent on coronary artery disease

Furthermore, there have been three sudden deaths during the past 25 New York marathons.¹¹ On the basis of similar assumptions to those for the London marathon this works out at a derived annual death rate of 1.68%. In comparison, the annual death rate from coronary heart disease for men in England and Wales in 1992 was 0.03% for age 35-44 years, 0.14% for age 45-54 years, and 0.47% for age 55-64 years.¹² Despite the small numbers, this suggests that the death rate from coronary heart disease during a marathon may be considerably greater than expected. However, as proposed more than 15 years ago,¹⁰ a proper epidemiological survey is needed to settle this issue.

Although taking regular physical exercise seems to protect against the triggering of a myocardial infarction after strenuous physical exertion,¹³ this may not always hold true. If participants in a marathon have pre-existing, albeit silent, coronary atherosclerosis, as is sometimes the case,¹⁰ the extreme conditions endured during the three to four hours of the race may precipitate myocardial infarction, presumably by causing rupture of a plaque. This does not, of course, diminish the value of the regular, less strenuous physical exercise undertaken by marathon runners and others during the rest of the year, which affords a considerable degree of protection against coronary heart disease in the long term.¹⁴

Postulated mechanisms for the protective effect of exercise include a reduction in blood pressure, decreased myocardial sensitivity to catecholamines, increased fibrinolysis, enhanced insulin sensitivity, and an improved lipid profile. There is increasing evidence

that the last of these mechanisms influences the composition of atheromatous plaques so as to render them less susceptible to haemodynamic stress and thus prevent the plaque rupture that frequently precedes myocardial infarction.¹⁵ Arrest of angiographically gauged progression of coronary artery disease requires the expenditure of about 6.27MJ a week whereas to induce regression of atherosclerotic lesions necessitates expending 9.20 MJ a week, equivalent to five to six hours of exercise.¹⁶ This has led to the proposal by the Centers for Disease Control and Prevention and the American College of Sports Medicine that every adult should undertake 30 minutes or more of moderate intensity physical activity on most, preferably all, days of the week.¹³

Discussion

MJW: How many marathon runners die during their training?

GRT: There are no statistics to draw on, but some undoubtedly do die, including the three runners described in South Africa by Noakes *et al* who were killed by a car while on a training run.¹⁰ Interestingly, two of these runners were incidentally found to have pronounced coronary atherosclerosis at postmortem examination.

MJW: What is the explanation for the increase in creatine kinase MB in skeletal muscle, and does it have any physiological difference from the normal muscle isotype?

GRT: Regenerating skeletal muscle fibres seem to be rich in creatine kinase MB, as are fetal skeletal myocytes, and the increase found in runners may simply reflect a state of chronic injury resulting from repetitive training.⁵

CMO: Do you think that sudden cardiac death during a marathon would always be caused by plaque rupture in a person with occult coronary disease or that ventricular fibrillation with cardiac arrest might also be caused by regional ischaemia without plaque rupture?

GRT: I am sure that the latter is a possibility, although I imagine that in this case the physician might have had symptoms of ischaemia during training.

JMBH: Systolic blood pressure rises during exercise, and in a marathon the rise would be maintained for three to four hours. Could this be a factor in causing rupture of atheromatous plaques?

GRT: I am sure it could, especially in someone who was hypertensive to start with, presumably by increasing haemodynamic shear stress on the vessel wall.

JS: I believe sudden death is much more common during exercise in those who are not physically trained. The classic is the Saturday morning squash player who dies suddenly. I have always presumed that this was due to acute arrhythmia rather than coronary occlusion. Do you know if there are any postmortem data on this?

GRT: Squash seems to be particularly risky in this respect, and, as you surmised, there is often no

evidence of a fresh thrombus at postmortem examination. In one series thrombus was present in only five of 20 deaths from squash with postmortem evidence of coronary atherosclerosis.¹⁷ This supports the suggestion that such deaths are often caused by an arrhythmia induced by myocardial ischaemia¹⁸ or catecholamines.¹⁹

CJB: You are very persuasive of the benefits of exercise and I am sure that your judgment is not clouded by the "endorphin buzz" induced by exercise and recognised by many athletes. However, as with all powerful medicines we need to know the risks. Do you know the state of your 61 year old's knees, ankles, and spine?

GRT: His ankles and spine are fine, but he has had problems with both of his anterior cruciate ligaments. However, these injuries were incurred during other activities and were not from running, although the latter probably compounded the damage.

The *BMJ* welcomes grand rounds from other hospitals.

- 1 Martin DE, Benario HW, Gynn RWH. Development of the marathon from Pheidippides to the present, with statistics of significant races. *Ann NY Acad Sci* 1977;301:820-52.
- 2 Siegel AJ, Silverman LM, Holman BL. Elevated creatine kinase MB isoenzyme levels in marathon runners. *JAMA* 1981;246:2049-51.
- 3 Siegel AJ, Silverman LM, Holman BL. Normal results of post-race thallium-201 myocardial perfusion imaging in marathon runners with elevated serum MB creatine kinase levels. *Am J Med* 1985;79:431-4.
- 4 Apple FS, Rogers MA, Casal DC, Lewis L, Ivy JL, Lampe JW. Skeletal muscle creatine kinase MB alterations in women marathon runners. *Eur J Appl Physiol* 1987;56:49-52.
- 5 Siegel AJ, Silverman LM, Evans WJ. Elevated skeletal muscle creatine kinase MB isoenzyme levels in marathon runners. *JAMA* 1983;250:2835-7.
- 6 Cummins P, Young A, Auckland ML, Michie CA, Stone PC, Shephstone BJ. Comparison of serum cardiac specific troponin-I with creatine kinase, creatine kinase-MB isoenzyme, tropomyosin, myoglobin and C-reactive protein release in marathon runners: cardiac or skeletal muscle trauma? *Eur J Clin Invest* 1987;17:317-24.
- 7 Ohman EM, Teo KK, Johnson AH, Collins PB, Dowsett DG, Ennis JT, *et al*. Abnormal cardiac enzyme responses after strenuous exercise: alternative diagnostic aids. *BMJ* 1982;285:1523-6.
- 8 Noakes TD, Rose AG, Opie LH. Hypertrophic cardiomyopathy associated with sudden death during marathon racing. *Br Heart J* 1979;41:624-7.
- 9 Noakes T, Opie L, Beck W, McKechnie J, Benchimol A, Desser K. Coronary heart disease in marathon runners. *Ann NY Acad Sci* 1977;301:593-619.
- 10 Noakes TD, Opie LH, Rose AG and Kleynhans PHT. Autopsy-proved coronary atherosclerosis in marathon runners. *N Engl J Med* 1979;301:86-9.
- 11 New York City marathon. The 25th running. *New York Times* 1994 Nov 7;sect D:6.
- 12 Office of Population Censuses and Surveys. *OPCS mortality statistics. Cause. England and Wales, 1992*. London: HMSO, 1993.
- 13 Curfman GD. Is exercise beneficial—or hazardous—to your heart? *N Engl J Med* 1993;329:1730-1.
- 14 Pate RR, Pratt M, Blair SN, Haskell WL, Macera CA, Bouchard C, *et al*. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* 1995;273:402-7.
- 15 Davies MJ. Acute coronary thrombosis—the role of plaque disruption and its initiation and prevention. *Eur Heart J* 1995;16 (suppl L):3-7.
- 16 Hambrecht R, Niebauer J, Marburger C, Grunze M, Kälberer B, Hauer K, *et al*. Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *JACC* 1993;22:468-77.
- 17 Northcote RJ, Evans ADB, Ballantyne D. Sudden deaths in squash players. *Lancet* 1984;i:148-50.
- 18 Northcote RJ, Ballantyne D. Cardiovascular implications of strenuous exercise. *Int J Cardiol* 1985;8:3-12.
- 19 Brady HR, Kinirons M, Lynch T, Ohman EM, Tormey W, O' Malley KM, *et al*. Heart rate and metabolic response to competitive squash in veteran players: identification of risk factors for sudden cardiac death. *Eur Heart J* 1989;10:1029-35.