Antimicrobials for acute otitis media? A review from the International Primary Care Network

Jack Froom, Larry Culpepper, Max Jacobs, Ruut A DeMelker, Larry A Green, Louk van Buchem, Paul Grob, Timothy Heeren

State University of
New York, Stony
Brook, NY 11794,
USA

Jack Froom,
professor of family
medicine

Boston University,
Boston, MA 02118,
USA

Larry Culpepper,
professor of family
medicine

Timothy Heeren,
associate professor of
biostatistics and
epidemiology

University of
Utrecht, 3584 CG
Utrecht,
Netherlands

Max Jacobs,
associate professor of
family medicine

Ruut A DeMelker,
professor of family
medicine

University of
Colorado, Denver,
CO 80220, USA

Larry A Green,
Woodward-Chisholm
chairman of family
medicine

St Elizabeth
Hospital, 500 LC
Tilburg,
Netherlands

Louk van Buchem,
otorhinolaryngologist

University of Surrey,
Guildford, Surrey
GU2 5YG

Paul Grob,
professor of general
practice

continued over

Increasing worldwide resistance of bacteria to antimicrobial drugs is causing a crisis, manifested by higher morbidity, mortality, and costs.¹ In 1992 the Institute of Medicine in the United States warned of the growing threat posed by resistant bacteria,² and in 1994 the Centers for Disease Control in Atlanta initiated a prevention strategy,³ linked to a global plan by the World Health Organisation.⁴ Proposed remedies include development of new antimicrobials, improved sanitation, and educating patients not to ask for antimicrobials when they are not useful and physicians to prescribe them conservatively.⁵

About 30% of British children under the age of 3 visit their general practitioner for acute otitis media each year⁶ and 97% receive antimicrobials.⁷ In America it is the most common reason for outpatient antimicrobial use. Because this use has uncertain benefits, it merits reconsideration.

Increasing resistance to antimicrobial agents has been reported for the three most common bacterial causes of otitis media (*Streptococcus pneumoniae*, *Haemophilus influenzae*, and *Moraxella catarrhalis*), but rates differ between countries.⁸⁻⁹ In England and Wales in 1990-5, resistance to penicillin by *S pneumoniae* increased from 1.5% to 3.9% and to erythromycin from 2.8% to 8.6%.¹⁰

Except in the Netherlands, antimicrobials are standard treatment for acute otitis media in most developed countries.⁷⁻¹¹ Although type and duration vary between countries, generally ampicillin, amoxycillin, or co-amoxiclav (amoxycillin-clavulanate) are preferred, with co-trimoxazole a low cost alternative.⁷ In the Netherlands, treatment of symptoms without antimicrobials has been adopted as routine initial treatment for otitis media,¹² and this policy is associated with decreased emergence of resistance among organisms commonly found in otitis media.¹³⁻¹⁴ This approach also is being adopted in Iceland.¹⁵

Does treatment improve outcomes?

Seven randomised blinded studies have compared antimicrobials with placebo in patients with acute otitis media (table 1).¹⁶⁻²³ Comparisons are difficult because inclusion criteria, outcome measures, duration of follow up, and the period when studies were conducted are different. Also some studies had insufficient power

Summary points

Bacterial resistance to antimicrobials is responsible for increasing morbidity, mortality, and costs

The most frequent use of antimicrobials in the United States is for otitis media

Evidence from randomised, placebo controlled trials that routine use of antimicrobials decreases the duration and severity of symptoms and prevents complications is weak

Treatment of acute otitis media differs worldwide, and careful use of antimicrobials in the Netherlands and Iceland seems to have reduced rates of resistance among organisms without compromising outcomes

The management of acute otitis media needs reassessing

to evaluate small to moderate differences. Nevertheless, since results are mixed and no study found large differences between placebo and antimicrobial groups, we conclude that the benefit of routine antimicrobial use for otitis media, judged by either short or long term outcomes, is unproved.

A recent meta-analysis concluded that one child out of seven benefits from treatment,²⁴ but only four of the 33 studies included had placebo or no medication groups,¹⁶⁻¹⁹⁻²¹⁻²⁵ and no justification was given for excluding several methodologically sound placebo controlled studies.¹⁸⁻²⁰⁻²² In contrast, after reviewing 50 trials, Claessen et al judged that methodological flaws precluded making recommendations.²⁶

Does treatment prevent complications?

Although preventing mastoiditis and meningitis is a rationale for antimicrobial treatment, little evidence exists that routine treatment is effective for this purpose. Mastoiditis, a commonly reported complication of otitis media in the preantimicrobial era, is now

Questions related to the routine use of antimicrobials for acute otitis media

- Are the course of and outcomes after acute otitis media improved by antimicrobial treatment?
- Do antimicrobials prevent complications of acute otitis media?
- Which children are at high risk of poor outcomes after acute otitis media? Do they benefit from antimicrobials?
- If antimicrobials are used, are the optimum type and duration of administration known?
- Have management options, other than routine antimicrobial administration, been widely used and with what results?
- Does decreased use of antimicrobials for acute otitis media influence the development of antimicrobial resistance?

uncommon. It is uncertain, however, whether the decrease is related to antimicrobial treatment, changes in clinical course and organism virulence, or increased host resistance. Also, uncomplicated otitis media often might not have come to medical attention in the era before antimicrobials, increasing the relative rate of reported complications.

Selective use of antimicrobials might yield the same benefit as routine use. In the Netherlands, among 4860 consecutive patients with acute otitis media who were not given antimicrobials, only two experienced mastoiditis; both responded to treatment with oral antimicrobial as outpatients.²⁷ Justification for routine antimicrobial treatment of otitis media to prevent meningitis is equally weak. None of the 4860 patients developed meningitis. Kilpi et al reported that positive blood cultures in children with bacterial meningitis were equally common in those treated and not treated with antimicrobials before admission (77% and 78%).²⁸ Even when otitis media and meningitis coexist, they are

likely to result from a common upper respiratory infection with haematogenous spread, rather than direct spread from ear to meninges.²⁹ It therefore remains uncertain whether either routine or selective antimicrobial treatment prevents the complications of mastoiditis and meningitis.

Do children at high risk benefit from antimicrobials?

Adverse consequences of otitis media include lack of bacteriological or clinical resolution, recurrent infection, persistent effusion, hearing loss, adenoidectomy, and insertion of a tympanostomy tube. No study has addressed whether antimicrobial treatment decreases the frequency of these in all or some of those with known risk factors.

The most important risk factors for poor outcome are young age and attendance at day care centres. In a study of two groups of children infected with *S pneumoniae*, comparable by sex, age at first attack, and frequency of earlier attacks, more penicillin resistant strains were isolated in those under 18 months ($P=0.003$).³⁰ Decreased rates of bacteriological or clinical resolution^{7 22 31} and increased rates of recurrence,³² adenoidectomy, and insertion of a tympanostomy tube³³ are other consequences reported in children under 2 years old.

Day care is a risk factor for poor outcomes, including recurrent infection,³⁴ admission to hospital, adenoidectomy, and insertion of a tympanostomy tube.³³ After other risk factors (age <2 years, white race, male sex, and history of tonsillitis, enlarged adenoids, or asthma) were controlled for, children in day care had a 50% higher chance of repeated ear infections than those not in day care.³⁵ In a nine country study, children in day care were more likely to have a history of poor hearing, tympanostomy tubes, tonsil-

Correspondence to: Professor Larry Culpepper, Department of Family Medicine, Dowling 5 South, 1 Boston Medical Center Place, Boston, MA 02118-2393, USA.

Table 1 Randomised double blind placebo controlled trials of antimicrobial treatment for acute otitis media

First author	Antimicrobials	No of subjects	Age (years)	Outcome intervals	Outcome measures	Results
Halsted (1968) ¹⁶	Ampicillin Penicillin-sulphamethoxazole	89	2 months- 66 months	24-72 hours, 14-18 days, 3 months	Symptomatic improvement, bacterial cure, recurrence	Increased treatment failure at 24-72 hours in placebo group (18.5% v 4.8%, $P<0.05$); no other differences
Howie (1972) ¹⁷	Erythromycin, triple sulphonamide, erythromycin-triple sulphonamide, ampicillin	280	0-2.5	2-7 days	Effusion, reduction in pathogens	No differences: all placebo group became afebrile and asymptomatic, though more produced positive cultures on tympanocentesis
Van Buchem (1981) ¹⁸	Ampicillin, ampicillin plus myringotomy, and myringotomy	171	2-12	2 and 7 days, 1 and 2 months, 1 and 2 years	Pain, fever, duration of discharge, otoscopy, hearing, recurrence, complications	No differences
Mygind (1981) ¹⁹	Penicillin	149	1-10	2-3 days, 7 days, 1 and 3 months	Pain, fever, analgesic use, discharge, otoscopy, tympanometry	Antimicrobials reduced pain on day 2. No other differences
Kaleida (1991) ²⁰	Amoxycillin, amoxycillin plus myringotomy, and myringotomy	536	2-12	2-3 days, 2 and 6 weeks, monthly to 1 year	Initial treatment failure (pain or fever), recurrence, effusion	In non-severe cases increased initial treatment failure (persistent fever $\geq 38^\circ$ orally or severe pain after 48 hours) in placebo group (7.7% v 3.9%) and increased effusion at 2 weeks; no other differences. In severe cases increased initial treatment failure in placebo group (23.5% v 4.1% and 3.1%), but at 6 weeks no differences in effusion or recurrence
Burke (1991) ²¹	Amoxycillin	232	3-10	1-8 days, 1-3 months, 1 year	Pain, crying, fever, school absence, effusion, recurrences	Initial response better in antimicrobial group (pain disappeared by day 3 in 75% of placebo group v 82% of antimicrobial group; treatment failures at one week 14.7% of placebo group v 1.7% of antimicrobial group). No differences after 1 week
Appelman (1991) ²² , Claessen (1994) ²³	Co-amoxiclav	121	0.5-12	3 days, 1 year	Pain, fever	No differences up to 1 year

lectomy, or adenoidectomy and to be referred to an otolaryngologist at the initial visit.³⁶

Other factors likely to contribute to poor recovery include multiple previous episodes,³⁷ bottle feeding,³⁸ history of ear infections in parents or siblings,³⁹ and use of a dummy.⁴⁰ Although exposure to tobacco smoke has been considered a contributor to poor outcome, the evidence is controversial and the effect weak at best.⁴¹ Although risk factors are known, no study has shown that antimicrobial treatment improves outcome in children at risk.

What is the optimal type and duration of administration?

In the nine country study, antimicrobials did not improve outcome at two months, and no differences in rates of recovery were found for either antimicrobial type or duration.⁷ The optimum duration of antimicrobial treatment for otitis media is uncertain. There is evidence that two days,⁴² three days,⁴³ five days,⁴⁴ and 10 days are equally effective. Treatment for more than a few days might have little local effect. In a study of antimicrobial penetrance of the middle ear, penicillin concentrations decreased by about 70% after the second day of treatment (compared with the first day), suggesting that penetration depends on inflammation.⁴⁵

No compelling evidence

After addressing these four questions, we conclude that existing research offers no compelling evidence that children with acute otitis media routinely given antimicrobials have a shorter duration of symptoms, fewer recurrences, or better long term outcomes than those who do not receive them. It also is not clear that routine compared with selective use of antimicrobials prevents complications. Thus it is prudent to reconsider routine use of antimicrobials for otitis media and to consider other approaches.

What are the effects of other management options?

Dutch family physicians use antimicrobials for upper respiratory tract infections more conservatively than doctors in other countries (see box).^{18 27} A World Health Organisation study of tonsillitis in 17 European countries found that 68% of Dutch patients received antimicrobials compared with 94% in other countries.⁴⁶ In 1990 the Dutch College of General Practitioners adopted a guideline for treating acute otitis media. It has received widespread use in the Netherlands, but its adoption elsewhere requires consideration of the current healthcare system, particularly the ability of parents to access care for their children initially and if they fail to improve after treatment of symptoms.

Has this guideline resulted in more complications in Dutch children? While no controlled study has addressed this question, van Buchem et al found that only 2.7% of 4860 children given symptomatic treatment and no antimicrobial drug developed persistent fever, pain, or persistent discharge after three to four days and only two developed mastoiditis.²⁷ Compared with cases in seven other countries where antimicrobial therapy is virtually universal, Dutch

Dutch guideline for the treatment of acute otitis media^{*12}

Patients 2 years and older

- Treatment of symptoms only (paracetamol with or without decongestant nose drops) for the first three days
- Re-evaluation if symptoms (pain or fever thought to be due to acute otitis media) continue for three days. At that time the doctor may continue additional observation or give an antimicrobial (amoxycillin, or erythromycin if amoxycillin is contraindicated) for seven days
- Special treatment for tympanic membrane perforation is not suggested unless it persists for 14 days, at which time a course of antimicrobials is suggested

Children between the ages of 6 months and 2 years

- Management is the same as for those 2 years and older, except for a mandatory contact (either telephone or visit) after 24 hours. If there is no improvement doctors may either start antimicrobials or wait an additional 24 hours
- Referral to an otolaryngologist suggested if patients in this age group appear to be seriously ill or do not improve after 24 hours of treatment with antimicrobials

*An English translation of the guideline is available from: The Dutch College of General Practitioners, Domus Medica, PO Box 323, 3502 GE Utrecht, Netherlands

patients had similar outcomes at two months.⁷ Thus Dutch experience suggests feasible alternatives to antimicrobial treatment exist.

What is the effect on antimicrobial resistance?

Antimicrobial use in children with otitis media results in the emergence of resistant organisms in those children and in the community. Among children previously given antimicrobials for otitis media, Harrison et al found a rate of ampicillin resistance in bacteria in effusions obtained during a subsequent bout of otitis media that was three times higher than normal.⁴⁷ Significant increases in resistant strains of *H influenzae*, *M catarrhalis*,⁴⁸ and *S pneumoniae*⁴⁹ have been reported among children with otitis media who have had previous antimicrobial treatment. Development and spread of multiply resistant pneumococci after treatment for otitis media have been documented in day care centres and surrounding communities,⁵⁰ including instances that have led to deaths from meningitis in children treated previously for uncomplicated acute otitis media.⁵¹

Although the organisms that cause otitis media are similar across countries,⁵² the Netherlands has a lower prevalence of resistant strains than other European countries. A study in 1989 in the Netherlands found that only 3% of 1100 *S pneumoniae* cultures showed decreased sensitivity to penicillin compared with more than 10% in most countries.^{14 53} Similarly, 4.7% of non-type B *H influenzae* isolates from respiratory tract infections were resistant to amoxycillin—less than the 10% overall rate in Europe.¹³

In Iceland, penicillin resistant pneumococci appeared in 1988 and increased to nearly 20% of pneumococcal infections by 1993.⁵⁴ After a campaign to decrease antimicrobial use, particularly for otitis media, and to control spread of infections in day care centres, this rate declined to 16.9% in 1994. The presence of resistant pneumococci among carriers who are day care attendees declined from 20% (21 of 104) in 1992 to 15% (19 of 126) in 1995.¹⁵ Nevertheless, it is not possible to state with certainty that decreased use of antimicrobials results in decreased resistance in endemic bacteria.

What needs to be done?

Research—Placebo studies indicate that more than 80% of children with acute otitis media recover without antimicrobials. Antimicrobials might be useful for some of the remainder, but evidence of benefit is lacking. A trial should be designed to study antimicrobial effectiveness in children at high risk for poor outcomes, using carefully chosen outcome measures, with sufficient numbers of patients for subgroup analyses.

Prevention of acute otitis media—Sufficient information exists to support aggressive encouragement of breast feeding and avoidance of tobacco smoke.^{38 55} There is insufficient evidence of effectiveness to recommend pneumococcal vaccine.⁵⁶ Over 70% of cases of acute otitis media are immediately preceded by viral respiratory infections,³¹ and there is convincing evidence that contaminated hands and surfaces are more important than airborne respiratory droplets in viral transmission.^{57 58} Campaigns to encourage day care centres to increase hand washing by staff and children and the use of virucidal agents to wipe surfaces might reduce viral transmission and subsequent otitis media.

Treatment of acute otitis media—Clinicians should immediately reconsider the routine use of antimicrobials for children with otitis media and consider treating symptoms with analgesics and observation for lack of improvement. Children less than 2 years old require frequent re-evaluation if not treated. Doctors are often uncertain about the diagnosis of otitis media⁷; in such circumstances in particular antimicrobials should be withheld. If antimicrobials are used they should be given for less than 10 days. In addition doctors should identify modifiable risks for poor outcome and work with parents to eliminate them. The potential benefits of these actions are profound. They could reduce costs without compromising outcomes while reducing the risk of bacterial resistance.

This paper is dedicated to the memory of Dr Max Jacobs, whose intellect, wit, and ability to solve problems internationally made our collaboration possible. He died suddenly on 21 December 1996.

Funding: Agency for Health Care Policy and Research, Grant No. RO1 HS07035-03.

Conflict of interest: None.

1 Neu HC. The crisis in antibiotic resistance. *Science* 1992;257:1036-8.

2 Lederberg J, Shope RE, Oaks SC, eds. *Emerging infections: microbial threats to health in the United States*. Washington, DC: Institute of Medicine, National Academy Press, 1992.

3 Centers for Disease Control and Prevention. *Addressing emerging infectious disease threats*. Atlanta: US Department of Health and Human Services, Public Health Service, 1994.

4 LeDuc JW. World Health Organization strategy for emerging infectious diseases. *JAMA* 1996;275:318-20.

- 5 Cohen ML. Epidemiology of drug resistance: implications for a post-antimicrobial era. *Science* 1992;257:1050-5.
- 6 Ross AK, Croft PR, Collins M. Incidence of acute otitis media in infants in a general practice. *J R Coll Gen Pract* 1988;38:70-2.
- 7 Froom J, Culpepper L, Grob P, Barteld A, Bowers P, Bridges-Webb C, et al. Diagnosis and antibiotic treatment of acute otitis media: report from International Primary Care Network. *BMJ* 1990;300:582-6.
- 8 Faden H, Doern G, Wolf J, Blocker M. Antimicrobial susceptibility of nasopharyngeal isolates of potential pathogens recovered from infants before antibiotic therapy: implications for the management of otitis media. *Pediatr Infect Dis J* 1994;13:609-12.
- 9 Henderson KW, Gilligan PH, Wait K, Goff DA. Nasopharyngeal carriage of antibiotic-resistant pneumococci by children in group day care. *J Infect Dis* 1988;157:256-63.
- 10 Johnson AP, Speller DCE, George RC, Warner M, Domingue G, Efstratiou A. Prevalence of antibiotic resistance to pneumococci in England and Wales: results of observational surveys in 1990 and 1995. *BMJ* 1996;312:1454-6.
- 11 De Melker RA, Kuyvenhoven MM. Management of upper respiratory infection in Dutch general practice. *Br J Gen Pract* 1991;41:504-7.
- 12 Appelman CLM, Bossen PC, Dunk JHM, van de Lisdonk EH, de Melker RA, van Weert HCPM. *Guideline. Dutch College of Family Doctors. Acute otitis media*. Utrecht: Dutch College of Family Doctors, 1990.
- 13 Machka K, Braveny I, Dabernat H, Dombusch K, van Dyck E, Kayser FH, et al. Distribution and resistance patterns of *Haemophilus influenzae*: a European Cooperative study. *Eur J Clin Microbiol Infect Dis* 1988;7:14-24.
- 14 De Neeling AJ, Hemmes JH, van Klingeren B. *Resistente tegen antibiotica bij routine isolaten van bacteriën in zeven streeklaboratoria* (English abstract). Bilthoven: Rijksinstituut Voor Volksgezondheid en Milieuhygiëne, 1993 (report No 253601001).
- 15 Stephenson J. Icelandic researchers are showing the way to bring down rates of antibiotic-resistant bacteria. *JAMA* 1996;275:175-6.
- 16 Halsted C, Lepow ML, Balassanian N, Emmerich J, Wolinsky E. Otitis media: clinical observations, microbiology, and evaluation of therapy. *Am J Dis Child* 1968;115:542-51.
- 17 Howie VM, Ploussard JH. Effectiveness of erythromycin estolate, triple sulfonamide, ampicillin, erythromycin estolate-triple sulfonamide, and placebo in 280 patients with acute otitis media under 2 and one-half years of age. *Clin Pediatr* 1972;11:205-14.
- 18 van Buchem FL, Dunk JHM, van't Hof MA. Therapy of acute otitis media: myringotomy, antibiotics, or neither: a double-blind study in children. *Lancet* 1981;ii:883-7.
- 19 Mygind N, Meistrup-Larsen KI, Thomsen J, Thomsen VF, Josefsson K, Sorensen H. Penicillin in acute otitis media: a double-blind, placebo-controlled trial. *Clin Otolaryngol* 1981;6:5-13.
- 20 Kaleida PH, Casselbrant ML, Rockette HE, Paradise JL, Bluestone CD, Blatter MM, et al. Amoxicillin or myringotomy or both for acute otitis media: results of a randomized clinical trial. *Pediatrics* 1991;87:466-74.
- 21 Burke P, Bain J, Robinson D, Dunleavy J. Acute red ear in children: controlled trial of non-antibiotic treatment in general practice. *BMJ* 1991;303:558-62.
- 22 Appelman CLM, Claessen JQPJ, Touw-Otten FWMM, Hordijk GJ, DeMelker RA. Co-amoxiclav in recurrent acute otitis media: placebo controlled study. *BMJ* 1991;303:1450-2.
- 23 Claessen JQPJ, Appelman CLM, Touw-Otten FWMM, de Melker RA, Hordijk GJ. Persistence of middle ear dysfunction after recurrent acute otitis media. *Clin Otolaryngol* 1994;19:35-40.
- 24 Rosenfeld RM, Vertrees JE, Carr J, Cipolle RJ, Uden DL, Giebink GS, et al. Clinical efficacy of antimicrobial drugs for acute otitis media: meta-analysis of 5400 children from thirty-three randomized trials. *J Pediatrics* 1994;124:355-67.
- 25 Laxdal OE, Merida J, Trefor Jones RH. Treatment of acute otitis media: a controlled study of 142 children. *Can Med Assoc J* 1970;102:263-8.
- 26 Claessen JQPJ, Appelman CLM, Touw-Otten FWMM, de Melker RA, Hordijk GJ. A review of clinical trials regarding treatment of acute otitis media. *Clin Otolaryngol* 1992;17:251-7.
- 27 Van Buchem FL, Peeters ME, van't Hof MA. Acute otitis media: a new treatment strategy. *BMJ* 1985;290:1033-7.
- 28 Kilpi T, Anttila M, Kallio MJ, Peltola H. Severity of childhood bacterial meningitis: duration of illness before diagnosis. *Lancet* 1991;338:406-9.
- 29 Eavey RD, Gao Y-Z, Schuknecht HF, Gonzalez-Pineda M. Otolologic features of bacterial meningitis of childhood. *Pediatrics* 1985;106(3):402-7.
- 30 Barry B, Gehanno P, Blumen M, Boucot I. Clinical outcome of acute otitis media caused by pneumococci with decreased susceptibility to penicillin. *Scand J Infect Dis* 1994;26:446-52.
- 31 Froom J, Culpepper L, Grob P, Barteld A, Bowers P, Bridges-Webb C, et al. Effect of patient characteristics and disease manifestations on the outcome of acute otitis media at two months. *Arch Fam Med* 1993;2:841-6.
- 32 Rasmussen F. Recurrence of acute otitis media at preschool age in Sweden. *J Epidemiol Community Health* 1994;48:33-5.
- 33 Alho OP, Koivu M, Sorri M, Oja H, Kilkku O. Which children are being operated on for recurrent acute otitis media? *Arch Otolaryngol Head Neck Surg* 1994;120:807-11.
- 34 Louhiala PJ, Jaakkola N, Ruotsalainen R, Jaakkola JJ. Form of day care and respiratory infection among Finnish children. *Am J Public Health* 1995;85:1109-12.
- 35 Hardy AM, Fowler MG. Child care arrangements and repeated ear infections in young children. *Am J Public Health* 1993;83:1321-5.
- 36 Froom J, Culpepper L. Otitis media in day care children: a report from the International Primary Care Network. *J Fam Pract* 1991;32:289-94.
- 37 Alho OP, Oja H, Koivu M, Sorri M. Risk factors for chronic otitis media with effusion in infancy. Each acute otitis media episode induces a high but transient risk. *Arch Otolaryngol Head Neck Surg* 1995;121:839-43.
- 38 Sassen ML, Brand R, Grote JJ. Breast-feeding and acute otitis media. *Am J Otolaryngol* 1994;15:351-7.

39 Teele DW, Klein JO, Rosner B. Epidemiology of otitis media during the first seven years of life in children in greater Boston: a prospective cohort study. *J Infect Dis* 1989;160:83-94.

40 Niemela M, Uhari M, Mottonen M. A pacifier increases the risk of recurrent acute otitis media in children in day care centers. *Pediatrics* 1995;96:884-8.

41 Blakley BW, Blakley JE. Smoking and middle ear disease: are they related? A review article. *Otolaryngol Head Neck Surg* 1995;112:441-6.

42 Meistrup-Larsen KI, Sorensen H, Johnsen NJ, Thomsen J, Mygind N, Sederberg-Olsen J. Two versus seven days penicillin treatment for acute otitis media. *Acta Otolaryngol* 1983;96:99-104.

43 Chaput de Saintonge DM, Levine DF, Savage IT, Gurgess GWS. Trial of three-day and ten-day course of amoxycillin in otitis media. *BMJ* 1982;284:1078-81.

44 Hendrickse WA, Kusmiesz H, Shelton S, Nelson JD. Five vs. ten days of therapy for acute otitis media. *Pediatr Infect Dis J* 1988;7:17-23.

45 Lundgren K, Ingvarsson L, Rudcrantz H. The concentration of penicillin-V in middle ear exudate. *J Pediatr Otorhinolaryngol* 1979;1:93-6.

46 Touw-Otten FW, Johansen KS. Diagnosis, antibiotic treatment and outcome of acute tonsillitis: report of a WHO regional office for Europe study in 17 European countries. *Fam Pract* 1992;9:255-64.

47 Harrison CJ, Marks MI, Welch DF. Microbiology of recently treated acute otitis media compared with previously untreated acute otitis media. *Pediatr Infect Dis J* 1985;4:641-6.

48 Faden H, Bernstein J, Stanievich J, Brodsky L, Ogra PL. Effect of prior antibiotic treatment on middle ear disease in children. *Ann Otol Rhinol Laryngol* 1992;101:87-91.

49 Ford KL, Mason EO Jr, Kaplan SL, Lamberth LB, Tillman J. Factors associated with middle ear isolates of *Streptococcus pneumoniae* resistant to penicillin in a children's hospital. *J Pediatrics* 1991;119:941-4.

50 Reichler MR, Alphin AA, Breiman RF, Schreiber JR, Arnold JE, McDougal LK, et al. The spread of multiply resistant *Streptococcus pneumoniae* at a day care center in Ohio. *J Infect Dis* 1992;166:1346-53.

51 Sloas MM, Barrett FF, Chesney PJ, English BK, Hill BC, TenoverFC, et al. Cephalosporin treatment failure in penicillin- and cephalosporin-resistant *Streptococcus pneumoniae* meningitis. *Pediatr Infect Dis J* 1992;11:662-6.

52 Bluestone CD, Klein JO. *Otitis media in infants and children*. Philadelphia: WB Saunders Company, 1988:45-8.

53 Van Klinger B, Michelen MF, Wagenvoort JHT. Beheersing van het resistentievraagstuk door het voeren van een antibioticum beleid. *Ned Tijdschr Geneesk* 1992;136:860-4.

54 Aarason VA, Kristinson KG, Sigurdsson JA, Stefansdottir G, Molstad S. Do antimicrobials increase carriage rate of penicillin resistant pneumococci in children? *BMJ* 1996;313:387-91.

55 Stenstrom R, Bernard PAM, Ben-Simhon H. Exposure to environmental tobacco smoke as a risk factor for recurrent acute otitis media in children under the age of five years. *Int J Pediatr Otorhinolaryngol* 1993;27:127-36.

56 Lee CJ, Wang TR. Pneumococcal infection and immunization in children. *Crit Rev Microbiol* 1994;20:1-12.

57 Gwaltney JM, Hendley JO. Transmission of experimental rhinovirus infection by contaminated surface. *Am J Epidemiol* 1982;116:828-33.

58 Murray BE. Can antibiotic resistance be controlled? *N Engl J Med* 1994;330:1229-30.

(Accepted 28 May 1997)

Feasibility of screening toddlers for iron deficiency anaemia in general practice

John A James, Gabrielle J Laing, Stuart Logan, Michael Rosedale

Montpelier Health Centre, Bristol BS6 5PT
John A James, general practitioner

Department of Child Health, St Leonard's, London N1 5LZ
Gabrielle J Laing, consultant in community paediatrics

Department of Epidemiology and Biostatistics, Institute of Child Health, London WC1N 1EH
Stuart Logan, senior lecturer in paediatric epidemiology

111 Pembroke Road, Bristol BS8 3EU
Michael Rosedale, general practitioner

Correspondence to: Dr James.

BMJ 1997;315:102-3

Iron deficiency is the commonest cause of anaemia in children in the United Kingdom and is reported to be associated with psychomotor delay; it is reversible by iron treatment.^{1,2} Screening for iron deficiency anaemia in toddlers could become a routine part of child health promotion.³

Screening at 13-14 months was introduced in an inner city practice (by JAJ) in 1987. Coverage exceeds 90%.⁴ We examined the feasibility of extending screening to other practices in the same city. Practices 1 (JAJ) and 3 serve deprived inner city communities with half of the population from ethnic minority groups; practice 2 serves a relatively affluent population; practices 4 and 5 have largely working class populations.

Subjects, methods, and results

The screening programme was agreed between JAJ and general practitioners from four other practices who expressed an interest in the programme. Screening for iron deficiency anaemia (haemoglobin concentration < 110 g/l, based on estimation by a

single laboratory) was offered to all children attending for routine measles, mumps, and rubella immunisation. All practice nurses were trained in thumb prick capillary blood sampling, and a sample was taken from children whose parents gave consent.

Numbers of children offered screening, parents refusing, samples on which estimations could not be performed (sample insufficient or clotted), and anaemic children were recorded for each practice. Screening continued for up to 24 months; practice 3 abandoned the project at six months because of time constraints, parental objections, and technical difficulties in obtaining blood samples.

Population coverage for each practice was estimated by multiplying the proportion of children offered the screen in whom a haemoglobin concentration was obtained by the immunisation coverage for that practice. Up to 32% of parents refused screening, and the a haemoglobin concentration was obtained in 39-100% of those offered screening. The estimated prevalence of anaemia was 23% overall (range 0-35%) (table 1).

Table 1 Toddlers offered and undergoing screening for iron deficiency anaemia and numbers who were anaemic. Values are numbers (percentages) unless indicated otherwise

Practice	Offered screening	Refusal	Sample inadequate	Haemoglobin concentration obtained	Coverage of MMR vaccine	Estimated coverage of population	No anaemic (%; 95% CI)
1	125	0	0	125 (100)	125/136 (92)	125/136 (92)	44 (35; 27 to 44)
2	170	26 (15)	0	144 (85)	170/177 (96)	144/177 (81)	22 (15; 10 to 22)
3	56	18 (32)	16 (29)	22 (39)	56/84 (67)	22/84 (26)	7 (32; 14 to 55)
4	68	5 (7)	3 (4)	60 (88)	68/97 (70)	60/97 (62)	17 (23; 18 to 41)
5	41	0	0	41 (100)	41/45 (91)	41/45 (91)	0 (0; 0 to 10)
All practices	460	49 (11)	19 (5)	392 (85)	460/539 (85)	392/539 (73)	90 (23; 19 to 28)

MMR=measles, mumps, and rubella.

Forty nine parents (11%) declined screening. Half felt the test was not necessary; others considered the test traumatic, and one quarter said they would "think about the test" and come back. None did.

Comment

Iron deficiency anaemia is common in toddlers, even in the more affluent population of practice 3 (prevalence 15.3%). The screening programme was successful in only one of the four new practices. Two practices achieved reasonable proportions of haemoglobin estimations, although in one with low uptake of immunisation the estimated population coverage rate was only around 70%. In the fourth practice, high rates of parental refusal and failure to obtain analysable samples led to abandonment of the programme after six months.

All four practices provide child health promotion for their patients and entered the study as eager volunteers. A successful screening programme must have high coverage; the population not screened generally has higher rates of abnormality. Although enthusiasts have reported successful screening programmes for iron deficiency anaemia in primary care, relatively poor coverage was achieved in three practices in spite of their commitment (practices 1, 2, and 3 together obtained haemoglobin concentrations in 77% of children offered screening, an estimated population coverage of 63%). An attempt to introduce screening

for iron deficiency anaemia into community clinics in a deprived London district was even less successful, with an uptake of under 10% in nearly 600 children invited for screening (GJ Laing, unpublished data).

Success would require that health professionals and parents were convinced of the benefits of detecting and treating iron deficiency anaemia; that staff were well trained and experienced; that adequate time was available; and that high uptake of measles, mumps, and rubella vaccine was maintained. The difficulties identified, in combination with the fluctuating nature of iron deficiency in this age group,³ suggest that population screening for iron deficiency anaemia is unlikely to be an appropriate strategy. Population based interventions may prove more effective.

Funding: South West Research and Development NHS Funding.

Conflict of interest: None.

- 1 Idjradinata P, Pollitt E. Reversal of developmental delays in iron deficient infants treated with iron. *Lancet* 1993;341:1-4.
- 2 Aukett MA, Parks YA, Scott PH, Wharton BA. Treatment with iron increases weight gain and psychomotor development. *Arch Dis Child* 1986;61:849-57.
- 3 Hall DM. *Health for all children*. 3rd ed. Oxford: Oxford University Press, 1996.
- 4 James JA, Lawson P, Male P, Oakhill A. Preventing iron deficiency in pre-school children by implementing an educational and screening programme in an inner city practice. *BMJ* 1989;299:838-40.
- 5 James J, Laing GJ, Logan S. Changing patterns of iron deficiency anaemia in the second year of life. *BMJ* 1995;311:230.

(Accepted 16 January 1997)

A talk that changed my practice My latterday Demosthenes

Bernard Dixon's article—"Sorry, you won't be able to see this ..." (*BMJ* 1996;313:1407) had a familiar ring. My Road to Damascus came at a meeting such as his. We had dozed through the mutterings of serial grim speakers when a young lady took the platform. Her talk was understandable and to the point, her slides simple, her delivery polished yet relaxed. The effect was electric, instantly piercing our postprandial stupor.

Her influence on me was even more profound. Here was someone who was either a "natural speaker," unlikely I thought, or one who had taken the trouble to acquire effective platform technique. The next week I found myself signing up as a beginner at the local school of public speaking. My knees were like jelly, but I need not have worried, so were those of the other novices.

All over again, I learnt about the muscles of respiration, only this time how to control them rather than how to describe them. The paranasal sinuses became resonators each adding its own characteristic timbre to the voice, and the chest became the biggest resonator of all, lending power to the delivery. Physiology became real too: how to control the feelings of fright and flight, how nervous tension affects voice production and what to do about it. By the end of term, I was deemed ready for the first test, the London Academy of Music and Dramatic Arts bronze medal in public speaking. On the day, with increasing dryness of the mouth, tightness of the chest, and weakness of the knees, terror knew no bounds. I was back as an undergraduate facing my first viva voce. I had great difficulty in silencing the Greek chorus situated somewhere behind my left ear which insisted

that this was optional and what I really needed was committal to a lunatic asylum. But I managed to deliver my two prepared short contrasting speeches to an imaginary audience and even to do the impromptu speech and sight reading some (rough) justice. The examiner was satisfied, and although I had vowed as I went in, "never more," I found myself continuing up the ladder, finally, some three years later, reaching the pinnacle, the Associate Diploma in Public Speaking. What next? The London Guildhall teaching diploma is beckoning.

What have I gained? I have acquired confidence in communication skills, not only in the mechanics of voice production and platform technique but also in planning, construction, and delivery of presentations. I have learnt the effective use of visual aids and how to use a microphone. I have learnt to keep in touch with the audience and to enlist the help of colleagues when faced with daunting assignments. Above all, I have learnt to enjoy the challenge.

During this time I have rubbed shoulders with architects, accountants, archivists, actresses, and bishops. Well, actually a church warden, but why spoil a good story for want of the truth? I have found time to return to literature, enjoying anew the magic of the written and spoken word. I have my latterday Demosthenes to thank for this, for her demonstration that the best speakers are those who combine ability with training. Who was she? I have no idea, but I thank her all the same.

Christine Waddell, *associate specialist in cytopathology, Birmingham*