How has fundholding in Northern Ireland affected prescribing patterns? A longitudinal study

Thérèse Rafferty, Keith Wilson-Davis, Hugh McGavock

See editorial by Keeley
and p 170

Drug Utilisation
Research Unit,
Department of
Therapeutics and
Pharmacology,
Queen's University
of Belfast, Belfast
BT9 7BL

Thérèse Rafferty,
computer officer
Keith Wilson-Davis,
deputy director
Hugh McGavock,
director

Correspondence to:
Miss Rafferty.

BMJ 1997;315:166-70

Abstract

Objective: To compare prescribing patterns in general practices before and after the introduction of fundholding in April 1993 to determine whether fundholding changed prescribing patterns among practices that joined the scheme.

Design: Analysis of prescribing data from the Drug Utilisation Research Unit's database for all practices in Northern Ireland during April 1989 to March 1996.

Setting: Northern Ireland.

Subjects: 23 first wave fundholders, 34 second wave fundholders, 9 third wave fundholders, and 268 non-fundholders.

Main outcome measures: Prescribing costs per 1000 patients, prescription items per 1000 patients, average cost per item, and rate of generic prescribing.

Results: Prescribing costs and frequency increased in all groups throughout the study. Among the fundholders the rate of increase in costs after fundholding was significantly lower than among non-fundholders. The rate of increase in cost per item fell, coinciding with a significant increase in the rate of generic prescribing. However, with regard to first wave fundholders, their yearly increase in costs in their third year as fundholders (1995-6) was similar to that of the non-fundholders. The earlier practices that joined the scheme seemed to differ in some important respects from those that joined later.

Conclusions: After fundholders joined the fundholding scheme their patterns of prescribing changed compared with those of non-fundholders: the rate of increase in costs fell and there was a significant rise in the rate of generic prescribing.

Introduction

Containing escalating health care budgets is an international problem. General practitioners' prescribing makes a big contribution to the rising health care budget and various measures have been used to influence their prescribing behaviour. These measures usually provide doctors with information relating to drugs and prescribing patterns. Some provide general information about a particular drug or group of drugs, encouraging restrained use.¹ Others provide prescribers with detailed analysis of their prescribing behaviour with specific instructions for change.² This type of information is sometimes presented to the pre-

scriber by a prescribing adviser, a technique which Soumerai et al found to be the most effective.³ Some countries penalise doctors who fail to comply with guidelines or to prescribe within certain boundaries. In the Netherlands overprescribing can result in non-reimbursement; in France doctors can be fined if they blatantly ignore guidelines. Other countries reward doctors for good practice. In parts of the United States doctors are rewarded with discretionary salary increments for using resources efficiently, and French doctors receive increased fees for complying with guidelines.⁴

In 1991 Britain introduced the fundholding scheme to contain costs. The scheme is voluntary and allows practices to hold budgets for several services including prescribing. Money can be moved between the various elements of the budget. The incentive of the scheme is that the practices are allowed to redirect any money left in the budget at the end of the year back into the practice to be used to benefit patients.⁵ The prescribing bill has the greatest potential for savings, and several studies have shown the effects of this scheme on prescribing costs among general practitioners in Scotland and England. Most show that fundholders can contain costs more effectively than non-fundholders⁶⁻⁷ and that costs are contained by increasing generic prescribing⁷⁻⁸ and decreasing the average cost per item rather than by giving fewer items.⁹ However, these initial achievements may not always be maintained.¹⁰

Fundholding was introduced in Northern Ireland in 1993. First wave fundholders started in April 1993, and further practices joined the scheme in April 1994 and April 1995. This study looks at the effects of fundholding on the prescribing patterns of practices that joined the scheme.

Methods

Our unit's prescribing database records the cost and number of prescriptions written for each preparation by each practice in every quarter. This database is built from data downloaded from the Central Services Agency, whose system was designed for reimbursing pharmacists. We compared the performance of first, second, and third wave fundholders with that of non-fundholders during April 1989 to March 1996 using the following prescribing measures: prescribing cost (£) per 1000 patients, prescription items per 1000

Table 1 Characteristics of fundholding and non-fundholding practices. Values are numbers (percentages) of practices unless stated otherwise

	Dispensing practices	No of partners					No of deprived patients/partner
		1	2	3	4	≥5	
Non-fundholders (n=268)	26 (10)	70 (26)	81 (30)	67 (25)	30 (11)	20 (7)	263
First wave (n=23)	1 (4)	0	1 (4)	4 (17)	7 (30)	11 (48)	177
Second wave (n=34)	3 (9)	3 (9)	8 (24)	9 (26)	5 (15)	9 (26)	200
Third wave (n=9)	1 (11)	0	1 (11)	2 (22)	4 (44)	2 (22)	290

Table 2 Mean cost (£) (mean % increase on previous year) per 1000 patients for fundholders and non-fundholders

	Year						
	89-90	90-1	91-2	92-3	93-4 (First wave)	94-5 (Second wave)	95-6 (Third wave)
Non-fundholders (n=268)	52 622	58 911 (12)	68 828 (17)	77 532 (13)	88 385 (14)	99 173 (12)	110 632 (12)
Fundholders:							
First wave (n=23)	50 701	56 804 (12)	66 404 (17)	72 924 (10)	75 362 (4)**	82 738 (10)**	91 329 (11)**
Second wave (n=34)	49 789	56 662 (15)	66 085 (17)	75 367 (14)	84 348 (12)	87 966 (5)**	94 281 (7)**
Third wave (n=9)	47 335	52 995 (12)	61 984 (17)	70 192 (13)	80 152 (15)	89 727 (13)	93 492 (4)*

*P≤0.01, **P≤0.001 compared with non-fundholders by the multiple comparison *t* test with Bonferroni correction.

Table 3 Mean number of prescription items (mean % increase on previous year) per 1000 patients for fundholders and non-fundholders

	Year						
	89-90	90-1	91-2	92-3	93-4 (First wave)	94-5 (Second wave)	95-6 (Third wave)
Non-fundholders (n=268)	9832	9989 (2)	10 494 (5)	10 791 (3)	11 473 (6)	11 992 (5)	12 455 (4)
Fundholders:							
First wave (n=23)	8550**	8635 (1)***	8 944 (4)***	9 113 (2)***	9 334 (38)***	9 756 (5)***	10 135 (4)***
Second wave (n=34)	9061*	9183 (2)*	9 712 (6)*	10 143 (4)	10 792 (7)	11 093 (3)*	11 482 (4)*
Third wave (n=9)	8407*	8567 (2)*	9 024 (5)*	9 342 (4)*	9 998 (7)*	10 423 (4)*	10 675 (3)*

*P≤0.05, **P≤0.01, ***P≤0.001 compared with non-fundholders by the multiple comparison *t* test with Bonferroni correction.

patients, average cost (£) per prescription item, and rate of generic prescribing.

We included all practices in Northern Ireland except for those that did not exist for the entire study period. Dispensing dentists and hospices were also excluded as no list size is recorded for them. The term "non-fundholder" refers to practices which were not fundholders at any point in the study.

Throughout the study period, practices in Northern Ireland received regular visits from a prescribing adviser. Practices also had equal access to prescribing reports similar to those issued by the Prescription Pricing Authority in England and to the COMPASS report (computerised online monthly prescribing analysed for science and stewardship), which gives specific indications where each practice can make savings.² When the fundholding scheme was introduced in 1993 the indicative prescribing scheme was already in place, but there were no positive or negative incentives for practices to prescribe within their budgets. No other incentives schemes to improve prescribing existed after the introduction of fundholding.

We analysed the data using the *SPSS* for Windows package. Differences were compared with the multiple comparison *t* test procedure with the Bonferroni correction.

Results

Table 1 gives the characteristics of the practices included in the study. List sizes ranged from 449 to 13 243 (mean 4479) among non-fundholders and from 1366 to 15 993 (mean 6887) among fundholders.

Prescribing costs per 1000 patients rose in all groups throughout the study (fig 1). The fundholders had slightly lower costs than the non-fundholders until they joined the fundholding scheme, although the differences were not significant (table 2). After this their costs were significantly lower than those of non-fundholders. The percentage yearly increases in cost per 1000 patients were similar for fundholders and non-fundholders before fundholding, but once practices became fundholding their costs were significantly lower than those for non-fundholders. Although the costs of first wave fundholders remained significantly lower than those of non-fundholders, in the third year of fundholding these practices had a yearly percentage increase similar to non-fundholders (table 2).

Prescribing items per 1000 patients rose in all groups throughout the study (fig 1). First and third wave fundholders prescribed significantly fewer prescription items per 1000 patients than non-fundholders throughout the study (table 3). The second wave fundholders prescribed significantly fewer items per 1000 patients than non-fundholders after becoming fundholders and for three of the five years previously. Fundholders and non-fundholders had similar yearly percentage increases in items per 1000 patients for most of the study, although in their first year of fundholding the first wave's yearly rate of increase was half that of the non-fundholders (table 3). In their second and third year of fundholding the rate of increase in items per 1000 patients among first wave fundholders was the same as for non-fundholders.

Cost per item rose in all groups (fig 1). The yearly figures show that before becoming fundholders the

Fig 1 Mean cost (£)/1000 patients, items/1000 patients, cost (£)/item, and generic percentage for fundholders and non-fundholders for each quarter from April/June 1989 to January/March 1996

first wave had a significantly higher cost per item than non-fundholders, but on joining the scheme their cost per item did not differ significantly from that of the non-fundholders (table 4). Second and third wave fundholders were similar in cost per item to the non-fundholders from 1989 to 1993, but on becoming fundholders their cost per item fell below that of the non-fundholders and the figure was significantly lower for the second wave in their second year as

fundholders. Fundholders had generally similar percentage yearly increases to those of non-fundholders until they joined the scheme, when their yearly increase was notably less than that of the non-fundholders.

Generic prescribing—The percentage of generic prescribing for non-fundholders remained around 25% throughout the study (table 5). The fundholders remained at a steady, though somewhat higher, level until they joined the scheme. On becoming fundholders their rate of generic prescribing significantly increased compared with non-fundholders. Before fundholding was introduced only small increases ($\leq 2\%$) in generic prescribing occurred among all groups. In the year they became fundholders, however, first wave fundholders had a yearly increase of 14%, the second wave had an increase of 11%, and the third wave an increase of 15%.

Discussion

This is one of few studies of prescribing behaviour that examines an almost complete population and includes behaviour for all practices before fundholding began. Our results confirm that fundholding has a significant effect on prescribing patterns. Fundholders contain costs more effectively than non-fundholders, and this change coincides with practices joining the fundholding scheme. The same pattern can be seen in all three waves of fundholding. This cost containment seems to have been achieved partly through an increase in generic prescribing, which is reflected in the cost per item. However, our data cannot show if there were other changes in prescribing patterns such as substitution of less expensive alternative drugs, or if prescriptions were written for shorter periods or for lower doses. A study looking at specific treatment groups using the defined daily dose method would give some indication of what changes occurred.

Differences between fundholders and non-fundholders

Before joining the fundholding scheme first wave fundholders had slightly lower costs per 1000 patients, fewer items per 1000 patients, and a higher cost per item compared with non-fundholders. Second and third wave fundholders had slightly lower costs per 1000 patients, fewer items per 1000 patients, and a similar cost per item compared with non-fundholders. Work in other regions suggests that the characteristics of the first wave fundholders are typical of practices in more affluent areas, which are larger and more organised than those in poorer areas.^{7 11} First wave fundholders were mainly larger practices, with almost

Table 4 Mean cost (£) (mean % increase on previous year) per item for fundholders and non-fundholders

	Year						
	89-90	90-1	91-2	92-3	93-4 (First wave)	94-5 (Second wave)	95-6 (Third wave)
Non-fundholders (n=268)	5.47	6.02 (10)	6.69 (11)	7.32 (10)	7.85 (7)	8.42 (7)	9.04 (8)
Fundholders:							
First wave (n=23)	6.02*	6.68 (11)*	7.52 (13)**	8.08 (8)*	8.18 (1)	8.59 (5)	9.12 (6)
Second wave (n=34)	5.56	6.25 (13)	6.89 (10)	7.54 (9)	7.93 (5)	8.05 (2)	8.34 (4)*
Third wave (n=9)	5.73	6.28 (10)	6.94 (11)	7.60 (10)	8.08 (7)	8.69 (8)	8.83 (2)

* $P \leq 0.01$, ** $P \leq 0.001$ compared with non-fundholders by the multiple comparison *t* test with Bonferroni correction.

Table 5 Mean generic prescribing rate (%) (mean % increase on previous year) for fundholders and non-fundholders

	Year						
	89-90	90-1	91-2	92-3	93-4 (First wave)	94-5 (Second wave)	95-6 (Third wave)
Non-fundholders (n=268)	25.32	24.85 (-2)	24.85 (0)	24.80 (0)	25.13 (2)	25.47 (2)	25.63 (1)
Fundholders:							
First wave (n=23)	27.02	26.53 (-2)	25.93 (-2)	26.42 (2)	29.93 (14)**	31.09 (4)**	31.42 (1)**
Second wave (n=34)	26.20	25.35 (-3)	25.47 (1)	25.51 (0)	26.37 (4)	29.12 (11)**	30.28 (4)**
Third wave (n=9)	26.27	26.14 (0)	26.19 (2)	26.00 (0)	25.58 (-1)	25.65 (0)	29.24 (15)*

*P<0.05, **P<0.001 compared with non-fundholders by the multiple comparison *t* test with Bonferroni correction.

50% having five or more partners per practice (table 1), partly as a result of the entry requirements (minimum list size of 7000 patients and no single handed practices).

Data on the number of deprived patients in each practice, which is based on a modification of the Jarman index,¹² suggests that first wave fundholders had the fewest deprived patients per partner. The fact that first wave fundholders are larger practices and have fewer deprived patients does not necessarily prove that they are in more affluent areas, considering the criticisms of the Jarman index.^{13 14} However, it does show that first wave fundholders are not typical practices, and this may have affected their response to the fundholding scheme. Comparing each fundholding group with the non-fundholding group for each quarter we found that the cost per 1000 patients of first wave fundholders started to differ significantly from that of the non-fundholders in the quarter before joining the fundholding scheme ($P=0.04$). Later fundholders differed significantly from the non-fundholders in the quarter they became fundholders, but not before.

It has been argued that the way in which budgets were set may have encouraged prospective fundholders to inflate their costs in order to obtain more money.¹⁵ Several studies, however, found no evidence of this.^{7 16} In Northern Ireland prospective fundholders had to carry out a prescribing study during their preparatory year to identify how they might prescribe more effectively once they became fundholders. Only very high cost practices were expected to show some improvement during their preparatory year. Any inflation of costs, therefore, would have occurred in the year before their preparatory year. Our results show no evidence of generalised wilful cost inflation, but five practices may have inflated their costs in the year before their preparatory year, one of which reduced its costs in the first year of fundholding and two in their second year of fundholding.

Possibilities for savings

Bradlow and Coulter found that none of the practices that they studied managed to reduce absolute costs on becoming fundholders.¹¹ In our study seven practices managed to reduce absolute prescribing costs in their first year of fundholding: 42 practices decreased their yearly increase on the previous year and 10 increased their yearly increase on the previous year. Nevertheless, in the third year of fundholding, first wave practices had a similar yearly percentage increase in prescribing costs to non-fundholders (table 2). Whether second and third waves will follow this pattern remains to be

seen, but the finding may suggest that the incentive for making further savings has diminished after the first two years of fundholding. In addition, in their second and third years of fundholding, the rate of increase in items per 1000 patients among first wave fundholders was the same as among non-fundholders, giving further evidence that the incentive for reducing prescribing frequency cannot be maintained.

Generic prescribing rates in Northern Ireland are lower than in England. Figures for 1995¹⁷ show that 55% of drugs were prescribed generically in England compared with 27% in Northern Ireland for the same year. However, 11% of the generically prescribed drugs in England were dispensed as proprietary drugs. This is because certain drugs are patented under a specific brand name and cannot be produced by another company until the patent expires. Drugs prescribed generically while still under patent are not recorded as generic prescribing in Northern Ireland. However, even after such drugs are excluded from the English figures, generic prescribing in Northern Ireland on average lags behind England by 17%. First wave fundholders had a generic prescribing rate of 31% in 1995, but this is still 13% lower than the rate in England. There is obviously room for improvement in generic prescribing in Northern Ireland.

Conclusions

We found that fundholding does affect prescribing in that the rate of increase in costs is reduced and the rate of generic prescribing is considerably increased. Each year's budget is reset to take account of changing factors in a practice from year to year such as, for example, pay and price increases. However, the amount left in the previous year's budget is deducted from this reset budget. This makes it increasingly difficult to make further changes each year and may also reduce the incentive to do so.

The savings that the practices can keep do not indicate what effect fundholding had on the overall regional drugs bill. Had fundholding not been introduced and costs continued to rise at the same rate as before, costs would have been higher than they actually were. So what savings did fundholders actually make? We calculated savings made in the first year of fundholding for each group by assuming that groups experienced the same percentage increase in costs in their first fundholding year as the previous year and by subtracting from this estimated cost the actual first year's costs. We estimate that the 23 first wave fundholders made savings of £1 267 748, the 34 second wave fundholders made savings of £1 476 394, and the nine third wave fundholders made savings of £451 903. This is a crude estimate and does not allow

Key messages

- The effects of fundholding on prescribing costs have been disputed
- Fundholders in Northern Ireland contained the rate of increase in prescribing costs more effectively than non-fundholders
- Fundholders increased their rate of generic prescribing by an average of 13% in the first year of fundholding
- The incentive to make further savings may have diminished after two years of fundholding

for increases in the cost of drugs (and so is probably an underestimate). However, it shows that general practitioners in Northern Ireland responded to the fundholding incentive and is clear evidence that fundholding alters the costs of prescribing.

Funding: The Drug Utilisation Research Unit is funded by the Department of Health and Social Services in Northern Ireland.

Conflict of interest: None.

- 1 Avorn J, Soumerai SB. Improving drug-therapy decisions through educational outreach: a randomized controlled trial of academically based detailing. *N Engl J Med* 1983;308:1457-63.

- 2 McGavock H, Wilson-Davis K, Rafferty T. A "compass" for general practitioner prescribers. *Health Trends* 1994;26:22-4.
- 3 Soumerai SB, McLaughlin TJ, Avorn J. Improving drug prescribing in primary care: a critical analysis of the experimental literature. *Milbank Q* 1989;67:268-317.
- 4 Bloor K, Freemantle N. Lessons from international experience in controlling pharmaceutical expenditure. 2. Influencing doctors. *BMJ* 1996;312:1525-7.
- 5 Walley T, Wilson R, Bligh J. Current prescribing in primary care in the UK: effects of indicative prescribing scheme and GP fundholding. *PharmacoEconomics* 1995;7:320-31.
- 6 Maxwell M, Heaney D, Howie JGR, Noble S. General practice fundholding: observations on prescribing patterns and costs using the defined daily dose method. *BMJ* 1993;307:1190-4.
- 7 Wilson RPH, Buchan I, Walley T. Alterations in prescribing by general practitioner fundholders: an observational study. *BMJ* 1995;311:1347-50.
- 8 Dowell JS, Snadden D, Dunbar JA. Changing to generic formulary: how one fundholding practice reduced prescribing costs. *BMJ* 1995;310:505-8.
- 9 Harris CM, Scrivener G. Fundholders' prescribing costs: the first five years. *BMJ* 1996;313:1531-4.
- 10 Steward-Brown S, Surender R, Bradlow J, Coulter A, Doll H. The effects of fundholding in general practice on prescribing habits three years after introduction of the scheme. *BMJ* 1995;311:1543-7.
- 11 Bradlow J, Coulter A. Effects of fundholding and indicative prescribing schemes on general practitioners' prescribing costs. *BMJ* 1993;307:1186-9.
- 12 Jarman B. Identification of underprivileged areas. *BMJ* 1983;286:1705-9.
- 13 Wilson-Davis K. Social deprivation and its effects on prescribing. *Prescriber* 1993;124:61-4.
- 14 Smith GD. Second thoughts on the Jarman index. *BMJ* 1991;302:359-60.
- 15 Glennerster H, Matsaganis M, Owens P. *A foothold for fundholding*. London: King's Fund Institute, 1992.
- 16 Healey AT, Reid JP. Do prospective fundholders inflate their prescribing costs? A study of Grampian fundholding and non-fundholding practices. *Health Bulletin* 1994;52:282-4.
- 17 Department of Health. *Statistical Bulletin* 1996;17:24.

(Accepted 28 May 1997)

What is to be done about fundholding?

Toby Gosden, David J Torgerson, Alan Maynard

See editorial by Keeley and p 166

National Primary Care Research and Development Centre, University of Manchester, Manchester M13 9PL

Toby Gosden, research associate

National Primary Care Research and Development Centre, Centre for Health Economics, University of York, Heslington YO1 5DD

David Torgerson, research fellow

National Primary Care Research and Development Centre, Department of Health Sciences, University of York Alan Maynard, honorary professor

Correspondence to: Mr Gosden.

BMJ 1997;315:170-1

The new Labour government has pledged to review the future of fundholding. What is the evidence base for the dissolution or reform of fundholding?

The evidence surrounding fundholding has been reviewed.¹⁻³ The authors of these reviews are in general agreement that although the relative scale of evaluating general practice fundholding is high, the results of fundholding are ambiguous. The evaluation has focused largely on the analysis of costs where there is some evidence that fundholding has been successful.⁴⁻⁹ The impact of general practice fundholding on patients' health has not been evaluated, and thus its impact on efficiency is unknown. The impact of this radical policy innovation on the quality of care, equitable provision of care, and responsiveness of care to patients' needs has been largely unresearched. Any results of evaluation, including evaluation of costs, are biased by the fact that fundholders are a self selected group differing from the generality of general practitioners. Despite this, rigorous studies with non-randomised designs can provide policy makers with useful information on the efficiency of different models of primary care.

The Audit Commission sought to determine whether fundholding was "worth" the large investment made to establish and run it.⁸ They concluded that so far the efficiency savings induced by this form of

Summary points

The evidence base on the performance of fundholding is incomplete as research on important issues such as outcomes of care is lacking; it provides no basis on which to make any hasty decisions as to the future of this policy

Careful evaluation of fundholding and all models of GP commissioning is required and is possible, ideally using trial methodology, to ensure that the scarce resources and the considerable investment that has been made in fundholding are not wasted

organisation were less than the set up and operating costs. Although the set up and administration costs of fundholding may be more than those of general practice commissioning models,¹⁰ there is no evidence about the relative benefits of these two forms of organisation. Many of the costs of fundholding are largely "sunk"—that is, it would be difficult to recoup the money spent. For example, general practitioners and practice managers have invested considerable time and effort in accumulating relevant expertise. Invest-

ment might have happened with or without fundholding, and although funds cannot be recouped they may produce considerable returns in future. Similar investments have created useful new skills in health authorities. This investment and other factors are leading to the creation of larger funds where skills are pooled, but their efficiency has yet to be evaluated.

If all that these investments have produced are once-only savings in areas such as prescribing, the case for retaining fundholding is weak. However, if cost containment and increased efficiency can occur only in the longer term, the returns to investment in management are yet to appear; thus it is not surprising that existing studies give us little confidence in the policy. Further time, experience, and evaluation are needed to evaluate this experiment properly.

The impact of fundholding on equity is thought to be deleterious, creating "two tierism."¹¹ However, a distinction has to be made between principle and practice. The principle is that the NHS treats patients in equal need (capacity to benefit per unit of cost) with equal treatment. In practice there are considerable variations in the provision of care. Whether, after fundholding, this would be more inequitable than before fundholding has to be evaluated fully.

Evaluation is possible

The remarkable aspect both of the internal market reforms and the 1996 white papers on primary care^{12 13} is the failure of a government that espoused "value for money" to establish performance benchmarks and to monitor the performance of policies such as general practice fundholding in relation to these benchmarks. The case for the maintenance or euthanasia of fundholding does not exist because the government is reluctant to evaluate the reforms. Rigorous evaluation of such reforms using randomised trial methodology is possible (as shown by the Rand experiment and the

American negative income tax experiments¹⁴), but it is rarely undertaken and usually ignored by government research and development programmes such as that recently established in Britain.

An incoming Labour government may repeat this error and waste resources on reforms such as locality purchasing which are not evidence based and not established to perform well in relation to clear policy goals. Labour, like its Conservative predecessors, will pursue efficiency (measured in terms of costs and benefits—that is, improvements in patients' health and equity). It must articulate these targets carefully and explicitly and rely on evidence rather than rhetoric to sustain its reforms. A failure to do this will waste scarce economic resources and prejudice patients' care.

- 1 Petty R. General practitioner fundholding: weighing the evidence. *Lancet* 1995;346:1139-42.
- 2 Dixon J, Glennerster H. What do we know about fundholding in general practice? *BMJ* 1995;311:727-30.
- 3 Gosden T, Torgerson DJ. The effect of fundholding on prescribing and referral costs: a review of the evidence. *Health Policy* 1997;40:103-14.
- 4 Stewart-Brown S, Surender R, Bradlow J, Coulter A, Doll H. The effects of fundholding in general practice on prescribing habits three years after introduction of the scheme. *BMJ* 1995;311:1543-7.
- 5 Burr A, Walker R, Stent S. Impact of fundholding on general practice prescribing patterns. *Pharm J* 1992;249(suppl):R8.
- 6 Whynes D, Baines D, Tolley K. GP fundholding and the costs of prescribing. *J Public Health Med* 1995;17:323-9.
- 7 Wilson RPH, Buchan I, Walley T. Alterations in prescribing by general practitioner fundholders: an observational study. *BMJ* 1995;311:1347-50.
- 8 Audit Commission. *A prescription for improvement: towards more rational prescribing in general practice*. London: HMSO, 1994.
- 9 Harris CM, Scrivener G. Fundholders' prescribing costs: the first five years. *BMJ* 1996;313:1531-3.
- 10 Pennington S. GPs enter first total commissioning scheme. *GP* 1995 April 21:21.
- 11 Association of Community Health Councils. Fundholding and access to hospital care. *Health News Briefing* 1994 March.
- 12 Secretaries of State for Health, Scotland, and Wales. *Choice and opportunity. Primary care: the future*. London: Stationery Office, 1996. (Cm3390.)
- 13 Secretaries of State for Health, Scotland, and Wales. *Primary care: delivering the future*. London: Stationery Office, 1996. (Cm 3512.)
- 14 Ferber I, Hirsch W. *Social experimentation and economic policy*. Cambridge: Cambridge University Press, 1982.

(Accepted 19 May 1997)

Any questions

The benefits of influenza immunisation

The head of an organisation is advised by a member of a profession supplementary to medicine that the workforce should be offered immunisation against influenza next year. Is this the right advice?

Probably not. The advice is presumably on the grounds either that immunising the workforce will reduce the level of sickness absence in the organisation during the winter months (so saving money for the organisation) or that it will be a staff benefit and prevent individual ill health. The evidence is conflicting, and less convincing from Britain than the United States, but on balance it seems that the rate of influenza among young, healthy people is significantly reduced in vaccinated versus unvaccinated groups.¹ The effect is mainly in years when there is a larger epidemic but these are not possible to predict in advance. Outside military or trial situations, however, only a proportion of the workforce (often a small proportion) will accept the vaccine. The low vaccine uptake, together with the incomplete efficacy of the vaccine, the relatively low attack rate of influenza even in epidemics (around 20%), and the high prevalence of other respiratory infections at the same time of year, mean that the reduction in sickness absence achieved by a routine immunisation programme is low in most years and will not offset the cost to the organisation of mounting the programme and purchasing the vaccine for its staff. If the vaccine programme is to be a perk then it should at least

focus on staff with respiratory, cardiac, or renal disease, diabetes mellitus, or immunosuppression. If the employer is a healthcare organisation there are special considerations. In the United States routine influenza vaccination is recommended for this group because of their contact with vulnerable patients.² In Britain, however, routine vaccination is recommended only for people with chronic disease, especially if elderly, and those living in long stay facilities, and not for healthcare or other key workers.³ Healthcare workers can be offered prophylaxis with amantadine during an epidemic.³ An employer is at liberty to offer staff annual influenza vaccination, even though this is not officially recommended in Britain or the United States, but this will not save money. If a benefit to the health of individual staff is the aim then the money might be better spent on providing healthy canteen food and exercise facilities and discouraging smoking.

Anne Cockcroft, consultant in occupational medicine, London

- 1 McNair A. *A reassessment of flu vaccine efficacy: the case for comprehensive immunisation programmes*. Influenza Monitoring and Advisory Bureau: Vox Prism, 1991.
- 2 Centers for Disease Control and Prevention. *Prevention and control of influenza. Recommendations of the Advisory Committee on Immunization Practices*. Atlanta: Centers for Disease Control and Prevention, 1993.
- 3 UK health departments. *Immunisation against infectious disease*. London: HMSO, 1996.