Sporadic Creutzfeldt-Jakob disease in the United Kingdom: analysis of epidemiological surveillance data for 1970-96

S N Cousens, M Zeidler, T F Esmonde, R De Silva, J W Wilesmith, P G Smith, R G Will

Abstract

Objective: To identify changes in the occurrence of Creutzfeldt-Jakob disease that might be related to the epidemic of bovine spongiform encephalopathy.

Design: Epidemiological surveillance of the United Kingdom population for Creutzfeldt-Jakob disease based on (a) referral of suspected cases by neurologists, neuropathologists, and neurophysiologists and (b) death certificates.

Setting: England and Wales during 1970-84, and whole of the United Kingdom during 1985-96.

Subjects: All 662 patients identified as sporadic cases of Creutzfeldt-Jakob disease.

Main outcome measures: Age distribution of patients, age specific time trends of disease, occupational exposure to cattle, potential exposure to causative agent of bovine spongiform encephalopathy.

Results: During 1970-96 there was an increase in the number of sporadic cases of Creutzfeldt-Jakob disease recorded yearly in England and Wales. The greatest increase was among people aged over 70. There was a statistically significant excess of cases among dairy farm workers and their spouses and among people at increased risk of contact with live cattle infected with bovine spongiform encephalopathy. During 1994-6 there were six deaths from sporadic Creutzfeldt-Jakob disease in the United Kingdom in patients aged under 30.

Conclusions: The increase in the incidence of sporadic Creutzfeldt-Jakob disease and the high incidence in dairy farmers in the United Kingdom may be unrelated to bovine spongiform encephalopathy. The most striking change in the pattern of Creutzfeldt-Jakob disease in the United Kingdom after the epidemic of bovine spongiform encephalopathy is provided by the incidence in a group of exceptionally young patients with a consistent and unusual neuropathological profile. The outcome of mouse transmission studies and the future incidence of the disease in the United Kingdom and elsewhere, will be important in judging whether the agent causing bovine spongiform encephalopathy has infected humans.

Introduction

The United Kingdom National Creutzfeldt-Jakob Disease Surveillance Unit was set up in May 1990 after

recognition of the epidemic of bovine spongiform encephalopathy in the late 1980s.¹ If bovine spongiform encephalopathy infects humans it was considered most likely to result in disease manifestations similar to those of Creutzfeldt-Jakob disease and that some or all of the following phenomena might be observed: (a) an increase in the overall incidence of Creutzfeldt-Jakob disease in the United Kingdom; (b) an excess of cases in groups most likely to have high exposure to the causative agent of bovine spongiform encephalopathy; (c) a change in the epidemiological pattern of Creutzfeldt-Jakob disease, such as a change in the age distribution; (d) a change in the clinical or neuropathological characteristics of Creutzfeldt-Jakob disease. The identification in 1996 of a group of comparatively young patients with Creutzfeldt-Jakob disease characterised by unusual clinical and neuropathological features² led the Spongiform Encephalopathy Advisory Committee to advise the United Kingdom government that "the most likely explanation at present is that these cases are linked to exposure to BSE."³

We present data on age specific changes in the incidence of Creutzfeldt-Jakob disease in the United Kingdom since 1970 and examine the incidence rates of the disease since 1990 in groups who might have been occupationally exposed to the causative agent of bovine spongiform encephalopathy.

Methods

Surveillance of Creutzfeldt-Jakob disease

A series of studies has attempted to identify all cases of Creutzfeldt-Jakob disease occurring in England and Wales since 1970 and in the United Kingdom since 1985. Methods of case ascertainment have been described.²⁻⁴ Briefly, cases of Creutzfeldt-Jakob disease occurring in England and Wales during 1970-9 were identified retrospectively at the end of that period⁵ and prospective surveillance instituted to detect cases during 1980-4.⁶ Prospective ascertainment of cases, extended to cover the whole of the United Kingdom, was reinstituted in May 1990 and is continuing. Cases occurring in the United Kingdom between 1985 and April 1990 were identified retrospectively.

Case ascertainment has been based throughout on direct notification of cases and on death certificates. Neurologists, neuropathologists, and neurophysiologists are asked to refer suspected cases of Creutzfeldt-Jakob disease, and the Office for National Statistics

Department of Infectious and Tropical Diseases, London School of Hygiene and Tropical Medicine, London

WC1E 7HT

S N Cousens,
senior lecturer

P G Smith,
professor of tropical
epidemiology and
department head

National
Creutzfeldt-Jakob
Disease

Surveillance Unit,
Western General
Hospital, Edinburgh
EH4 2XU

M Zeidler,
research registrar

T F Esmonde,
research registrar

R De Silva,
research registrar

R G Will,
consultant neurologist

Central Veterinary
Laboratory,
Addlestone, Surrey
KT15 3NB

J W Wilesmith,
head of epidemiology
department

Correspondence to:
Mr Cousens
(s.cousens@lshtm.
ac.uk).

BMJ 1997;315:389-96

supplies death certificates coded under rubrics 339.9 and 781.7 (*International Classification of Diseases*, eighth revision (ICD-8)) and 046.1 and 331.9 (ICD-9). A few cases are referred by other routes.

Since May 1990 suspected cases have been seen, when possible, by a neurologist from the national surveillance unit and a close relative has been interviewed to collect epidemiological information, including the current and previous occupations of the patient.

Case definition

The diagnostic criteria for Creutzfeldt-Jakob disease have been essentially the same since surveillance began in 1970.⁷ In summary, classification as a definite case requires neuropathological confirmation of spongiform change, or (since 1993) immunocytochemical confirmation of the presence of prion protein, or identification of scrapie associated fibrils on electron microscopy. Currently about 70% of patients with suspected Creutzfeldt-Jakob disease come to necropsy. In the absence of neuropathological investigation patients are classified as probable cases of Creutzfeldt-Jakob disease if they present with progressive dementia, a typical electroencephalogram, and at least two of the following: myoclonus; visual or cerebellar signs; pyramidal or extrapyramidal signs; akinetic mutism.

Cases are classified as iatrogenic when the patients have a known risk factor for accidental transmission (for example, pituitary hormone recipients). The criteria for inherited Creutzfeldt-Jakob disease have been updated to include information available from genetic analyses. Cases are now classified as inherited if the patients carry a disease specific mutation of the prion protein gene or have a history of probable or definite Creutzfeldt-Jakob disease in a first degree relative. Molecular biological data were available for 53% of cases identified since May 1990. All other cases are classified as sporadic. Probably during 1970-90 roughly half of inherited cases of Creutzfeldt-Jakob disease—that is, one or two a year—may have been classified as sporadic because genetic analyses were not carried out during the period.

Groups with potentially high occupational exposure

We do not know how or indeed if bovine spongiform encephalopathy is transmitted to humans. We identified three routes through which transmission might be hypothesised to occur.

Route 1 might be via contact with ruminant derived meat and bone meal contaminated by bovine spongiform encephalopathy (the likely source of the bovine epidemic).⁸ Occupational groups identified to be at high risk of this exposure were workers on farms with cattle, pigs, poultry, sheep, or goats (as ruminant derived meat and bone meal may have been fed to these animals) and workers in feed mills producing meat and bone meal.

Route 2 might be via contact with live cattle infected with bovine spongiform encephalopathy. Occupational groups identified to be at high risk of this exposure were workers on farms with cattle, veterinarians, and abattoir workers. Three different populations of farm workers with increasing likelihood of exposure to live infected cattle were identified—namely, workers on

all farms with cattle (dairy or beef herds); workers on farms with dairy cattle (the incidence of bovine spongiform encephalopathy has been much higher in dairy herds)⁹; workers on dairy or beef farms with a confirmed case of bovine spongiform encephalopathy.

Route 3 might be via contact with brain or spinal cord from dead infected cattle. Occupational groups identified to be at high risk of this exposure were veterinarians, abattoir workers, workers in rendering plants, and butchers. Though cattle may be slaughtered on farms by licensed personnel and dressed by farmers, we have no evidence that this is widespread and farmers were therefore excluded from this occupational group.

The numbers of workers on different types of farms during 1990-4 were obtained from the annual agricultural censuses for England and Wales, Northern Ireland, and Scotland (1990-5). Data on the numbers of workers on farms affected by bovine spongiform encephalopathy were obtained by linking agricultural census data with the database on the disease held at the Central Veterinary Laboratory. The number of veterinarians, butchers, and meat cutters in Great Britain was estimated from the 10% sample of the 1991 census by using the recommended conversion factor of 10.16. The number of abattoir workers could not be derived from published census data for Great Britain,

Fig 1 Yearly numbers of deaths from Creutzfeldt-Jakob disease in England and Wales during 1970-96 and in Scotland and Northern Ireland during 1985-96 (data exclude known iatrogenic and inherited cases). Horizontal lines represent numbers of deaths corresponding to yearly death rates of 0.5/million (lower) and 1.0/million (upper)

Table 1 Numbers of deaths from Creutzfeldt-Jakob disease in 10 year age groups for each two year period in England and Wales (from 1970) and United Kingdom* (from 1985)

Age group (years)	Two year period														Total†
	1970-1	1972-3	1974-5	1976-7	1978-9	1980-1	1982-3	1984-5	1986-7	1988-9	1990-1	1992-3	1994-5	1996†	
10-19	0	0	0	0	0	1	0	0	0	0	0	0	1	1 (1)	3 (1)
20-29	0	0	0	0	0	0	0	0	0	0	0	0	1	3 (0)	4 (0)
30-39	1	0	0	2	2	1	1	4	1	0	1	0	1	4 (0)	18 (0)
40-49	2	0	2	1	1	2	1	0	3	2	2	3	4	4 (0)	27 (0)
50-59	7	9	11	6	11	13	12	10	5	13	18	12	14	4 (1)	145 (1)
60-69	9	13	10	22	17	24	20	28	22	17	28	38	32	8 (2)	288 (2)
≥70	2	2	2	4	9	4	12	16	18	17	9	29	37	12 (0)	173 (0)
Total	21	24	25	35	40	45	46	58	49	49	58	82	90	36 (4)	658 (4)

Table excludes cases known to be iatrogenic or inherited.

*United Kingdom includes cases from Scotland and Northern Ireland.

†Figures in parentheses are numbers of patients alive on 1 January 1997.

Table 2 Rate of increase in mortality from sporadic Creutzfeldt-Jakob disease by age group

Age group (years)	Average % increase in mortality over two year period (95% CI)	P value
40-49	6.6 (-4.8 to 19.5)	0.27
50-59	4.1 (-0.1 to 8.8)	0.08
60-69	7.3 (3.9 to 10.8)	<0.001
≥70	21.3 (15.4 to 27.5)	<0.001

but information on these workers was provided by the Ministry of Agriculture, Fisheries and Food.

Data for Northern Ireland on veterinarians and on workers employed in the slaughtering of animals and the production of meat for 1993 were provided by the Department of Economic Development, Belfast. We could not derive the number of workers in rendering plants or feed mills producing meat and bone meal from published census data for Great Britain. However, the numbers of workers in these groups are small compared with the number of farm workers (see below) and their exclusion from calculation of the expected number of cases is therefore of no consequence. (The number of feed mill workers is estimated at about 3000 (United Kingdom Agricultural Supply Trade Association, personal communication, 1996).)

Expected numbers of cases in different occupational groups

Expected numbers of deaths from Creutzfeldt-Jakob disease between 1 May 1990 and 31 December 1996 were calculated by multiplying age specific and sex specific death rates for the United Kingdom population by the estimated numbers of people in the different occupational groups. The age and sex distribution of farm workers was assumed to follow that reported for farm workers in England and Wales in the European Commission's structure survey of 1990. The age and sex distributions of veterinarians and of butchers and meat cutters were assumed to follow those of the wider occupational groups within which they were classified in the 1991 census.

Observed and expected numbers of cases were compared in the assumption that the observed number of cases followed a Poisson distribution.

Results

Between 1 January 1970 and 31 December 1996, 708 definite and probable cases of Creutzfeldt-Jakob

disease were identified. Five were in patients alive on 31 December 1996. In all, 540 (76%) cases were classified as definite. Twenty three cases were classified as iatrogenic and 23 as inherited, both of which groups were excluded from analysis. Among the 662 sporadic cases, 15 were of the variant form of the disease.

Distribution of cases by age, sex, and time period

In England and Wales the yearly number of deaths from Creutzfeldt-Jakob disease increased from around 10 at the beginning of the 1970s to around 40 in the 1990s (fig 1). No increase in yearly numbers of deaths was evident for Scotland and Northern Ireland after 1985 (fig 1).

Figure 2 shows the average yearly age specific and sex specific death rates over the study period. Below 40 years of age, death rates were extremely low (<0.1/million yearly). Death rates increased substantially in the 50-59 year age group and reached a peak of around 2.0/million yearly in the 60-69 year age group before declining in people aged 70 and over.

Table 1 shows the numbers of deaths in 10 year age groups for each two year period from 1970 to 1996. Mortality from Creutzfeldt-Jakob disease increased substantially over the period in patients aged 70 and over and by successively smaller proportions in those aged 60-69 and 50-59 (fig 3; table 2). The increase in patients aged 40-49 years was not significant ($P = 0.27$). Under the age of 40 numbers of cases were small, but of the seven patients who died under the age of 30, six

Fig 2 Age and sex specific average yearly death rates from Creutzfeldt-Jakob disease in United Kingdom during 1970-96

Fig 3 Trends in age specific death rates from Creutzfeldt-Jakob disease in England and Wales during 1970-96. For age group 40-49 years death rates for 1972-3 and 1984-5 are not plotted as there were no deaths in this age group in the periods

had onset of the disease after 1 January 1994. All six died in 1995 or 1996.

Sporadic disease in groups with potential occupational exposure

Six cases of sporadic Creutzfeldt-Jakob disease (five definite, four in men) were identified from 1 May 1990 in people whose occupation at disease onset was in one of the groups identified above. Four have been the subject of case reports.¹⁰⁻¹³ Ages at death in the six cases were 54, 54, 59, 61, 64, and 64 years. All five patients subjected to neuropathological investigation showed

changes typical of Creutzfeldt-Jakob disease. None showed the "florid" plaques characteristic of the variant form of the disease.² The sixth, probable case followed a clinical course typical of sporadic Creutzfeldt-Jakob disease with a characteristic electroencephalogram.

Four of the six patients lived or worked on dairy farms (on three of which there had been confirmed cases of bovine spongiform encephalopathy), and two lived or worked on farms with beef suckler herds (one with a confirmed case of bovine spongiform encephalopathy). Two of the six cases were in spouses of farmers. All patients had lived or worked on farms throughout their working lives. Table 3 shows the distribution of these cases by occupational group. There was a significant excess of cases among animal farm workers (6 observed, 2.4 expected; $P=0.03$). All these cases were in cattle farmers, four of whom worked on farms with a case of bovine spongiform encephalopathy. This excess of cases was highly significant (4 observed, 0.58 expected; $P=0.003$). Table 4 shows the distribution of observed and expected cases by hypothesised route of transmission. There was an excess of cases among workers potentially exposed to ruminant derived meat and bone meal and among workers exposed to live cattle infected with bovine spongiform encephalopathy. Two cases among farmers were on farms without a case of bovine spongiform encephalopathy compared with an expected number of 1.78 ($P=0.5$).

Discussion

In our introduction we identified four changes in the pattern of Creutzfeldt-Jakob disease that might occur if exposure to the bovine agent caused spongiform encephalopathy in humans. Arguably all of these

Table 3 Distribution of expected and observed cases of Creutzfeldt-Jakob disease among occupational groups potentially at increased risk of infection with bovine spongiform encephalopathy

Occupational group	No of subjects in group*	No of cases of Creutzfeldt-Jakob disease 1 May 1990 to 31 December 1996		Yearly incidence/million	P value†
		Expected	Observed		
Farm workers on farms with cattle, pigs, poultry, sheep, or goats	447 162	2.36	6	2.01	0.03
Farm workers on farms with cattle (dairy or beef)	354 242	1.87	6	2.54	0.01
Farm workers on farms with dairy cattle	149 436	0.79	4	4.02	0.01
Farm workers on farms with confirmed case of bovine spongiform encephalopathy	109 643	0.58	4	5.47	0.003
Veterinarians	8 225	0.03	0	0	1.0
Abattoir workers, butchers, meat cutters	68 300‡	0.15	0	0	1.0

*Numbers given for farm workers include spouses and represent an average over period.

†P values indicate probability of obtaining observed number of cases or more under null hypothesis that there is no occupational risk of Creutzfeldt-Jakob disease.

‡Includes an estimated 48 300 butchers and meat cutters in Great Britain (1991 census), 12 400 staff employed in red meat slaughtering (Ministry of Agriculture, Fisheries and Food), and 7600 staff employed in slaughtering animals and production of meat in Northern Ireland (Department of Economic Development, Belfast).

Table 4 Distribution of expected and observed cases of Creutzfeldt-Jakob disease by hypothesised route of exposure to agent of bovine spongiform encephalopathy

Type of potential contact with agent of bovine spongiform encephalopathy	No of subjects exposed*	No of cases of bovine spongiform encephalopathy 1 May 1990 to 31 December 1996		Yearly incidence/million	P value†
		Expected	Observed		
Contact with ruminant derived meat and bone meal (farm workers on farms with cattle, pigs, poultry, sheep, or goats)	447 162	2.36	6	2.01	0.03
Contact with live cattle infected with bovine spongiform encephalopathy (farm workers on farms with confirmed case of bovine disease, veterinarians, abattoir workers)	132 408	0.64	4	4.53	0.004
Contact with brain and spinal cord of dead cattle (butchers and meat cutters)	68 300	0.15	0	0	1.0

*Numbers given for farm workers include spouses and represent an average over period.

†P values indicate probability of obtaining observed number of cases or more under null hypothesis that there is no occupational risk of Creutzfeldt-Jakob disease.

Table 5 Incidence of Creutzfeldt-Jakob disease in countries other than United Kingdom over time

Country	Yearly incidence/million
Chile:	
1955-72	0.10
1973-7	0.31
1978-83	0.69
France:	
1968-77	0.34
1978-82	0.58
1992-4	0.81
Germany:	
1979-90	0.31
1993-4	0.68
United States:	
1973-7	0.26
1986-8	0.83

Table 6 Incidence rates of Creutzfeldt-Jakob disease in dairy farmers in different European countries

Country	Period	No of cases	Yearly incidence/million
France	1992-4	4	4.8
Germany	1993-4	2	3.8
Italy	1993	2	9.4
Netherlands	1993-4	0	0
United Kingdom	1990-6	4	4.0

changes may have occurred. There are, however, important caveats that inhibit clear interpretation of our findings.

The number of sporadic cases of Creutzfeldt-Jakob disease recorded in England and Wales has increased. During a period of prospective surveillance before the advent of bovine spongiform encephalopathy (1980-4) the yearly number of deaths from sporadic Creutzfeldt-Jakob disease in England and Wales averaged 24.8. During prospective surveillance after the epidemic of bovine spongiform encephalopathy (1990-6) the yearly number of deaths averaged 33.6. However, similar increases in the incidence of Creutzfeldt-Jakob disease have been observed in countries where the bovine disease is rare or absent (table 5).^{14 15} These increases may reflect improved case ascertainment rather than real increases in incidence. It has long been suspected that the age distribution of Creutzfeldt-Jakob disease in the United Kingdom (fig 2) and elsewhere—with a decline in incidence in people over 70—might reflect under ascertainment of cases in older age groups.¹⁶ In the United Kingdom the greatest increase in incidence has been in people aged over 70. For many diseases case ascertainment is likely to be poorest among old people and best among age groups in which death from any cause is unusual. Thus aside from the recent increase in cases in young people, the age specific time trends suggest that improved ascertainment may explain much or all of the apparent rise in the incidence of Creutzfeldt-Jakob disease since 1970.

We found significant excesses of cases of Creutzfeldt-Jakob disease among workers on dairy farms and on farms with a confirmed case of bovine spongiform encephalopathy. None of the six patients who might have been occupationally exposed to the agent of bovine spongiform encephalopathy presented with the variant form of Creutzfeldt-Jakob

disease, which might indicate a different aetiology from the variant form. However, evidence from iatrogenic cases that route of infection may be a determinant of the clinical and pathological pattern of disease¹⁷ means that we cannot exclude entirely the possibility that “occupational” cases and the variant form of the disease arise from exposure to the same agent by different routes. One possible explanation for the apparent excess risk among farmers is that case ascertainment of Creutzfeldt-Jakob disease in the United Kingdom has been better in this group than among other groups because of awareness of a possible link between the bovine and human diseases. Moreover, though the incidence of Creutzfeldt-Jakob disease among dairy farmers in the United Kingdom is higher than in the general population, it does not seem remarkable when compared with the incidence among dairy farmers in other European countries where bovine spongiform encephalopathy is rare or absent (table 6).¹⁵ This suggests that dairy farmers may be at increased risk of Creutzfeldt-Jakob disease for reasons other than exposure to the causative agent of the bovine disease. We emphasise also that, though some farm workers seem to be at increased risk of Creutzfeldt-Jakob disease compared with the general population, their absolute risk remains extremely low, the yearly incidence of the disease being well below 10 cases/million.

Our analyses are based on occupation at the time of disease onset, as national data are not available on the numbers of people ever employed in particular occupational groups. We therefore excluded from the analyses one patient who had worked in an abattoir for 10 months, six years before disease onset in 1995; one patient who had worked on a farm with dairy and beef herds until retiring in 1991, four years before disease onset in 1995; and one patient with the variant form of the disease who had worked as a butcher between 1985 and 1987. Another case of interest that was similarly excluded was of a man who transported meat and bone meal from an animal feed compounder to farms for many years up to 1991 but who changed his job before onset of Creutzfeldt-Jakob disease in 1993.

Cases with unusual features

During 1994-6, 10 cases of sporadic Creutzfeldt-Jakob disease occurred in the United Kingdom in patients under the age of 30. Based on past experience in the United Kingdom (two cases in the preceding 24 years) and elsewhere² this is unusual. That nine of these patients together with five others aged 30, 32, 34, 39, and 48 years at onset also exhibited a previously undescribed neuropathological pattern renders this finding even more remarkable and raises important questions. Were these cases really “new” or had similar cases occurred previously without being identified? If these cases were new why did they occur in the United Kingdom in the mid-1990s?

Bruton et al reviewed the Runwell Hospital Brain Archive and identified 19 cases of spongiform encephalopathy, only 11 of which had been diagnosed before death.¹⁸ All the patients died before 1985, and none showed the neuropathological changes characteristic of the variant form of Creutzfeldt-Jakob disease. The authors suggested that prion diseases may be more common than supposed. Only about 4% of elderly

patients dying with dementia come to necropsy.¹⁹ Taken with our finding of a substantial increase over time in the number of cases identified in older age groups, it seems likely that some cases of Creutzfeldt-Jakob disease in this age group have been missed. To maintain the hypothesis that the phenomenon of cases in young people with striking neuropathological changes is not new requires that young people dying of progressive neurological disorders over the past several decades came to necropsy sufficiently rarely for such cases to have been missed entirely. Since 1979 about 46% of patients aged under 45 dying of dementias with which Creutzfeldt-Jakob disease might be confused have come to necropsy, with no evidence that necropsy rates have increased over that time.¹⁹ Thus though we cannot exclude the possibility that such cases were missed, it seems to us an unlikely explanation.

If we accept for the present that these young cases are a new phenomenon, why did they appear in the United Kingdom in the mid-1990s? Bovine spongiform encephalopathy has been advanced as the most likely explanation.³ Other workers have asked why in the absence of direct evidence of a link bovine spongiform encephalopathy should be singled out ahead of microwave ovens, high voltage power lines, or organophosphorous sheep dips.²⁰ These exposures are not confined to the United Kingdom and therefore seem less plausible explanations than bovine spongiform encephalopathy. The one case of the variant form of Creutzfeldt-Jakob disease reported in France²¹ does not compromise the plausibility of the link with bovine spongiform encephalopathy. France has been one of the biggest markets for exported British beef, beef products, and cattle (B Schreuder, J Wilesmith, O Straub, personal communication, 1996).²²

One argument against the plausibility of a link between bovine spongiform encephalopathy and the variant form of Creutzfeldt-Jakob disease is that the variant disease has been observed mainly in young people. The lack of cases in older age groups could be due to misdiagnosis, reduced susceptibility, or age related exposure to the agent.² That the variant form of the disease may have been missed in old people seems plausible but that it would be missed in patients in their late 40s and early 50s seems less so. That age related exposure explains the observed age distribution of the cases is also difficult to sustain, as the link between age and exposure would have to be very strong, and we have identified no exposure related to bovine spongiform encephalopathy that meets this condition.

In summary, the epidemiological data available suggest that the distribution of Creutzfeldt-Jakob disease in the United Kingdom population may be changing. Neuropathological data suggest that a new form of the disease may have appeared which could partly but not wholly explain the epidemiological anomalies observed. There is no direct evidence that links bovine spongiform encephalopathy with these changes, but recent observations accord with this hypothesis.²³⁻²⁴ The results of transmission studies from human Creutzfeldt-Jakob disease cases to mice and the future incidence of Creutzfeldt-Jakob disease in the United Kingdom and elsewhere will be important in judging whether the causative agent of bovine spongiform encephalopathy has infected humans.

Key messages

- The overall incidence of Creutzfeldt-Jakob disease in the United Kingdom has increased in recent years
- The largest increases have been in the oldest age groups and probably represent improved case ascertainment rather than a real change in incidence
- The comparatively high incidence of Creutzfeldt-Jakob disease in dairy farmers in the United Kingdom is comparable to that observed in dairy farmers in countries where bovine spongiform encephalopathy is rare or absent
- The observation of a group of comparatively young patients with Creutzfeldt-Jakob disease characterised by unusual neuropathological features during 1994-6 remains unexplained
- The results of mouse transmission studies and the future incidence of Creutzfeldt-Jakob disease in the United Kingdom and elsewhere will be important in judging whether the causative agent of bovine spongiform encephalopathy has infected humans

Hill et al have recently reported the findings of a Western blot analysis of protease resistant prion protein (PrP^{RES}) from five of the six farmers reported here.²⁵ On the basis of fragment size and glycoform ratio of this protein, all five were classified as having type I or type II PrP^{RES}. These are types associated with the classical form of Creutzfeldt-Jakob disease and not with the variant form. These authors conclude that their results, in conjunction with the phenotype of these patients, do not support a link with occupational exposure to the agent of bovine spongiform encephalopathy.

We thank J Mackenzie for data management; W B Matthews for his foresight in initiating surveillance of Creutzfeldt-Jakob disease in the United Kingdom in 1980 and for subsequent advice; R Knight and R Harries-Jones for invaluable contributions to surveillance of the disease in England and Wales during 1970-84; and James Ironside, Jeanne Bell, Linda McCardle, and Caroline Barrie for neuropathological data. We also thank all neurologists, neuropathologists, and neurophysiologists in Britain, without whose help Creutzfeldt-Jakob disease surveillance would not be possible.

Funding: The Creutzfeldt-Jakob disease surveillance unit is funded by the Department of Health and the Scottish Home and Health Department and supported by the Biotechnology and Biological Sciences Research Council (grant No 15/BS204814).

Conflict of interest: None.

- 1 Wells GAH, Scott AC, Johnson CT, Gunning RF, Hancock RD, Jeffrey M, et al. A novel progressive spongiform encephalopathy in cattle. *Vet Rec* 1987;121:419-20.
- 2 Will RG, Ironside JW, Zeidler M, Cousens SN, Estibeiro K, Alperovitch A, et al. A new variant of Creutzfeldt-Jakob disease in the UK. *Lancet* 1996;347:921-5.
- 3 Spongiform Encephalopathy Advisory Committee. *Statement*. London, HMSO, 1996.
- 4 Cousens SN, Harries-Jones R, Knight R, Will R, Smith PG, Matthews WB. Geographical distribution of cases of Creutzfeldt-Jakob disease in England and Wales 1970-84. *J Neurol Neurosurg Psychiatry* 1990;53:459-65.
- 5 Will RG, Matthews WB. A retrospective study of Creutzfeldt-Jakob disease in England and Wales 1970-1979. I: clinical features. *J Neurol Neurosurg Psychiatry* 1984;47:134-40.

- 6 Harries-Jones R, Knight R, Will RG, Cousens S, Smith PG, Matthews WB. Creutzfeldt-Jakob disease in England and Wales, 1980-84: a case-control study of potential risk factors. *J Neurol Neurosurg Psychiatry* 1988; 51:1113-9.
- 7 Will RG. Surveillance of prion diseases in humans. In: Baker H, Ridley RM, eds. *Methods in molecular medicine: prion diseases*. Totowa: Humana Press, 1996:119-37.
- 8 Wilesmith JW, Wells GAH, Cranwell MP, Ryan JMB. Bovine spongiform encephalopathy: epidemiological studies. *Vet Rec* 1988;123:638-44.
- 9 Wilesmith JW. Bovine spongiform encephalopathy: epidemiological factors associated with the emergence of an important new animal pathogen in Great Britain. *Semin Virol* 1994;5:179-87.
- 10 Sawcer SJ, Yuill GM, Esmonde TFG, Estibeiro P, Ironside JW, Bell JE, et al. Creutzfeldt-Jakob disease in an individual occupationally exposed to BSE. *Lancet* 1993;341:642.
- 11 Davies PTG, Jahtar S, Ferguson IT, Windl O. Creutzfeldt-Jakob disease in an individual occupationally exposed to BSE. *Lancet* 1993;342:680.
- 12 Smith PEM, Zeidler M, Ironside JW, Estibeiro P, Moss TH. Creutzfeldt-Jakob disease in a dairy farmer. *Lancet* 1995;346:898.
- 13 Young GR, Fletcher NA, Zeidler M, Estibeiro K, Ironside JW. Creutzfeldt-Jakob disease in a beef farmer. *Lancet* 1996;348:610-1.
- 14 Brown P, Cathala F, Raubertas RF, Gajdusek DC, Castaigne P. The epidemiology of Creutzfeldt-Jakob disease: conclusions of a 15-year investigation in France and review of the world literature. *Neurology* 1987;37:895-904.
- 15 National CJD Surveillance Unit. *Fourth annual report to the Department of Health*. Edinburgh: NCJDSU, 1995.
- 16 Will RG, Matthews WB, Smith PG, Hudson C. A retrospective study of Creutzfeldt-Jakob disease in England and Wales. II: Epidemiology. *J Neurol Neurosurg Psychiatry* 1986;49:749-55.
- 17 Bell JE, Ironside JW. Neuropathology of spongiform encephalopathies in humans. *Br Med Bull* 1993;49:738-77.
- 18 Bruton CJ, Bruton RK, Gentleman SM, Roberts GW. Diagnosis and incidence of prion (Creutzfeldt-Jakob) disease: a retrospective archival survey with implications for further research. *Neurodegeneration* 1995;4:357-68.
- 19 Aylin P, Rooney C, Drever F, Coleman M. Increasing mortality from Creutzfeldt-Jakob disease in England and Wales since 1979: ascertainment bias from increase in post-mortems? *Popul Trends* 1996;85:34-8.
- 20 Wickham A. CJD and BSE: advisory committee's conclusion was based on "absence of any credible alternative." *BMJ* 1996;312:1038.
- 21 Chazot G, Broussolle E, Lapras CI, Blättler T, Aguzzi A, Kopp N. New variant of Creutzfeldt-Jakob disease in a 26-year-old French man. *Lancet* 1996;347:1181.
- 22 Meyer LA. *Impact of BSE on UK bone-in beef exports and UK beef fatstock prices*. Wageningen: Wageningen Agricultural University, 1996. (MSc thesis.)
- 23 Lasmézas CI, Deslys J-P, Demaimay R, Adjou KT, Lamoury F, Dormont D. BSE transmission to macaques. *Nature* 1996;381:743-4.
- 24 Collinge J, Sidle KCL, Meads J, Ironside J, Hill AF. Molecular analysis of prion strain variation and the aetiology of "new variant" CJD. *Nature* 1996;383:685-90.
- 25 Hill AF, Will RG, Ironside J, Collinge J. Type of prion protein in UK farmers with Creutzfeldt-Jakob disease. *Lancet* 1997;350:188.

(Accepted 19 March 1997)

Commentary: Age related exposure of patients to the agent of BSE should not be downplayed

Sheila M Gore

There have been two periods of prospective surveillance of Creutzfeldt-Jakob disease in the United Kingdom—during 1980-4, covering England and Wales only, and from 1 May 1990, covering the whole of the United Kingdom. Approximating these, table 1 summarises by age groups the cases reported for 1980-5 and 1990-5, together with the respective reference populations. Case ascertainment in patients under 70 was 20% higher in 1990-5 than in 1980-5 but more than doubled for patients aged 70 and over; in 1990-5, 2.16 cases per million population aged 70 and over were notified yearly. Comparison of rates of Creutzfeldt-Jakob disease among patients aged 70 and over from comparable prospective surveillance studies elsewhere in the European Union and internationally could give reassurance that there were no reasons other than differentially enhanced ascertainment which accounted for the doubling—such as enhanced dietary exposure to the causative agent of bovine spongiform encephalopathy at older ages in the United Kingdom. More generally, because the incidence of Creutzfeldt-Jakob disease varies so greatly with age group, it makes little sense epidemiologically to present international comparisons of incidence

unless they are age group specific or age standardised. I therefore endorse the decision in France that routine reporting of sporadic cases and the variant forms of Creutzfeldt-Jakob disease should be by 10 year age groups.¹

With only one sporadic case of Creutzfeldt-Jakob disease in a patient under 40 diagnosed yearly in the United Kingdom before 1996, how many suspected cases in patients under 40 were actually referred to the surveillance unit in 1980-5 and 1990-5, of whom 11 were diagnosed as having sporadic Creutzfeldt-Jakob disease? And did the referral pattern change in 1996 because of the emergence of the variant form of the disease? Partial answers are given in a recent letter in the *Lancet* but this concerns a broader age group (under 50 at referral to the surveillance unit) and gives data for April 1996 to February 1997 only.²

In discussing the 14 confirmed cases of the variant form of Creutzfeldt-Jakob disease, all but one occurring in patients under 40 at the onset of clinical illness, Will et al downplay age related exposure to the agent of bovine spongiform encephalopathy as an explanation for the observed age distribution. They identified no exposure related to the bovine disease

MRC Biostatistics Unit, Cambridge CB2 2SR

Sheila M Gore, senior statistician

sheila.gore@mrc-bsu.cam.ac.uk

Table 1 Distribution of cases of Creutzfeldt-Jakob disease by age group and surveillance period

Age group (years)	1980-5 England and Wales			1990-5 United Kingdom			Relative rate	
	Reference population (millions)	Cases of Creutzfeldt-Jakob disease		Reference population (millions)	Cases of Creutzfeldt-Jakob disease			
		No	Yearly incidence/million		No	Yearly incidence/million		
10-39	22.140	7	0.05	23.325	4	0.03	}	1.17
40-49	5.734	3	0.09	7.547	9	0.20		
50-59	5.453	35	1.07	5.964	44	1.23		1.15
60-69	5.080	72	2.36	5.663	98	2.88		1.22
≥ 70	5.308	32	1.00	5.796	75	2.16		2.15

that was sufficiently strongly related to age. However, consumption of certain foods is strongly age related (p 404)³. For example, in 1986-7, burgers and kebabs were consumed by 45% of 16-24 year olds but by only 13% of 50-64 year olds. By contrast, the consumption of beef or even sausages is not age related.⁴

Let us turn now from young adults to farmers. Will et al confirm that the occurrence of sporadic Creutzfeldt-Jakob disease in farmers is more than happenstance.⁵ Molecular marker studies of bovine spongiform encephalopathy reported by Collinge et al effectively proved that the variant form of Creutzfeldt-Jakob disease is caused by the same agent that causes the bovine disease.⁶ Hence as well as transmission studies with brain tissue taken at necropsy from farmers dying of Creutzfeldt-Jakob disease, molecular marker studies will be important in determining whether some of the cases in farmers were likely to have been due to occupational exposure to the infectious agent in cattle. It is worth recalling that in July 1996 (that is, before Collinge's work was reported) the degree of belief in the proposition that bovine spongiform encephalopathy causes the variant form of Creutzfeldt-Jakob disease averaged 5.4 (SD 2.8) on a scale of 0 (nil belief) to 10 (absolute certainty).⁷

Will et al report that in addition to the six cases of Creutzfeldt-Jakob disease in lifelong farmers, there had been three other confirmed cases since 1 May 1990 in people who belonged formerly to the occupational group comprising abattoir workers, butchers, meat cutters, and rendering plant workers. This occupational

group together with veterinarians accounted for no more than $s=78\,000$ workers yearly. The "ever employed" person years in this occupational group to be counted in the surveillance period 1990-5 depend on turnover of staff. If all 78 000 remained in the occupational group throughout 1981-95 they would contribute only $6s=6 \times 78\,000=0.468$ million person years to the surveillance period 1990-5. If two thirds of employees stayed for 15 years, one sixth were replaced every year, and one sixth turned over after three years, then the person years contribution would increase considerably. Serious consideration needs to be given to the proper estimation of "ever employed" and "continuously employed" denominators if full use is to be made of occupational histories collected in cases of Creutzfeldt-Jakob diseases.

- 1 Alperovitch A, Brendel JB, Chalclain J, Delasncio-Laupretre N, Deslys J-P, Dormont D, et al. Situation en France. Incidence de la maladie de Creutzfeldt-Jakob en France. *Bull Epidemiol Hebdom* 1996;48:207-8.
- 2 Will RG, Knight RSG, Zeidler M, Stewart G, Ironside JW, Cousens SN, et al. Reporting of suspect new variant Creutzfeldt-Jakob disease. *Lancet* 1997;349:847-8.
- 3 Gore SM, Bingham S, Day N. Age related dietary exposure to meat products from British dietary surveys of teenagers and adults in the 1980s and 1990s. *BMJ* 1997;315:404-5.
- 4 Gregory JR, Foster K, Tyler H, Wiseman M. *Dietary and nutrition survey of British adults, 1986-7*. London: HMSO, 1990.
- 5 Gore SM. More than happenstance: Creutzfeldt-Jakob disease in farmers and young adults. *BMJ* 1995;311:1416-8.
- 6 Collinge J, Sidle KCL, Meads J, Ironside J, Hill AF. Molecular analysis of prion strain variation and the aetiology of "new variant" CJD. *Nature* 1996;383:685-90.
- 7 Gore SM. Address to the Parliamentary and Scientific Committee. More than happenstance: CJD in farmers and young adults. *Sci Parliament* 1996;53:2-3.

Birth weight and risk of cardiovascular disease in a cohort of women followed up since 1976

Janet W Rich-Edwards, Meir J Stampfer, JoAnn E Manson, Bernard Rosner, Susan E Hankinson, Graham A Colditz, Walter C Willett, Charles H Hennekens

Department of Medicine, Brigham and Women's Hospital and Harvard Medical School, Boston, MA 02215, USA

Division of Preventive Medicine

Janet W Rich-Edwards, instructor

JoAnn E Manson, associate professor

Charles H Hennekens, professor

continued over

BMJ 1997;315:396-400

Abstract

Objective: To examine the association between birth weight and non-fatal adult cardiovascular disease while controlling for potential confounders such as socioeconomic group and adult lifestyle.

Design: Retrospective self report of birth weight in an ongoing longitudinal cohort of nurses followed up by postal questionnaire every two years.

Setting: Nurses' health study, a cohort of 121 700 women followed up since 1976.

Main outcome measures: Non-fatal cardiovascular disease, including myocardial infarction, coronary revascularisation, and stroke.

Results: Among the 70 297 women free of cardiovascular disease at baseline who reported birth weight in the 1992 questionnaire there were 1309 first cases of non-fatal cardiovascular disease. Increasing birth weight was associated with decreasing risk of non-fatal cardiovascular disease. There were 1216 first cases of non-fatal cardiovascular disease among women who were singletons and had been born full

term; their relative risks adjusted for several cardiovascular risk factors were 1.49 (95% confidence interval 1.05 to 2.10) for birth weight < 2268 g (< 5 lb 0 oz); 1.25 (0.98 to 1.61) for birth weight 2268-2495 g (5 lb 0 oz to 5 lb 8 oz); 1.12 (0.98 to 1.27) for birth weight > 2495-3175 g (> 5 lb 8 oz to 7 lb 0 oz); 1.00 (referent) for birth weight > 3175-3856 g (> 7 lb 0 oz to 8 lb 8 oz); 0.96 (0.80 to 1.15) for birth weight > 3856-4536 g (> 8 lb 8 oz to 10 lb 0 oz); and 0.68 (0.46 to 1.00) for birth weight > 4536 g (> 10 lb 0 oz) (P value for trend = 0.0004). The inverse trend was apparent for both coronary heart disease and stroke.

Conclusions: These data provide strong evidence of an association between birth weight and adult coronary heart disease and stroke.

Introduction

As interest has grown in early determinants of cardiovascular disease, it has been hypothesised that indicators of adult cardiovascular risk are manifest