Fortnightly review

A regular review of the long term follow up of depression

J Angst

Depression is a common and disabling psychiatric disorder with a lifetime prevalence in the community estimated at 17%.¹ Outcome studies show that depressive disorders often recur and may become chronic in up to 25% of patients. For these patients the issue is whether they can reasonably expect to recover completely, while the challenge for clinicians is to recognise the course of the disorder and to manage it appropriately. I have reviewed data from long term outcome studies to clarify the issues surrounding recovery from depression and to help identify appropriate management. Studies were selected from a literature search of the full Medline database (1966 to date) and my own collection of publications.

Depression and disability

The medical outcomes study collated data from 11 242 outpatients in the United States.^{2,3} It showed that depressive symptoms, with or without major depressive disorder (*Diagnostic and Statistical Manual of Mental Disorders*, third edition, revised, DSM-III-R), impaired functional ability and wellbeing as much as the most common chronic medical conditions such as diabetes, chronic lung disease, hypertension, and heart disease. After two years' follow up approximately 40% of patients with major depression were still affected and functionally impaired, while those with chronic minor depression (dysthymia) had the worst outcome—54% had had a major depressive episode during this period.³ Partial recovery of functional capacity was the rule.³

The World Health Organisation collaborative study, a cross national study of over 5000 outpatients, examined the relationship between psychiatric status and functional disability in different countries.⁴ The results showed that psychiatric disorder is common in primary care and is associated with substantial functional disability. One of the strongest associations overall was functional disability with major depression (as assessed by the World Health Organisation's primary care version of the composite international diagnostic interview). The relationship between psychopathology and disability was not explained by physical health status.

These studies indicate that many depressed patients have functional disability for some years after an index episode. What then is the prognosis for recovery from depression?

Summary points

Depression limits daily functioning and wellbeing considerably

It often recurs

Up to 25% of patients develop chronic depression

Prophylaxis has been recommended for patients who have had several episodes of major depression

Many depressed patients receive either no treatment or inadequate treatment after an initial episode

Increased attention must be paid to maintenance treatment

Psychiatric
University Hospital,
Postfach 68, Zurich
8, CH-8029
Switzerland

J Angst,
professor of psychiatry

BMJ 1997;315:1143-6

Lifetime prognosis

Recovery rates

Before examining recovery, it is important to define the terms commonly used to describe the course of depressive illness. These are adapted from Frank et al⁵ and are presented in table 1.

LUCY BRISTOW

Table 1 Definition of the terms used according to the Hamilton depression rating scale (HAM-D), the Beck depression inventory (BDI), and the schedule for affective disorders and schizophrenia (SADS)

Term	HAM-D	BDI	SADS
Episode —A period of >D days, during which the patient consistently has full range of symptoms to meet the criteria for the disorder	D=14 days	D=28 days	D=28 days
Partial remission —A period during which an improvement is observed such that the patient no longer has the full range of symptoms, but continues to show more than minimal symptoms			
Response —Point at which partial remission begins			
Full remission —A relatively brief period (>E but <F) during which an improvement is observed such that the patient no longer meets full syndromal criteria for the disorder	E=2 weeks F=6 months	E=3 weeks F=4 months	E=2 weeks F=8 weeks
Recovery —A remission that lasts for F or longer	F=6 months	F=4 months	F=8 weeks
Relapse —A return of symptoms satisfying the full syndrome criteria for an episode which occurs during the period of full or partial remission, but before recovery			
Recurrence —The occurrence of a new episode during recovery			

Adapted from Frank et al.⁵

Two comprehensive reviews have considered recovery in follow up studies in psychiatric settings.^{6 7} Piccinelli and Wilkinson reviewed 51 follow up studies of depressed adults treated in inpatient or outpatient psychiatric services, and noted outcome in terms of recovery, recurrence, and persistent depression.⁷ Average recovery rates were calculated after one year and more than 10 years of follow up, and figures for each study were weighted in relation to the sample size. The authors estimated that 50-60% of patients with major depression recovered at least briefly over the short term (one year) and that up to 90% showed recovery for at least a short time at up to five years.⁷ However, rates fell when sustained recovery was considered—a weighted average of 43% had recovered at one year but only 24% at 10 or more years.⁷

In an earlier review of 29 studies up to the mid 1980s, I considered the clinical course of depressive disorder in relation to diagnostic subgroups.⁶ Bipolar and unipolar affective disorders seemed to show similar rates of recovery.⁶ Depression did not seem to have a worse outcome in the elderly than in younger adults, and 35 to 60% of elderly patients showed good recovery.⁶ In a recent community study 31% of elderly patients without dementia were considered recovered at five year follow up, a figure which agrees broadly with rates reported from clinical studies.⁸

Double depression

An estimated 25% of patients with acute major depression have an underlying chronic minor depression (dysthymia), a condition which has been termed double depression.^{9 10} In these patients it is important to distinguish between recovery from an acute depressive episode to a state of chronic minor depression and recovery to a state with no affective symptoms. Fewer than 40% of patients with double depression became free of symptoms after one or two years of follow up, although recovery from the acute episode of major depression was reported as 83% at one year and 97% at two years.¹⁰ Moreover, dysthymic patients had a higher rate and frequency of relapse than patients with major depression alone and their recovery from the chronic minor depression was slow.¹⁰

Recurrence

Piccinelli and Wilkinson estimated that approximately 25% of patients had a recurrence of major depression within one year of an index episode, and the proportion increased to around 75% at 10 or more

years.⁷ Numerous authors have investigated the duration of intervals between successive episodes of depression. There is wide agreement that the first interval between episodes is longer than subsequent ones^{6 11} and that a later age of onset of depression correlates with shorter first and subsequent intervals.⁶

Chronicity

Chronicity (defined as non-remitting episodes of major depression of at least two years' duration) is well documented, and most large studies report that this occurs in 10-25% of depressed patients.^{6 7 12} Similar rates have been reported in both unipolar depression and bipolar disorder.¹³⁻¹⁵

A prospective five year follow up study of patients with chronic major depression reported that almost 25% had not had two or more symptom free months in this period, and that recovery was substantially delayed in chronic depression.¹⁶ Data from the National Institute of Mental Health collaborative study on the psychobiology of depression indicated that the rate of recovery from an index episode of depression declined over time. It was approximately 70% within one year but 81%, 87%, and 88% in years 2, 4, and 5, respectively.¹⁷

Dysthymia is, by definition, chronic, and DSM-IV criteria for the disorder require the presence of depressed mood for at least two years, although with symptoms less severe than those of major depressive disorder (subsyndromal depression). Despite the reduced severity of symptoms, the functional impairment caused by dysthymia was comparable to that of major depressive disorder,^{18 19} underlining the clinical importance of this condition.

Treatment

Clearly, appropriate treatment strategies are needed. When medication for an acute episode of major depression is stopped too soon after an initial response, symptoms reappear in about 50% of patients (relapse of depression). Most investigators recommend that medication should be continued after remission, until the patient has been free of major symptoms for at least four to six months.²⁰⁻²³ A longer course of maintenance treatment is strongly recommended for patients with residual symptoms and a history of chronic disorder or several depressive episodes, or both. In patients with double depression, intensive treatment after recovery from the acute episode of major depression is necessary, as continued chronic

minor depression in these people seems to predispose them to rapid relapse.¹⁰

Maintenance treatment

The World Health Organisation Mental Health Collaborating Centres have recommended prophylactic maintenance treatment in patients who have had several episodes of major depression over five years²⁴ but the optimum duration of maintenance treatment is still under debate. There are study data supporting two or three years of maintenance treatment,²⁵⁻²⁶ but information on which to base recommendations for longer periods is limited.

The efficacy of medication plus psychotherapy in prophylactic maintenance treatment of depression has been established in a number of controlled studies, and examples of the rate of recurrence reported at up to three years of follow up are given in table 2.²⁷⁻³² Maintenance treatment with antidepressants produces a substantial reduction in the recurrence of depression, and even patients with chronic disorder (chronic major depression, dysthymia and double depression) respond to appropriate medication.³⁰

Psychotherapy

Interpersonal psychotherapy as maintenance treatment was investigated in a three year outcome study in depressed patients with repeated episodes of depression (Pittsburgh maintenance therapies in recurrent depression study).²⁸⁻³¹ Monthly interpersonal psychotherapy sessions designed specifically to improve social adjustment were clearly shown to be a useful adjunct to medication in prolonging the interval between recurrent episodes of depression.

Electroconvulsive treatment

Maintenance electroconvulsive therapy may be practised reasonably regularly but it is rarely reported.³³ In a small study of nine patients with thought or mood disorders, maintenance electroconvulsive therapy given for longer than six months was shown to be effective in producing full or partial remission and was well tolerated in all patients.³⁴ Few clinicians support long term electroconvulsive therapy, although it may prove a useful alternative for patients who cannot tolerate psychotropic medication.³⁵

Psychotherapy versus pharmacotherapy

Interpersonal psychotherapy and cognitive behavioural therapy have been successful in treating acute episodes of major depression but they make heavy demands on resources and are therefore costly options for long term treatment. Scott and Freeman considered treatment outcome and economic costs in outpatients with non-psychotic major depression and concluded that the cost savings of treatment based in primary care outweighed the advantages of specialist treatment.³⁶ Furthermore, the introduction of the selective serotonin reuptake inhibitors, which are as effective as the older tricyclic antidepressants but are often better tolerated by patients, has widened the choice of acceptable drug treatments available to the primary care physician.³²

Duration of maintenance

Investigators agree that maintenance treatment with antidepressants can reduce the risk of recurrence.

Table 2 Rates of recurrence of depression during controlled trials of maintenance treatment

Study	Treatment	Proportion with recurrence
Montgomery et al ²⁷ (1988)*	Pharmacotherapy Placebo	26† 57†
Kupfer ²⁸⁻²⁹ (1992, 1993)* three years before randomisation	Pharmacotherapy Placebo	9‡ 67‡
Kocsis et al ³⁰ (1996) *26 weeks before randomisation	Pharmacotherapy Placebo	15‡ 52‡
Frank et al ³¹ (1990)* ≥20 weeks before randomisation	Pharmacotherapy Interpersonal psychotherapy Pharmacotherapy+interpersonal psychotherapy	22§ 61§ 24§
Reynolds et al ³² (1995) (late life depression)	Pharmacotherapy Interpersonal psychotherapy	20† 50†

*Patients received treatment for the time indicated before being randomised to continue treatment or switch to placebo.

†After one year of follow up.

‡After two years of follow up.

§After three years of follow up.

Therefore, patients at greatest risk need to be identified and prescribed an adequate level of medication. Studies indicate that even three years of treatment with a full dose of an antidepressant may be insufficient to prevent depression recurring when treatment is stopped in these patients, and they may need prophylaxis for five years or longer.²⁹

Indications for maintenance treatment

- Highly recurrent depressive disorder:
Longstanding history of yearly episodes of depression
Short illness cycle length (less than three years)
Several episodes over a five year period
- Severe incapacitating episodes of depression
- History of protracted episodes of depression
- Chronic major depression
- Double depression
- Residual dysthymia

Effective treatment

Although depression is common and disabling, the naturalistic, collaborative study of depression and the depression patient research in European society study showed that depressed patients consistently received either no medication or ineffective dosages of medication.³⁷⁻⁴⁰ More attention must also be paid to maintenance treatment, and the advent of antidepressants that are better tolerated by patients means that patient's compliance with long term treatment strategies has improved.

The serious consequences of depressive disorders indicate the need for swift and effective intervention by doctors. Moreover, they need to consider the recovery of depressed patients in broader terms, paying attention to assessing the effect of depression on the patients' daily functioning as well as the improvement in symptoms.

I thank Dr Christine McKillop for help with a literature search and editorial support.

Funding: None.

Conflict of interest: None.

- 1 Blazer DG, Kessler RC, McGonagle KA, Swartz MS. The prevalence and distribution of major depression in a national community sample: the national comorbidity survey. *Am J Psychiatry* 1994;151:979-6.

- 2 Wells KB, Stewart A, Hays RD, Burnam A, Rogers W, Daniels M, et al. The functioning and well-being of depressed patients. Results from the medical outcomes study. *JAMA* 1989;262:914-9.
- 3 Wells KB, Burnam MA, Rogers W, Hays R, Camp P. The course of depression in adult outpatients. Results from the medical outcomes study. *Arch Gen Psychiatry* 1992;49:788-94.
- 4 Ormel J, VonKorff M, Üstün B, Pini S, Korten A, Oldehinkel T. Common mental disorders and disability across cultures: results from the WHO collaborative study on psychological problems in general health care. *JAMA* 1994;272:1741-8.
- 5 Frank E, Prien RF, Jarrett RB, Keller MB, Kupfer DJ, Lavori PW, et al. Conceptualisation and rationale for consensus definitions of terms of major depressive disorder: remission, recovery, relapse, and recurrence. *Arch Gen Psychiatry* 1991;48:851-5.
- 6 Angst J. Clinical course of affective disorders. In: Helgason T, Daly RJ, eds. *Depressive illness: prediction of course and outcome*. Berlin: Springer-Verlag, 1988:1-47.
- 7 Piccinelli M, Wilkinson G. Outcome of depression in psychiatric settings. *Br J Psychiatry* 1994;164:297-304.
- 8 Kivela S-L. Long-term prognosis of major depression in old age: a comparison with prognosis of dysthymic disorder. *Int Psychogeriatr* 1995;7(suppl):69-82.
- 9 Keller MB, Shapiro RW. "Double depression": superimposition of acute depressive episodes on chronic depressive disorders. *Am J Psychiatry* 1982;139:438-42.
- 10 Keller MB, Lavori PW, Endicott J, Coryell W, Klerman GL. "Double depression": two-year follow-up. *Am J Psychiatry* 1983;140:689-94.
- 11 Haghighat R. Lifelong development of risk of recurrence in depressive disorders. *J Affect Disord* 1996;41:141-7.
- 12 Scott J. Chronic depression. *Br J Psychiatry* 1988;153:287-97.
- 13 Cassano GB, Masetti L, Perugi G, Soriani A, Mignani V, McNair DM, et al. A proposed new approach to the clinical subclassification of depressive illness. *Pharmacopsychiatry* 1988; 21:19-23.
- 14 Marneros A, Deister A, Rohde A. *Affektive, schizoaffektive und schizophrene Psychosen*. Berlin: Springer-Verlag, 1991.
- 15 Angst J, Preisig M. Course of a clinical cohort of unipolar, bipolar and schizoaffective patients. Results of a prospective study from 1959 to 1985. *Schweizer Arch Neurol Psychiatrie* 1995;146:5-16.
- 16 Coryell W, Endicott J, Keller M. Outcome of patients with chronic affective disorder: a five year follow up. *Am J Psychiatry* 1990;147:1627-33.
- 17 Keller MB, Lavori PW, Mueller TI, Endicott J, Coryell W, Hirschfeld RMA, et al. Time to recovery, chronicity and levels of psychopathology in major depression. *Arch Gen Psychiatry* 1992;49:809-16.
- 18 Leader JB, Klein DN. Social adjustment in dysthymia, double depression and episodic major depression. *J Affect Disord* 1996;37:91-101.
- 19 Howland RH. General health, health care utilization, and medical comorbidity in dysthymia. *Int J Psychiatry* 1993;23:211-38.
- 20 Lecrubier Y. Long-term treatment of depression. *Eur Psychiatry* 1993;8(suppl 1):13-16.
- 21 Montgomery SA. Prophylaxis in recurrent unipolar depression; a new indication for treatment studies. *J Psychopharmacol* 1989;3(suppl 2):47-53.
- 22 Prien RF. Efficacy of continuation drug therapy of depression and anxiety: issues and methodologies. *J Clin Psychopharmacol* 1990;10(suppl 3):865-905.
- 23 Thase ME. Relapse and recurrence in unipolar major depression: short-term and long-term approaches. *J Clin Psychiatry* 1990;51:51-7.
- 24 WHO Mental Health Collaborating Centres. Pharmacotherapy of depressive disorders: a consensus statement. *J Affect Disord* 1989;17:197-8.
- 25 Montgomery SA, Montgomery DB. Prophylactic treatment in recurrent unipolar depression. In: Montgomery S, Rouillon F, eds. *Perspectives in psychiatry*. Vol 3. Chichester: J Wiley, 1992:53-79.
- 26 Solomon DA, Bauer MS. Continuation and maintenance pharmacotherapy for unipolar and bipolar mood disorders. *Psychiatr Clin North Am* 1993;16:515-40.
- 27 Montgomery SA, Dufour H, Brion S, Gailledreau J, Laqueille X, Ferrey G, et al. The prophylactic efficacy of fluoxetine in unipolar depression. *Br J Psychiatry* 1988;153(suppl):69-76.
- 28 Kupfer DJ, Frank E, Perel JM, Comes C, Mallinger AG, Thase ME, et al. Five year outcome for maintenance therapies in recurrent depression. *Arch Gen Psychiatry* 1992;49:769-73.
- 29 Kupfer DJ. Management of recurrent depression. *J Clin Psychiatry* 1993;54:29-33.
- 30 Kocsis JH, Friedman RA, Markowitz JC, Leon AC, Miller NL, Gniwesch L, et al. Maintenance therapy for chronic depression. A controlled clinical trial of desipramine. *Arch Gen Psychiatry* 1996;53:769-74.
- 31 Frank E, Kupfer DJ, Perel JM, Comes C, Jarrett DB, Mallinger AG, et al. Three-year outcomes for maintenance therapies in recurrent depression. *Arch Gen Psychiatry* 1990;47:1093-9.
- 32 Reynolds CF, Frank E, Perel JM, Mazumdar S, Kupfer DJ. Maintenance therapies for late life recurrent major depression: research and review circa 1995. *Int Psychogeriatr* 1995;7:27-39.
- 33 Kramer BA. Maintenance ECT: a survey of practice (1986). *Convulsive Therapy* 1987;3:260-8.
- 34 Stiebel VG. Maintenance electroconvulsive therapy for chronic mentally ill patients: a case series. *Psychiatric Serv* 1995;46:265-8.
- 35 Ezion T, Levy A, Levin Y, Neumann M. Should electroconvulsive therapy be used as an ambulatory preventive treatment. *Isr J Psychiatry Relat Sci* 1990;27:168-74.
- 36 Scott AIF, Freeman CPL. Edinburgh primary care depression study: treatment outcome, patient satisfaction, and cost after 16 weeks. *BMJ* 1992;304:883-7.
- 37 Keller MB, Klerman GL, Lavori PW, Fawcett JA, Coryell W, Endicott J. Treatment received by depressed patients. *JAMA* 1982;248:1848-55.
- 38 Keller MB, Lavori PW, Klerman GL, Andreasen NC, Endicott J, Coryell W, et al. Low levels and lack of predictors of somatotherapy and psychotherapy received by depressed patients. *Arch Gen Psychiatry* 1986;43:458-66.
- 39 Keller MB. Undertreatment of major depression. *Psychopharmacol Bull* 1990;24:75-80.
- 40 Lépine J-P on behalf of the DEPRES Steering Committee. European perspective on depression: the patient's view. *Primary Care Psychiatry* 1997;3(suppl 1):S3-6.

(Accepted 19 May 1997)

Lesson of the week

Delayed closure of injuries to the hand caused by blasts helps to preserve function

I Grant

Injuries to the hand caused by blasts are associated with contamination and should not be closed during initial surgery

BMJ 1997;315:1146-7

Proper treatment of hand injuries can greatly reduce the chance of disability in people who are otherwise healthy. The importance of the initial management of such injuries has been recognised by those caring for people who have been wounded in wars over the past 60 years.¹ The hand surgeon Alfred Swanson drew on his experience in the Vietnam war to formulate fundamental rules for managing hand wounds.² He concluded that injuries to the hand caused by a blast are best left open, with primary closure being performed after three to seven days.

An injury caused by a blast introduces a large amount of contaminating material into the tissues. Some material will remain no matter how scrupulously a debridement is performed. The primary closure of blast wounds invites infection.³ During the healing process wounds become oedematous and swell, particularly in

injuries caused by blasts. Tissue injured by a blast may develop a temporary loss of perfusion but retain the capacity for revascularisation if the patient is treated properly. Primary closure of injuries caused by blasts creates localised areas of tension in the tissue adjacent to the sutures. This tension, coupled with the entrapment of fluids within the wound, compromises tissue perfusion and hinders the recovery of damaged tissue. Animal studies have shown that delayed primary closure improves tissue perfusion during the early stages of wound healing even in ischaemic wounds; an ischaemic wound in which closure is delayed for three days has more mechanical strength after 10 days than a wound that is closed at the time of injury.⁴

Injuries caused by blasts are rare in civilian practice.⁵ However, an understanding of their management should not be restricted to those working in war

zones or in countries affected by landmines. Terrorist bombing campaigns, for example, can occur in any part of the world. I describe the consequences of neglecting established principles of management of injuries from blasts.

Case report

A 45 year old farm worker injured his left hand when a tractor tyre exploded while he was changing it. When the tyre exploded during inflation his palm was against the wheel rim and received the force of the blast. He presented to his local accident and emergency department six hours later with a painful, moderately swollen hand. A ragged 1 cm laceration was noted on his palm at the base of his little finger (fig 1). There was no disruption of the normal posture of the hand.

Radiographs showed comminuted fractures of the fourth and fifth metacarpals and the proximal phalanges of the ring and little fingers (fig 2). The fractures were immobilised with percutaneous Kirschner wires under regional anaesthesia. The wound was lavaged and closed. Antibiotics and prophylaxis against tetanus were given and the patient was instructed to keep his hand elevated.

Ten days later the patient returned with fever, purulent discharge from his wound, and signs of infection of the flexor tendon sheath. Further surgery was required. The flexor tendon sheaths were opened and irrigated, and the wounds were thoroughly debrided and left open for primary closure three days later. The patient received intense physiotherapy, yet it was four months before his hand was comfortable and mobile enough for him to return to work on his farm.

Discussion

About one in three injuries seen in hospital accident and emergency departments affect the hand.⁶ The future functioning of the injured hand may depend on the skill with which primary care is administered, yet the initial management of such injuries is often performed by comparatively junior staff. The need for

Fig 1 Injury to patient's hand caused by blast from tractor tyre exploding

Fig 2 Radiograph of comminuted fractures of fourth and fifth metacarpals and proximal phalanges of ring and little fingers

adequate support and education of staff is shown by this case. The failure to correctly identify the severity of the injury and to institute the appropriate initial management almost certainly contributed to the patient's prolonged disability and loss of income. Hand injuries caused by blasts are associated with considerable force and contamination; they should not be closed at the time of the initial surgery.

Blond McIndoe
Centre, Queen
Victoria Hospital,
East Grinstead,
Sussex RH19 3 DZ
I Grant,
surgical research
fellow
mcindoe@dia.
pipex.com

- 1 Brown PW. War wounds of the hand revisited. *J Hand Surg [Am]* 1995;20:S61-7.
- 2 Swanson AB. The treatment of war wounds to the hand. *Clin Plast Surg* 1975;2:615-26.
- 3 Brown PW. The prevention of infection in open wounds. *Clin Orthop* 1973;96:42-50.
- 4 Quirinia A, Viidik A. The effect of delayed primary closure on the healing of ischaemic wounds. *J Trauma Inj Crit Care* 1996;41:1018-23.
- 5 Frykberg ER, Tepas JJ. Terrorist bombings: lessons learned from Belfast to Beirut. *Ann Surg* 1988;208:569.
- 6 Nicoll J, Cole RP, Clarke JA. Audit of emergency throughput in regional plastic surgery unit. *Ann R Coll Surg Engl* 1994;76:161-3.

(Accepted 19 May 1997)

Medical work experience—luxury or necessity?

I am a 16 year old student and over recent years have become interested in pursuing a career in medicine. But I have a message for some sections of the medical establishment about the difficulties I had in finding out more about medicine.

For the past two years I have been studying for my GCSEs. I discovered that most universities require A level chemistry to study medicine. As this might not have been my first choice I decided to check out my vocational hunch in some real work experience.

I talked to my local general practitioner and then wrote to my local community hospital. Neither I or my father could persuade the senior administrator to let me have even the most modest of experiences in the hospital.

Not to be put off, I turned to my grandfather, a well respected former general practitioner on the south coast. He explained my situation to his local hospitals, and as a result I have just completed a wonderful programme of two weeks in a variety of real work environments. This not only enabled me to observe doctors, but also gave me the chance to see how physiotherapists, occupational therapists, biochemists, health visitors, and receptionists worked.

I began in a large hospital, where I was made very welcome and given four different daily programmes. At the start of my second

week I went to a local cottage hospital, followed by two days in a general practitioner's surgery; one looking at the administrative side, and the other shadowing a district nurse. Despite its less glamorous image, this was a complex organisation, which seemed to require more formality and control to make it work effectively.

At the end of the fortnight I was left with some interesting reflections. I found the majority of staff helpful and enthusiastic to share their knowledge and opinions of medicine. It concerned me, however, that so few doctors advised me to study medicine, yet they would not tell me specifically why. I was left wondering about the reason for such scepticism.

I loved the experience of the new world I had been allowed to see, and it will certainly increase my motivation in future study. Hospitals must recognise the importance of work experience. Many seem ignorant of the fact that there are students who want and need to find out more about medical work experience, yet they are denied the chance. If students have a keen interest in medicine, you, the establishment, should give them the chance to test it out at the earliest opportunity.

Victoria Purser lives in Huntingdon