estimated risk of thrombotic stroke in users of second versus third generation pills. The United Kingdom and the continental countries had similar findings. These odds ratios should be assessed against the backdrop of the small absolute risk they entail and in the context of the clear benefits of use of oral contraceptives for women of reproductive age. The annual event rate is between 1-1.6 stroke events per 10 000 women aged 25-44—that is, 1 stroke per 12 000 women. Three strokes per 100 000 women per year may be attributable to the use of oral contraceptives. This risk could be controlled by avoiding prescription of the pill in women who have important cardiovascular risk factors such as high blood pressure and might be lessened by appropriate management of these risk factors.

The investigators were accountable only to the Scientific Reference Board (members listed in reference 1).

Transnational case-control study. Principal investigators: WO Spitzer (Montreal, 1993-5), LAJ Heinemann (Berlin, 1996 to date). Scientific review board: U Bergman (Stockholm), M Breckwoldt (Freiburg), J Collins (Toronto), F Kemper (Munster), J LeLorier (Montreal), S MacLeod (Toronto), K MacRae (London, until May 1995), W Ray (Nashville), J Schlesselman (Pittsburgh). Centre directors: Austria—H Concin (Bregenz), G Stark (Graz), K Pfeiffer (Innsbruck), R Weitgasser (Salzburg); France—B Begaud (Bordeaux), B Boissel (Lyons). Germany—J Haerting, F Lautenschlaeger (Halle), K Püschel (Hamburg), F Hoffmann (Jena), BP Robra (Magdeburg), M Lustermann (Nordhausen), I Schulzki (Schwerin), S Boethig (Zwickau). Switzerland—F Gutzwiller (Zurich). United Kingdom—L de

Caestecker (Glasgow), C McCollum (till August 1995), P Hannaford (from August 1995) (Manchester), R Mann (Southampton). Publication committee: I Guggenmoos-Holzmann (Berlin); R Bruppacher (Basle); M Lewis (Berlin); M Thorogood (London), LAJ Heinemann (Berlin). Data management centre: M Lewis, D Kühl-Habich, C Klindworth, A Assmann (Potsdam) (for more extensive list of collaborators see reference 1).

Funding: Unconditional grant from Schering AG, Berlin.

Conflict of interest: Funding by Schering AG.

- 1 Spitzer WO, Lewis MA, Heinemann LAJ, Thorogood M, McRae KD. Transnational Research Group on Oral Contraception and the Health of Young Women. Third generation oral contraceptives and risk of venous thromboembolic disorders: an international case-control study. *BMJ* 1996;312:83-8.
- 2 Lewis MA, Spitzer WO, Heinemann LAJ, McRae KD, Bruppacher R, Thorogood M. Transnational Research Group on Oral Contraception and the Health of Young Women. Third generation oral contraceptives and risk of venous thromboembolic disorders: an international case-control study. *BMJ* 1996;312:88-90.
- 3 Spitzer WO, Thorogood M, Heinemann L. Trinational case-control study of oral contraceptives and health. *Pharmacoepidemiol Drug Safety* 1993;2:21-31.
- 4 World Health Organisation Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. A multinational case-control study of cardiovascular disease and steroid hormone contraceptives. Description and validation of methods. *J Clin Epidemiol* 1995;48:1513-47.
- 5 Stata. *Statistical software package, version 4.0*. College Station, Tx: Stata, 1995.
- 6 WHO Collaborative Study of Cardiovascular Disease and Steroid Hormone Contraception. Ischaemic stroke and combined oral contraceptives: results of an international, multicentre, case-control study. *Lancet* 1996;348:498-507.
- 7 Heinemann L. Epidemiological assessment of cardiovascular effects of hormonal contraception. In: F Michael, ed. *Safety requirements for contraceptive steroids*. New York: Cambridge University Press, 1989:113-25. (Accepted 29 July 1997)

Underreporting of mortality from RhD haemolytic disease in Scotland and its implications: retrospective review

Charles R Whitfield, Alaeddin Raafat, Stanislaw J Urbaniak

See editorial by van Dijk
Department of Obstetrics and Gynaecology, the Queen Mother's Maternity Hospital, University of Glasgow, Glasgow G3 8SH

Charles R Whitfield, *emeritus regius professor of midwifery*

Academic Transfusion Medicine Unit, Department of Medicine and Therapeutics, University of Aberdeen, Regional Transfusion Centre, Foresterhill, Aberdeen AB9 2ZW

Alaeddin Raafat, *honorary lecturer in transfusion medicine*
Stanislaw J Urbaniak, *honorary reader in transfusion medicine*

Correspondence to: Professor Whitfield

BMJ 1997;315:1504-5

Clarke et al surveyed the decline in RhD haemolytic disease in England and Wales from 1977-92 by reviewing the statistics of the Office of Population Censuses and Surveys; during that time deaths attributable to RhD haemolytic disease had fallen from 18.4 to 1.3 per 100 000 live births.¹ We surveyed RhD alloimmunisation in mothers resident in Scotland between 1987 and 91 and confirmed that fetal loss from RhD haemolytic disease was greater than that listed by the General Register Office in Scotland. Underreporting has serious implications for recognising, understanding, and preventing this potentially lethal disease.

Subjects, methods, and results

Data on deaths attributable to RhD haemolytic disease during 1987-91 were obtained from: death certificates; details of pregnancies in Scottish residents who had antibody to RhD antigen and were managed at the Queen Mother's Hospital; and a retrospective study identifying women in whom RhD antibodies were detected for the first time during 1987-91 (table). The definition of mortality from RhD haemolytic disease included all abortions after fetal death, stillbirths, and neonatal deaths attributed to RhD incompatibility or its treatment and deaths after 28 days of life in which RhD haemolytic disease or its treatment was the primary cause.²

Only four deaths were listed by the General Register Office in Scotland; all were neonatal deaths occurring in women immunised against RhD before 1987. Four deaths occurred in the cohort but none were listed through the General Register Office in Scotland. Twelve other pregnancies managed at the Queen Mother's Hospital during this time (mothers alloimmunised before 1987) resulted in deaths related to RhD haemolytic disease. Ten of the pregnancies ended with abortions after fetal death, and two resulted in liveborn infants treated in utero by repeated blood transfusions. These two infants were delivered electively more than two months preterm but died from complications on the first and 95th days of life.

Thirteen of the 20 deaths resulted from alloimmunisation against RhD antigen during a first pregnancy. In four cases maternal RhD antibodies were first detected within two days of delivery, and in the remaining nine during gestation. There was also good documentation of antepartum immunisation in relation to at least two of the deaths from second pregnancies.

Comment

A review of the case notes of deaths recorded as Rh haemolytic disease in England and Wales showed that disease was often caused by antibodies other than RhD antibody or was wrongly coded. Our data show that

Details of deaths from RhD haemolytic disease during 1987-91 and mother's alloimmunisation

Antibody to RhD first detected			Deaths from RhD haemolytic disease		
Year	Order of pregnancy	Gestation (weeks)	Year	Order of pregnancy	Category of gestation (No of weeks)
1985	2nd	9	1987	3rd	Aborted (23)
1985	1st	39	1987	3rd	Late neonatal death (30)
1983	1st	34	1987	2nd	Early neonatal death (31)
1983	1st	34	1989	3rd	Postneonatal death (31)†
1983	2nd	14	1988	3rd	Early neonatal death (28)
1983	1st	36	1987	3rd	Aborted (27)
1981	2nd	26	1988	4th	Aborted (26)
1988	2nd	31	1988	2nd	Stillborn (31)
1988	2nd	31	1990	3rd	Aborted (19)
1989	1st	29	1989	1st	Stillborn (30)
1979	1st	NK	1989	3rd	Early neonatal death (33)
1979	1st	NK	1990	5th	Aborted (19)
1977	2nd	36*	1989	6th	Aborted (21)
1977	2nd	36*	1990	7th	Aborted (16)
1984	1st	40	1990	3rd	Aborted (25)
1987	1st	33	1990	2nd	Stillborn (31)
1986	1st	40*	1990	5th	Aborted (25)
1986	1st	40*	1991	6th	Aborted (19)
1986	1st	38*	1990	4th	Aborted (23)
1986	1st	38*	1991	5th	Early neonatal death (29)

*Antibody to RhD first detected post partum. †Died 95th day. NK=not known.

five times as many deaths from RhD haemolytic disease were uncertified as were certified through the General Register Office in Scotland. The major cause of underreporting deaths was that certification data exclude abortions. Extrapolation of all the deaths from RhD haemolytic disease identified in Scotland to the population of the United Kingdom would suggest about 50 deaths per year (six per 100 000 live births).

Antepartum prophylaxis against RhD antigen is used widely outside the United Kingdom, and in North America alloimmunisation has been reduced to less than 0.3% of women at risk.³ The standard British dose of 500 international units of anti-D immunoglobulin administered at 28 and 34 weeks' gestation reduced alloimmunisation from 0.95% in historical controls to 0.32%,⁴ and we have recently shown that this approach could generate net savings to the NHS in Scotland by avoiding the costs of intervention in alloimmunised women and their babies.⁵

Our data support the introduction of routine antepartum prophylaxis as 65% (13/20) of deaths from RhD haemolytic disease were the eventual results of antepartum sensitisation in the mothers' first pregnancies and 65% of women seen at the Queen Mother's Hospital were alloimmunised during and not after a pregnancy.²

Funding: This study was financed in part by the Scottish Office Department of Health, project grant K/OPR/2/2D155. Conflict of interest: None.

- Clarke C, Hussey RM. Decline in deaths from Rhesus haemolytic disease of the newborn. *J Roy Coll Phys* 1994;28:310-1.
- Murphy KW, Whitfield CR. Rhesus disease in this decade. *Contemp Rev Obstet Gynaecol* 1994;6:61-7.
- Bowman JM. The prevention of Rh immunization. *Transfusion Med Rev* 1988;2:129-50.
- Thornton JG, Page C, Foote G, Arthur GR, Tovey LAD, Scott JS. Efficacy and long term effects of antenatal prophylaxis with anti-D immunoglobulin. *BMJ* 1989;298:1671-3.
- Vick S, Cairns J, Urbaniak SJ, Whitfield CR, Raafat A. Cost effectiveness of antenatal anti-D prophylaxis. *Health Econ* 1996;5:319-28. (Accepted 11 March 1997)

A case I will never forget

The wrong diagnosis

I will never forget Stan. It is now over 30 years since he died, but the circumstances of his death haunt me still.

He had been unwell for a couple of weeks, so we took him to the vet. Like most Welsh terriers he seemed to have the brains of a rocking horse, but for a couple of weeks his behaviour had been positively bizarre. He was chewing everything in the house: shoes, furniture, golf balls, and even the telephone wire—unhelpful behaviour in a GP's house.

The vet didn't examine him. He didn't need to. "This is typical behaviour that we vets see in dogs who feel unloved. This is his way of saying that you should pay him more attention. Show him more affection and his behaviour will improve."

We slunk off, feeling thoroughly guilty at treating Stan so badly, even though in our heart of hearts we knew that we'd always cared for him as lovingly as we could. From then on we resolved to mend our ways, but his behaviour got worse and worse.

We took him back to the vet. This time a locum was there; he examined the dog carefully, and found that he had acute lymphocytic leukaemia, which was giving him a severe sore throat. The only way that he could ease his discomfort was by chewing unusual textures—like telephone wires and golf balls.

Some lessons you never forget.

David Haslam, *general practitioner, Cambridgeshire*

We welcome filler articles up to 600 words on topics such as *A memorable patient*, *A paper that changed my practice*, *My most unfortunate mistake*, or any other piece conveying instruction, pathos, or humour. If possible the article should be supplied on a disk. Permission is needed from a patient or a relative if an identifiable patient is referred to. We also welcome contributions for "Endpieces," consisting of quotations of up to 80 words (but most are considerably shorter) from any source, ancient or modern, which have appealed to the reader.