Published in final edited form as:

Curr Opin Cell Biol. 2007 October ; 19(5): 508–514.

Making and breaking contacts: The cellular biology of cadherin regulation

Alpha S. Yap¹, Matthew S. Crampton¹, and Jeff Hardin²

¹ *Division of Molecular Cell Biology, Institute for Molecular Bioscience, The University of Queensland, St. Lucia, Brisbane, Queensland, Australia 4072*

² *Laboratory of Genetics, Department of Zoology, University of Wisconsin, Madison, WI53706, USA*

Summary

Cadherin-mediated cell-cell interactions are dynamic processes and cadherin function is tightly regulated in response to cellular context and signaling. Ultimately, cadherin regulation is likely to reflect the interplay between a range of fundamental cellular processes, including surface organization of receptors, cytoskeletal organization and cell trafficking, that are coordinated by signaling events. In this review we focus on recent advances in understanding how interplay with membrane trafficking and other cell-cell junctions can control cadherin function. The endocytosis of cadherins, and their post-internalization fate, influence surface expression and metabolic stability of these adhesion receptors. Similarly, at the surface, components of tight junctions provide a mode of cross-talk that regulates assembly of adherens junctions.

Introduction: cadherin regulation and tissue organization

Cadherin adhesion molecules, notably those of the classical cadherin subfamily, critically determine tissue organization both in health and disease. Cadherin function is dynamic and regulated by developmental and cellular signals. Moreover, cadherins exert their functional impact at several different biological levels, ranging from conferring resistance to detachment, controlling the morphogenesis of contacts as cells integrate into populations, and influencing tissue patterning and cohesion. Thus, it is perhaps not surprising that many signaling pathways and cellular processes affect cadherin biology. In this review we will focus on recent developments in two areas that emphasize how interconnected mechanisms influence cadherin function: membrane trafficking and junctional cross-talk.

Cadherin trafficking and the metabolic fate of cadherins

It has long been recognized that membrane trafficking must link the biosynthesis of cadherins with their eventual degradation in lysosomes. Like other integral membrane proteins, newly-synthesized cadherins are packaged in the Golgi, transported to the cell surface and eventually internalised and degraded [1,2]. But in-between is a complex network of trafficking pathways that provide alternative routes for the endocytosis and subsequent fate of internalised cadherin (Fig 1).

Regulating internalisation: The yin and yang of adhesion and endocytosis

Constitutive endocytosis of a number of classical cadherins has been observed in cells that display apparently stable cell-cell contacts [3,4]. Basal levels of cadherin internalisation would be expected to support their metabolic turnover and perhaps contribute to local remodelling of contacts. Endocytosis can also be substantially altered in response to external stimuli. For example, E-cadherin internalisation is increased when subconfluent epithelial cultures are treated with EGF [5] while VEGF induces the internalisation of VE-cadherin [6•]. Conversely, uptake of N-cadherin in cultured hippocampal neurons was decreased upon stimulation of NMDA receptors [7•]. This stabilized N-cadherin in synapses with the potential to affect synaptic plasticity. Thus, cadherin endocytosis is itself a dynamic process that can be physiologically, and perhaps pathologically, altered depending on cellular context.

Moreover, cadherin internalisation may occur through different pathways. Constitutive endocytosis of E-cadherin and VE-cadherin appears to occur via a clathrin-mediated pathway [3,8•,9], but a number of clathrin-independent pathways for cadherin endocytosis have also been identified [5,10,11]. Notably, macropinocytosis is implicated in EGF-stimulated internalisation of E-cadherin in subconfluent epithelial cells [5]. Different pathways may then dominate in different cellular contexts; it is further possible that cadherin endocytosis may be regulated by activating different uptake mechanisms. The extent to which this occurs remains to be defined. Even so, it is clear that cadherin internalization can be influenced by negative and positive mechanisms.

Perhaps the best characterized inhibitor of cadherin endocytosis is its associated protein, p120-ctn. One major mechanism for p120-ctn to support cadherin adhesion is by inhibiting clathrin-mediated endocytosis [8•,12], thereby stabilizing cadherins at the cell surface [13,14]. Additionally, Rac activity and actin cytoskeletal reorganization through the actin cross-linker, IQGAP, inhibited internalisation in a cell-free model of E-cadherin endocytosis [9]. How these molecules block cadherin endocytosis, and whether they are mechanistically linked, remains unknown. One possibility is that dynamic actin reorganization is necessary to facilitate clathrin-mediated endocytosis [15], consistent with the observation that actin-disrupting agents promoted E-cadherin endocytosis [9]. Interestingly, p120-ctn can regulate Rac signaling with the potential to signal to the actin cytoskeleton [16].

Conversely, endocytosis is a complex, multi-step process that involves discrete events such as cargo selection, budding and fission of vesicles that are subject to tight regulation [17]. A key step is the recognition of target proteins by the endocytic machinery. The cytoplasmic tails of several classical cadherins bear a number of di-leucine and tyrosine-based motifs that are potentially recognized by adaptor proteins of the clathrin-based endocytic apparatus [1,2]. Agents that inhibit cadherin endocytosis, such as p120-ctn [2,12], might then work by preventing these motifs being recognized and bound by the endocytic machinery. Alternatively, signals that can down-regulate cadherin function might stimulate cadherin internalisation by inducing pro-endocytic modifications to the cytoplasmic tail. A striking example occurs in endothelia, where VEGF activates a signaling cascade involving Rac and PAK that leads to the phosphorylation of a conserved serine motif in the cytoplasmic tail of VE-cadherin [6•]. This then recruits β -arrestin leading to the clathrin-dependent uptake of the cadherin. Monoubiquitylation can also promote endocytosis [18] and the E3 ligase, Hakai, binds E-cadherin in a tyrosine kinase-dependent fashion [19], providing an alternative mechanism to target cadherins for internalization. At this point it is noteworthy that Hakai did not bind N-cadherin; this emphasizes that while much work has been done with E-cadherin, the trafficking of other cadherins may have significant differences. Indeed, the serine residue necessary for β -arrestin-mediated internalisation of VE-cadherin is not found in either E- or N-cadherin [6•].

Net cadherin internalisation is thus likely to reflect the balance between signals that promote endocytosis and those that inhibit it. This then raises the question whether there are dominant factors that control its internalisation. Indeed, cadherin adhesion itself appears to be one such factor. This was first suggested by the observation that cadherin internalisation increased when cell-cell contacts were broken by chelating extracellular calcium [3,20]. Similarly, EGF induced internalisation of E-cadherin by macropinocytosis in subconfluent, but not confluent, cultures [5]. Consistent with this, Izumi et al. (2004) reported that it was the free cadherin, rather than the cadherins engaged in adhesion, that underwent endocytosis in cell-free assays. Moreover, homophilic ligation of cellular cadherins with recombinant ligands inhibited internalisation in their experiments [9]. Together, these observations suggest that productive adhesive ligation of cadherins prevents their endocytosis, perhaps by activating Rac signaling and remodelling the actin cytoskeleton. Adhesion and endocytosis would then be closely opposed processes: Endocytosis would tend to decrease cadherin function by removing free cadherins from the cell surfaces, making them unavailable to engage in adhesive interactions. Productive cadherin ligation, on the other hand, would stabilize the cadherin by preventing endocytosis, thereby prolonging its residence time on the cell surface.

Alternatively, the Troyanovsky group recently proposed that endocytosis actively inhibits the cadherin adhesive machinery. The key observations were that maneuvers, such as hypertonic stress, that inhibit cadherin internalisation prevented the disassembly of cadherin *trans*-dimers that occurred upon chelation of extracellular calcium [21•]. Such *trans*-dimers are thought to reflect the adhesive interactions of cadherins on neighbouring cells, implying that endocytosis targets the adhesive machinery itself. This contrasts with the first model, where endocytosis would target the unliganded cadherin. It should be noted, however, that many non-specific maneuvers that inhibit endocytosis can influence cadherin function by alternative mechanisms. For example, hypertonic stress induces tyrosine phosphorylation of cortactin [22], which contributes to cadherin junction integrity [23,24] (Helwani et al., unpublished). These disparate models thus remain to be resolved and warrant further investigation. Nonetheless, they both emphasize the close functional and mechanistic relationship between cadherin adhesion and endocytosis.

Controlling the fate of cadherins after internalization

Once internalised, cadherins may either be recycled back to the cell surface or trafficked to late endosomes and lysosomes for degradation [1]. Traffic along these disparate pathways is also subject to cellular regulation [17]. Traffic to lysosomes entails passage through Rab5-enriched and Rab7-enriched compartments and dominant-negative mutants of either of these GTPases block transport of E-cadherin to lysosomes [25•]. In contrast, membrane proteins can be recycled to the plasma membrane by two routes: directly from sorting endosomes shortly after internalisation or after transport to the recycling endosome, a longer-lived compartment that is characteristically enriched in the Rab11 GTPase [26,27•]. Rab11 and the exocyst complex appear to participate in a pathway for cadherin recycling. Disruption of the *sec5* subunit of the exocyst caused *Drosophila* E-cadherin to accumulate in Rab11-enriched endosomes [28•], while Rab11 interacts with components of the exocyst providing a potential molecular basis for this functional relationship [28•,29•]. Thus transport through recycling endosomes appears to constitute one, but not necessarily the sole, pathway for cadherin recycling. Subsequent transport to the cell surface is likely to involve microtubule-based transport [30] and phosphoinositide signaling [31]. Interestingly, in mammalian and *Drosophila* epithelial cells, the recycling pathway may be important for targeted delivery of E-cadherin to the cell surface. Expression of Rab11 mutants caused the apical mis-expression of E-cadherin in MDCK cells [27•] while disruption of the *sec5* subunit of the exocyst in flies prevented transcytosis of DE-cadherin to adherens junctions in the dorsal thorax of the pupae [28•].

Faced with such disparate alternative pathways, what determines which route is taken after cadherins are internalised (Fig 1)? One possibility is that different uptake pathways lead to different fates. For example, uptake of EGF receptor by clathrin-mediated endocytosis may principally mediate recycling, while clathrin-independent uptake may serve as the major pathway for its degradation [32]. However, E-cadherin that was internalised by EGF-induced macropinocytosis appeared to be recycled to the cell surface [5]. Moreover, cadherins that are internalised by either clathrin-mediated or clathrin-independent entry mechanisms often appear in early endosomes that stain for EEA-1 [4,11]. This suggests that cadherins may enter a common compartment shortly after endocytosis, implying that the choice of fates is decided at, or after, exit from this early endosomal compartment.

Recent analysis of how Src signaling affects E-cadherin transport may be informative in analysing how such decisions may be made. Palacios et al. (2005) demonstrated that activation of signaling by a temperature-sensitive (ts) Src mutant downregulated E-cadherin expression in MDCK epithelial cells [25•]. This involved shuttling of internalised cadherin to lysosomes for degradation. Thus one mechanism for mutant Src to inhibit E-cadherin would be to promote its traffic for degradation, presumably at the expense of recycling to the cell surface. Interestingly, ts-Src affected multiple steps along the pathway leading to lysosomal degradation, including promoting GTP loading of Rab 5 and Rab 7 [25•]. As well, the authors reported that mutation of a predicted ubiquitylation site in the cadherin cytoplasmic domain prevented cadherin degradation without affecting its initial endocytosis. This suggests that for E-cadherin, like other membrane proteins [18], ubiquitylation may be an important signal for transport to lysosomes.

These observations thus carry two implications. Firstly, they suggest that there may be overarching cell signals that determine the post-endocytic fate of cadherins to define the choice between recycling and lysosomal shuttling. These will ultimately influence the metabolic half-life of the cadherin and its steady-state levels in the cell, independent of any changes in gene expression or protein synthesis. Based on the example of other membrane proteins, the decision to follow a path to lysosomal degradation may be key, as recycling is often a default pathway [17]. Secondly, such overarching signals may act by targeting multiple regulatory points found along the trafficking pathway of the cadherin, rather than necessarily at a single decision point, thereby biasing traffic along a pathway.

Taken together, cadherin trafficking – notably regulation of its post-endocytic fate – provides a very attractive mechanism to influence cadherin function. Indeed, it has been invoked in a number of morphogenetic processes where dynamic remodelling of cell-cell interactions is thought to happen [33•,34•]. Several limits to our current knowledge need to be noted, however. Firstly, it is difficult to specifically regulate cadherin trafficking. Our armamentarium to date has been limited to global manipulation of molecules such as Rab 5 and sec 5, whose activity is not specific for cadherin trafficking. Thus, the functional significance of altered cadherin traffic seen in these circumstances is formally correlative, rather than causal. Ongoing efforts to define the molecular mechanisms of cadherin traffic will be essential for progress in this area.

Secondly, we do not yet know how extensive is the contribution of regulated trafficking to dynamic control of cadherin function. Cadherin trafficking has the clear potential to affect both the surface expression and metabolic stability of cadherins. However, in vertebrate cells, cadherin adhesion can be significantly reduced without demonstrable changes in either the total or surface levels of cadherin, as occurs when either Myosin II or microtubule dynamics are acutely perturbed [35,36]. Changes in cadherin trafficking are unlikely to be responsible for the altered cadherin function seen under these circumstances. A key parameter may be the biological time scale. Functionally significant alterations in cadherin trafficking have been

most commonly observed associated with changes in cell state (such as induction of epithelial-to-mesenchymal transformation by expression of mutant Src) or during morphogenesis in fast-developing organisms, such as *Drosophila*. It is possible that altered cadherin trafficking principally regulates cadherin function on relatively longer time scales, whereas rapid regulation occurs by other mechanisms, such as lateral clustering or cytoskeletal regulation.

Finally, although we have focused on the trafficking of the cadherin adhesion molecule itself, it is also quite plausible that membrane trafficking mediates the local delivery and turnover of other molecules necessary for cadherin biology. E-cadherin can bind elements of the exocyst complex [28•], providing the capacity for it to mark sites for vesicle docking and delivery [29•,37]. Exocytosis can deliver molecules such as Rho GTPases to the cell surface [32], with the potential to regulate cell signaling and cytoskeletal organization at cadherin adhesions. It is therefore possible that, rather than solely controlling cadherin transport and expression, membrane traffic could also affect cadherin function through the accessory proteins that functionally couple signaling and cytoskeletal activity to cadherin adhesion.

Tight junctions and cadherin regulation: cross-talk at the surface

Just as there is cross talk between trafficking and cadherin function, similar functional interplay occurs between junctions at the cell surface. It has long been thought that cadherin function contributes to the assembly of other specialized cell-cell junctions, notably elements of the apical junctional complex in epithelia (tight junctions, gap junctions, desmosomes) [38]. But, as well, there is increasing evidence to suggest that tight junction (TJ) components influence cadherin-based adherens junctions (AJ).

This is exemplified by the ZO (Zonula occludens) family of MAGUK scaffolds, peripheral membrane proteins that were amongst the first components of the TJ to be identified [39]. That these are necessary for TJ assembly was recently established by the Tsukita group who derived epithelial cells lacking all members of this family (ZO1-3) in which TJs are totally absent [40]•. Surprisingly, although cadherin contacts appeared intact in fully established monolayers of ZO-deficient cells [40]•, cells also showed a defect in the efficiency with which they formed AJ [41]•. In a Ca^{2+} switch assay, these cells show a delay in the transition from spot-like AJs to more belt-like structures with associated linear actin cables. Initial accumulation of spots occurred with normal kinetics, along with recruitment of afadin and Par-3/Par-6/Tiam-1, suggesting that the defects are specific to junctional maturation. That this requirement for ZO proteins may be specific to AJ-related functions was bolstered by structure-function studies, which suggested that a distinct region of ZO-1 is required for spot-to-belt maturation as opposed to claudin polymerisation [41]•. This suggested a role for ZO proteins in the dynamic process of AJ maturation.

A similar theme emerges from recent work on tracheal development in *Drosophila*, whose genome is predicted to contain one orthologue of ZO-1, *polychaetoid/pyd* [42]. Interestingly, Pyd seems to be important in determining the fate of tracheal cells that normally undergo branch fusion [43]•, a role consistent with studies in vertebrates, where ZO-1 appears to sequester the proliferation-promoting transcription factor, ZONAB (reviewed in [44]). In addition, however, *pyd* mutants show defective rearrangement of tracheal cells [43]•. Dynamic analysis with a Pyd::GFP construct reveals that it localizes to AJs. Taken together, these data suggest that Pyd regulates remodeling of AJs during tracheal morphogenesis.

How, then, might TJ components, and perhaps TJs themselves, affect cadherin function and AJ assembly? Although it is still early, two possibilities already present themselves. Firstly, such cross-talk may represent interactions between cadherin complexes and TJ components in early cell-cell adhesions, before definitive junctions have formed. ZO-1 can bind α -catenin and

has often been seen to co-localize with E-cadherin in early spot-like cell-cell contacts. Only later, as TJs form, does ZO-1 shift apically [45]. The ZO-1-binding region of α -catenin appears to be important for strong cadherin adhesion [46]. Moreover, ZO-1 can bind actin filaments [47] and could possibly serve as a cadherin-associated protein to link nascent adhesions to the actin cytoskeleton. Along similar lines, MAGI (MAGUK with inverted domain structure)-1, is also found both in nascent adhesions and in TJ. Recently, Sakurai et al [48]•, reported that MAGI-1 accumulates at VE-cadherin contacts, coinciding with activation of the small GTPase Rap1, which can regulate cadherin adhesions [49]. Of note, MAGI-1 can bind the Rap1 activator, PDZ-GEF and also associate with the actin-binding proteins, synaptopodin and α -actinin-4, providing other ways of linking to the cytoskeleton.

Alternatively, definitive TJ components may affect the biogenesis or maintenance of cadherin contacts themselves. Here it is noteworthy that the polarity determinants, *Drosophila* Stardust (Std) and its mammalian orthologue, PALS1/MPP5, are found in septate junctions and TJs, respectively [50]. These are best understood for their roles in determining apical polarity, but it has been known for many years that *Std* mutants also have defects in AJs [51,52]. Very recent work has uncovered a similar role for MPP5/PALS1 in zebrafish cardiac development [53]; loss of MPP5 function perturbs the cohesion of cardiac precursors, again suggesting that MPP5 somehow affects adhesion. How might a complex that does not physically colocalize with AJs in mature epithelia regulate AJ formation? Recent work from the Margolis laboratory using a strong PALS1 knockdown (KD) cell line suggests one clue [54]•. AJ, as well as TJ, are perturbed in PALS1(KD) cells and AJ components accumulate in cadherin-positive vesicles that appear to fail to undergo exocytic delivery to the surface. Moreover, exocyst components are also mislocalized in the KD cells. Thus depletion of this TJ component appears to affect E-cadherin exocytosis. An interesting speculation is that MPP5/PALS1 spatially regulates the exocyst, perhaps forming with Stardust/Crumbs/PATJ part of a “nascent polarizing” apparatus upstream of the exocyst that is required when cells first make cell-cell contacts. Dynamic analysis of the Crumbs complex vis-à-vis TJ and AJ components in nascent epithelia would help to resolve this question.

Conclusions

In summary, we sought in this review to highlight the ways in which cadherin function is determined by, and regulated through, the interplay between the adhesion receptor and the cellular systems of membrane traffic and tight junctions. Of course, these are not the only processes that influence cadherin biology. We have elided issues such as surface clustering of receptors, cytoskeletal activity and cell signaling in our discussion. Here it may be useful to emphasize that cadherin function encompasses biophysical, cellular and tissue levels. Nor need these be commensurate: for example, inhibiting Arp2/3-dependent actin nucleation perturbs the morphogenesis of cadherin contacts without significantly affecting surface adhesion [55]. Thus different cellular processes may regulate distinct functional parameters as cells use cadherins to interact with one another. Ultimately, a comprehensive understanding of cadherin regulation will need to encompass this diversity, both of effector mechanisms and biological outcomes.

Acknowledgements

We thank all the members of our laboratories for their support, advice and good cheer. We especially thank Tina Tuskey for helpful discussions regarding ZO proteins and Robert Parton and Jenny Stow for trying to teach us about membrane traffic. J.H. is supported by NIH grant GM58038; A.S.Y. and M.S.C. are funded by the NHMRC (Australia).

References

1. Bryant DM, Stow JL. The ins and outs of E-cadherin trafficking. *Trends Cell Biol* 2004;14:427–434. [PubMed: 15308209]
2. Xiao K, Oas RG, Chiasson CM, Kowalczyk AP. Role of p120-catenin in cadherin trafficking. *Biochim Biophys Acta* 2007;1773:8–16. [PubMed: 16949165]
3. Le TL, Yap AS, Stow JL. Recycling of E-cadherin: a potential mechanism for regulating cadherin dynamics. *J Cell Biol* 1999;146:219–232. [PubMed: 10402472]
4. Xiao K, Allison DF, Kottke MD, Summers SA, Sorescu GP, Faundez V, Kowalczyk AP. Mechanisms of VE-cadherin processing and degradation in microvascular endothelial cells. *J Biol Chem* 2003;278:19199–19208. [PubMed: 12626512]
5. Bryant DM, Kerr MC, Hammond LA, Joseph SR, Mostov KE, Teasdale RD, Stow JL. EGF induces macropinocytosis and SNX1-modulated recycling of E-cadherin. *J Cell Sci* 2007;120:1818–1828. [PubMed: 17502486]
- 6. Gavard J, Gutkind JS. VEGF controls endothelial-cell permeability by promoting the beta-arrestin-dependent endocytosis of VE-cadherin. *Nat Cell Biol* 2006;8:1223–1234. [PubMed: 17060906]
This paper identifies a pathway for stimulated endocytosis of VE cadherin in endothelial cells. The authors demonstrate that VEGF induces cadherin endocytosis. They characterize a Rac/PAK pathway that phosphorylates a conserved motif in the cadherin cytoplasmic tail, leading to β -arrestin binding and internalization.
- 7. Tai C-Y, Mysore SP, Chiu C, Schuman EM. Activity-regulated N-cadherin endocytosis. *Neuron* 2007;54:771–785. [PubMed: 17553425]As an alternative mode of regulating cadherin endocytosis, the authors report that synaptic activity inhibits the uptake of N-cadherin in neurons, providing a mechanism to modulate synaptic strengthening.
- 8. Xiao K, Garner J, Buckley KM, Vincent PA, Chiasson CM, Dejana E, Faundez V, Kowalczyk AP. p120-Catenin Regulates Clathrin-dependent Endocytosis of VE-Cadherin. *Mol Biol Cell* 2005;16:5141–5151. [PubMed: 16120645]This paper demonstrates that p120-ctn inhibits VE-cadherin endocytosis, providing a mechanism to explain the capacity of p120-ctn to stabilize surface expression of cadherin.
9. Izumi G, Sakisaka T, Baba T, Tanaka S, Morimoto K, Takai Y. Endocytosis of E-cadherin regulated by Rac and Cdc42 small G proteins through IQGAP1 and actin filaments. *J Cell Biol* 2004;166:237–248. [PubMed: 15263019]
10. Akhtar N, Hotchin NA. Rac1 regulates adherens junctions through endocytosis of E-cadherin. *Mol Biol Cell* 2001;12:847–862. [PubMed: 11294891]
11. Paterson AD, Parton RG, Ferguson C, Stow JL, Yap AS. Characterization of E-cadherin endocytosis in isolated MCF-7 and chinese hamster ovary cells: the initial fate of unbound E-cadherin. *J Biol Chem* 2003;278:21050–21057. [PubMed: 12657640]
12. Miyashita Y, Ozawa M. Increased internalization of p120-uncoupled E-cadherin and a requirement for a dileucine motif in the cytoplasmic domain for endocytosis of the protein. *J Biol Chem* 2007;282:11540–11548. [PubMed: 17298950]
13. Davis MA, Ireton RC, Reynolds AB. A core function for p120-catenin in cadherin turnover. *J Cell Biol* 2003;163:525–534. [PubMed: 14610055]
14. Xiao K, Allison DF, Buckley KM, Kottke MD, Vincent PA, Faundez V, Kowalczyk AP. Cellular levels of p120 catenin function as a set point for cadherin expression levels in microvascular endothelial cells. *J Cell Biol* 2003;163:535–545. [PubMed: 14610056]
15. Qualmann B, Kessels MM. Endocytosis and the cytoskeleton. *Int Rev Cytol* 2002;220:93–144. [PubMed: 12224553]
16. Goodwin M, Kovacs EM, Thoreson MA, Reynolds AB, Yap AS. Minimal mutation of the cytoplasmic tail inhibits the ability of E-cadherin to activate Rac but not phosphatidylinositol 3-kinase: direct evidence of a role for cadherin-activated Rac signaling in adhesion and contact formation. *J Biol Chem* 2003;278:20533–20539. [PubMed: 12672818]
17. Maxfield FR, McGraw TE. Endocytic recycling. *Nat Rev Mol Cell Biol* 2004;5:121–132. [PubMed: 15040445]

18. d'Azzo A, Bongiovanni A, Nastasi T. E3 ubiquitin ligases as regulators of membrane protein trafficking and degradation. *Traffic* 2005;6:429–441. [PubMed: 15882441]
19. Fujita Y, Krause G, Scheffner M, Zechner D, Leddy HE, Behrens J, Sommer T, Birchmeier W. Hakai, a c-Cbl-like protein, ubiquitinates and induces endocytosis of the E-cadherin complex. *Nat Cell Biol* 2002;4:222–231. [PubMed: 11836526]
20. Kamei T, Matozaki T, Sakisaka T, Kodama A, Yokoyama S, Peng Y-F, Nakano K, Takaishi K, Takai Y. Coendocytosis of cadherin and c-Met coupled to disruption of cell-cell adhesion in MDCK cells - regulation by Rho, Rac and Rab small G proteins. *Oncogene* 1999;18:6776–6784. [PubMed: 10597286]
- 21. Troyanovsky RB, Sokolov EP, Troyanovsky SM. Endocytosis of cadherin from intracellular junctions is the driving force for cadherin adhesive dimer disassembly. *Mol Biol Cell* 2006;17:3484–3493. [PubMed: 16760429] This report presents provocative evidence to suggest that endocytosis may actively inhibit molecular mechanisms responsible for cadherin adhesion.
22. Kapus A, Szaszi K, Sun J, Rizoli S, Rotstein OD. Cell shrinkage regulates Src kinases and induces tyrosine phosphorylation of cortactin, independent of the osmotic regulation of Na⁺/H⁺ exchangers. *J Biol Chem* 1999;274:8093–8102. [PubMed: 10075710]
23. Helwani FM, Kovacs EM, Paterson AD, Verma S, Ali RG, Fanning AS, Weed SA, Yap AS. Cortactin is necessary for E-cadherin-mediated contact formation and actin reorganization. *J Cell Biol* 2004;164:899–910. [PubMed: 15024035]
24. El Sayegh TY, Arora PD, Fan L, Laschinger CA, Greer PA, McCulloch CA, Kapus A. Phosphorylation of N-cadherin-associated cortactin by Fer kinase regulates N-cadherin mobility and intercellular adhesion strength. *Mol Biol Cell* 2005;16:5514–5527. [PubMed: 16176974]
- 25. Palacios F, Tushir JS, Fujita Y, D'Souza-Schorey C. Lysosomal targeting of E-cadherin: a unique mechanism for the down-regulation of cell-cell adhesion during epithelial to mesenchymal transitions. *Mol Cell Biol* 2005;25:389–402. [PubMed: 15601859] This paper demonstrates that down-regulation of cadherin expression by constitutively-active Src may occur at several points in the itinerary of cadherin endocytosis to the lysosome, implying that CA Src may bias cadherin traffic towards a degradative fate.
26. Ullrich O, Reinsch S, Urbe S, Zerial M, Parton RG. Rab11 regulates recycling through the pericentriolar recycling endosome. *J Cell Biol* 1996;135:913–924. [PubMed: 8922376]
- 27. Lock JG, Stow JL. Rab11 in Recycling Endosomes Regulates the Sorting and Basolateral Transport of E-Cadherin. *Mol Biol Cell*. 2005
- 28. Langevin J, Morgan MJ, Sibarita JB, Aresta S, Murthy M, Schwarz T, Camonis J, Bellaiche Y. Drosophila exocyst components Sec5, Sec6, and Sec15 regulate DE-Cadherin trafficking from recycling endosomes to the plasma membrane. *Dev Cell* 2005;9:355–376. [PubMed: 16224820]
- 29. Beronja S, Laprise P, Papoulas O, Pelikka M, Sisson J, Tepass U. Essential function of Drosophila Sec6 in apical exocytosis of epithelial photoreceptor cells. *J Cell Biol* 2005;169:635–646. [PubMed: 15897260] Collectively, these three papers provide important insights into the molecular mechanisms and itinerary necessary for cadherin trafficking both in mammalian cells and in Drosophila embryos. In particular, they identify a Rab11 compartment that may participate both in the biosynthetic pathway and for cadherin recycling.
30. Mary S, Charrasse S, Meriane M, Comunale F, Travo P, Blangy A, Gauthier-Rouviere C. Biogenesis of N-Cadherin-dependent Cell-Cell Contacts in Living Fibroblasts Is a Microtubule-dependent Kinesin-driven Mechanism. *Mol Biol Cell* 2002;13:285–301. [PubMed: 11809840]
31. Ling K, Bairstow SF, Carbonara C, Turbin DA, Huntsman DG, Anderson RA. Type I{gamma} phosphatidylinositol phosphate kinase modulates adherens junction and E-cadherin trafficking via a direct interaction with {micro}1B adaptin. *J Cell Biol* 2007;176:343–353. [PubMed: 17261850]
32. Polo S, Di Fiore PP. Endocytosis conducts the cell signaling orchestra. *Cell* 2006;124:897–900. [PubMed: 16530038]
- 33. Classen AK, Anderson KI, Marois E, Eaton S. Hexagonal packing of Drosophila wing epithelial cells by the planar cell polarity pathway. *Dev Cell* 2005;9:805–817. [PubMed: 16326392]
- 34. Ulrich F, Krieg M, Schotz EM, Link V, Castanon I, Schnabel V, Taubenberger A, Mueller D, Puech PH, Heisenberg CP. Wnt11 functions in gastrulation by controlling cell cohesion through Rab5c and E-cadherin. *Dev Cell* 2005;9:555–564. [PubMed: 16198297] Together, these papers identify

membrane trafficking pathways, involving Rab 11 and Rab 5, as critical for morphogenetic cellular movements. They further demonstrate coordinated changes in cadherin, providing organismal evidence that regulated trafficking of cadherin may play an important role in morphogenesis.

35. Shewan AM, Maddugoda M, Kraemer A, Stehbins SJ, Verma S, Kovacs EM, Yap AS. Myosin 2 Is a Key Rho Kinase Target Necessary for the Local Concentration of E-Cadherin at Cell-Cell Contacts. *Mol Biol Cell* 2005;16:4531–4532. [PubMed: 16030252]
36. Stehbins SJ, Paterson AD, Crampton MS, Shewan AM, Ferguson C, Akhmanova A, Parton RG, Yap AS. Dynamic microtubules regulate the local concentration of E-cadherin at cell-cell contacts. *J Cell Sci* 2006;119:1801–1811. [PubMed: 16608875]
37. Yeaman C, Grindstaff KK, Nelson WJ. Mechanism of recruiting Sec6/8 (exocyst) complex to the apical junctional complex during polarization of epithelial cells. *J Cell Sci* 2004;117:559–570. [PubMed: 14709721]
38. Gumbiner B, Stevenson B, Grimaldi A. The role of the cell adhesion molecule uvomorulin in the formation and maintenance of the epithelial junctional complex. *J Cell Biol* 1988;107:1575–1587. [PubMed: 3049625]
39. Stevenson BR, Siliciano JD, Mooseker MS, Goodenough DA. Identification of ZO-1: a high molecular weight polypeptide associated with the tight junction (zonula occludens) in a variety of epithelia. *J Cell Biology* 1986;103:755–766.
- 40. Umeda K, Ikenouchi J, Katahira-Tayama S, Furuse K, Sasaki H, Nakayama M, Matsui T, Tsukita S, Furuse M, Tsukita S. ZO-1 and ZO-2 independently determine where claudins are polymerized in tight-junction strand formation. *Cell* 2006;126:741–754. [PubMed: 16923393] This work provides the first full analysis of cultured cells depleted of all three Zonula Occludens (ZO) proteins (ZO-1, ZO-2, ZO-3), and provides clear evidence for their functional importance in formation of tight junctions (TJs) in at least some types of cultured cells. Surprisingly, ZO- cells are well polarized, but lack TJs, a defects that can be rescued by expression and forced dimerization of a membrane-tethered form of ZO-1.
- 41. Ikenouchi J, Umeda K, Tsukita S, Furuse M, Tsukita S. Requirement of ZO-1 for the formation of belt-like adherens junctions during epithelial cell polarization. *J Cell Biol* 2007;176:779–786. [PubMed: 17353356] This study shows that, in addition to their effects on tight junction (TJ) formation, the Zonula Occludens proteins ZO-1 and ZO-2 are also involved in the transition from point-like spots of cadherin-based adhesion to belt-like adherens junctions (AJs) in vitro. Rac1 activation is surprisingly low in the ZO- cells, but can be rescued by a fragment of ZO-1 different from that required for formation of TJs.
42. Chen CM, Freedman JA, Bettler DR, Manning SD, Giep SN, Steiner J, Ellis HM. Polychaetoid is required to restrict segregation of sensory organ precursor from proneural clusters in *Drosophila*. *Mech Dev* 1996;57:215–227. [PubMed: 8843398]
43. Jung AC, Ribeiro C, Michaut L, Certa U, Affolter M. Polychaetoid/ZO-1 is required for cell specification and rearrangement during *Drosophila* tracheal morphogenesis. *Curr Biol* 2006;16:1224–1231. [PubMed: 16782014] This paper identifies functional roles for the *Drosophila* ZO-1 homologue *polychaetoid* (pyd), which was identified using a gene chip approach to find targets of the transcription factor *trachealess*, in the tracheal system. pyd is involved in cell fate specification, but it is also involved in junctional remodeling during tracheal morphogenesis.
44. Matter K, Aijaz S, Tsapara A, Balda MS. Mammalian tight junctions in the regulation of epithelial differentiation and proliferation. *Curr Opin Cell Biol* 2005;17:453–458. [PubMed: 16098725]
45. Ando-Akatsuka Y, Yonemura S, Itoh M, Furuse M, Tsukita S. Differential behaviour of E-cadherin and occludin in their colocalization with ZO-1 during the establishment of epithelial cell polarity. *J Cellular Physiol* 1999;179:115–125. [PubMed: 10199550]
46. Itoh M, Nagafuchi A, Moroi S, Tsukita S. Involvement of ZO-1 in cadherin-based cell adhesion through its direct binding to α Catenin and actin filaments. *J Cell Biol* 1997;138:181–192. [PubMed: 9214391]
47. Fanning AS, Ma TY, Anderson JM. Isolation and functional characterization of the actin binding region in the tight junction protein ZO-1. *Faseb J* 2002;16:1835–1837. [PubMed: 12354695]
- 48. Sakurai A, Fukuhara S, Yamagishi A, Sako K, Kamioka Y, Masuda M, Nakaoka Y, Mochizuki N. MAGI-1 Is Required for Rap1 Activation upon Cell-Cell Contact and For Enhancement of Vascular Endothelial Cadherin-mediated Cell Adhesion. *Mol Biol Cell* 2005;17:966–976. [PubMed: 16030252]

16339077]This work shows that in cultured vascular endothelial cells, MAGI-1, which can bind to both β -catenin and the Rap1 guanine nucleotide exchange factor, PDZ-GEF/Rap-GEF, is important for junction-localized Rap1 activation, based on in vivo FRET activity assays.

49. Hogan C, Serpente N, Cogram P, Hosking CR, Bialucha CU, Feller SM, Braga VM, Birchmeier W, Fujita Y. Rap1 regulates the formation of E-cadherin-based cell-cell contacts. *Mol Cell Biol* 2004;24:6690–6700. [PubMed: 15254236]
50. Shin K, Fogg VC, Margolis B. Tight junctions and cell polarity. *Annu Rev Cell Dev Biol* 2006;22:207–235. [PubMed: 16771626]
51. Tepass U. Crumbs, a component of the apical membrane, is required for zonula adherens formation in primary epithelia of *Drosophila*. *Dev Biol* 1996;177:217–225. [PubMed: 8660889]
52. Grawe F, Wodarz A, Lee BS, Knust E, Skaer H. The *Drosophila* genes crumbs and stardust are involved in the biogenesis of adherens junctions. *Development* 1996;122:951–959. [PubMed: 8631272]
53. Rohr S, Bit-Avragim N, Abdelilah-Seyfried S. Heart and soul/PRKCi and nagie oko/Mpp5 regulate myocardial coherence and remodeling during cardiac morphogenesis. *Development* 2006;133:107–115. [PubMed: 16319113]
- 54. Wang Q, Chen X-W, Margolis B. PALS1 regulates E-cadherin trafficking in mammalian epithelial cells. *Mol Biol Cell* 2007;18:874–885. [PubMed: 17182851]This paper examines effects of knockdown of the vertebrate Stardust homolog, Protein Associated with Lin Seven 1 (PALS1), and finds that E-cadherin trafficking to the cell surface is affected. Exocyst complex componentns are mislocalized in the knockdown cells, which could account for E-cadherin trafficking defects.
55. Verma S, Shewan AM, Scott JA, Helwani FM, Elzen NR, Miki H, Takenawa T, Yap AS. Arp2/3 Activity Is Necessary for Efficient Formation of E-cadherin Adhesive Contacts. *J Biol Chem* 2004;279:34062–34070. [PubMed: 15159390]

Figure 1. The trafficking itineraries of classical cadherins

Newly synthesized cadherins are trafficked from the Golgi to the plasma membrane. Upon delivery to the surface, cadherins may engage in productive adhesive interactions (1) or be endocytosed (2). Following internalisation, cadherins are transported to early or sorting endosomes. Two alternative routes are available from this compartment: recycling back to the cell surface (3), a path may commonly involve an intermediate step in the recycling endosomal compartment; or transport through late endosomes for degradation in the lysosome (4). Which of these alternatives is taken can influence surface expression and metabolic stability of the cadherin.