Published in final edited form as:

Biologicals. 2007 October ; 35(4): 367–371.

A summary of the workshop on Passive Immunization Using Monoclonal Antibodies for HIV/AIDS:

Held at the National Institute of Allergy and Infectious Diseases, Bethesda on March 10, 2006

Geetha P. Bansal

Division of AIDS, National Institute of Allergy and Infectious Diseases, NIH, Bethesda, Maryland

Workshop goals

Passive immunization with antibodies has been shown to prevent or treat a wide variety of infectious diseases. In the early use of this technology, antibodies were administered as human or animal plasma. Improvements in technology led to the process of producing the antibodies as a purified immunoglobulin fraction from pooled human plasma for intravenous (IVIG) or intramuscular (IG) use. These immunoglobulin fractions contained specificities against multiple antigens and thus only a fraction of the specificities was directed against any one disease agent. The exquisite specificity of monoclonal antibodies (MAbs) has made it possible to develop antibody products with increased potency and ease of administration. Although several MAbs for infectious diseases are in clinical trials, only one product (against respiratory syncytial virus) is licensed for use. Currently, no MAb products have been licensed for use for therapy or prophylaxis against HIV infection. However, rational arguments can be advanced for the use of passive immunization to prevent HIV infection, especially to reduce mother-to-child transmission (MTCT) of HIV, and in immediate post exposure situations. The key arguments questioning the feasibility of this approach relate to the alternate availability and use of effective anti-retroviral drugs, and to the high cost of production and distribution of MAb products.

A one-day workshop was held in March 2006 to evaluate the potential role of passive immunization using MAbs for therapeutic or prophylactic outcomes in HIV/AIDS, and to evaluate the feasibility and acceptability of using MAbs in adults, children and infants. An additional goal was to discuss whether there were roles for anti-HIV MAbs and passive immunization studies other than in clinical settings, and if the field needed to pursue them. This workshop had representation from the perspectives of basic science and clinical research, production and manufacturing, cost consideration and regulatory issues, among others. The expected output was a better understanding of the importance of MAbs for passive immunization against HIV and to evaluate the possibility of identifying protective epitopes, designing appropriate protection models and clarifying regulatory requirements.

At a previous workshop on Immunoprophylaxis for HIV-1 in Pediatrics: Moving Concepts to Reality on Vaccines and Passive Immunity held in 2002, the discussion focused on neonatal immunoprophylaxis strategies and how to incorporate our understanding of the pathogenesis of HIV infection in MTCT into the design of immunoprophylaxis protocols, a review of antibodies and vaccine candidates currently or imminently available, available cohorts of

Address Correspondence to Geetha P. Bansal, Ph. D., Division of AIDS, National Institute of Allergy and Infectious Diseases, 6700-B Rockledge Drive, Room 5120, Bethesda, MD 20892, Phone 301-496-5042, Fax 301 - 402- 3684, e-mail: gbansal@niaid.nih.gov.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

subjects for clinical trials and potential barriers to implementation of neonatal immunoprophylaxis trials. The workshop reviewed both active and passive immunization regimens, as well as ideas for a combination approach [1].

As a result of the 2002 workshop, a few anti-HIV MABs were pursued and a few small studies of safety and effectiveness have been conducted in adults. At that time, a limited number of well characterized human anti-HIV MABs were available that exhibited potent and broad neutralizing activity. MABs b12, F105 and 2G12 bind to the surface of gp120 envelope glycoprotein. MABs 2F5 and 4E10 recognized linear epitopes near the membrane-spanning region of gp41 [cross referenced in 1]. The currently described March 2006 workshop set out to address a limited number of questions as listed in Table 1. The outcome from the discussions is summarized in this report.

Passive immunization studies

Because of the exquisite specificity of antigen recognition by MABs, the ability to characterize them more reproducibly, to generate quantitative and qualitatively defined amounts consistently and for safety reasons (as compared to harvesting from polyclonal plasma donations), the preference for MAB products for passive protection in clinical settings is clear. MABs can be used to provide evidence for *in vivo* protective effect, for identification of vaccine epitopes as a preventive strategy in newborns or for post-exposure prophylaxis.

In some countries, women of childbearing age are particularly at high risk of HIV infection and primary prevention strategies are desperately needed. While microbicides and vaccines offer the best hope for protecting women against HIV infection, none are currently available. Simian models have been very helpful in understanding the mechanisms and early events in vaginal SIV transmission, and topical or systemic administration of MABs has provided proof of the importance of neutralizing antibodies in protection from infection (2, 12). These studies provide a strong rationale for developing preventive vaccines that elicit strong neutralizing antibodies or the development of microbicides that incorporate MABs to protect women.

World wide, MTCT of HIV is the primary mode of pediatric infection. Antiretroviral therapy (ART) can significantly prevent MTCT in the peripartum period; however, the utility of these regimens may be short-lived due to the emergence of ART resistance. In addition, perinatal ART does not prevent MTCT through breast-feeding, which still continues to be a major mode of transmission. The need for preventing MTCT during the breast feeding period needs special attention especially in financially constrained societies where for several reasons, such as lack of affordable and acceptable options, breast-feeding cannot be discouraged or discontinued [3]. In addition, infants born to HIV+ women do not have a fully developed immune system and may well benefit from passive immunization. More work needs to be done to understand the biology of developing immune responses to HIV in children <6 months of age, such as understanding the neutralization sensitivity of vertically or horizontally transmitted viruses [4].

Studies in juvenile macaques showing that passively introduced neutralizing antibodies resulted in enhanced *de novo* production of neutralizing antibodies in infected animals [5] provide another reason for exploring the value of passive immunization with MABs in a passive-active immunization strategy.

Passive immunization and protection studies in non-human primates (NHP) using the broadly neutralizing MABs 2F5, 2G12, F105 and b12 indicate that a cocktail of 2 or more MABs is more potent than single MABs, and that passively transferred antibody administered prior to or within a few hours after challenge is more effective at inducing sterilizing immunity [6,7].

It is possible that the administration of neutralizing MAbs at the time of birth, and monthly during the neonatal period, in combination with ART, may provide sterilizing protection.

Strategies for developing useful MAbs for HIV/AIDS

Very few MAbs have been used to study vaccine epitopes, and most of these have been generated against Clade B HIV. Much of the information required for vaccine design or for epitope definition will necessarily need to be generated via induction of neutralizing MAbs and passive protection studies. There is a need for generating MAbs to other Clades in order to define protective epitopes of HIV-1 isolates in endemic regions. The field needs more standardized and cohesive virus panels for screening MAbs; these would also be useful for testing vaccine responses.

The availability of consensus Clades B and C virus panels will be useful to screen for broadly neutralizing MAbs [8,9]. It was also suggested that in view of the differences in the neutralization epitopes between transmitted and parent viruses [10], it may be a good idea to make a consensus panel of newly transmitted viruses, including those passed from mothers to infants during delivery or post-partum through breast milk. Much work is needed in understanding the nature of the transmitted virus. There are a number of questions that need to be addressed with respect to the critical epitopes on the virus that are associated with transmission and which can be targets of neutralizing MAbs. However, it is also unclear if it will be possible to find a conserved epitope that will be present in all transmitted viruses, such as between different mother-infant pairs or hetero-sexual donor-recipient pairs. To add another level of complexity, it is unclear as to whether critical epitopes involved in virus transmission are different as virus is transmitted across different tissues, and whether transmission occurs by cell-associated virus or cell-free virus. Thus, for all these reasons, from the discussions in the context of the utility of compiling a newly transmitted virus panel, it was clear that because there was not sufficient information available regarding the epitopes seen in different transmission settings, whether the search for breadth of reactivity of neutralizing MAbs against HIV would now be further stretched to include not just regional Clades but different transmitted isolates.

Safety of *in vivo* administered MAbs

While the *in vivo* studies in NHP models and Phase I studies in humans have been encouraging for contemplating the use of these broadly reacting anti-HIV MAbs in clinical settings (especially for breast feeding infants), the chance discovery that these antibodies exhibit cross-reactive binding to self antigens has begun a whole new process of inquiry into the safety issues associated with the *in vivo* use of MAbs [11]. MAb 4E10 has not been studied by itself in NHP, and there is limited data from three juvenile macaques studied with 2F5 alone in NHP [12]. In light of the “anti-phospholipid” reactivity of these antibodies *in vitro*, a study to assess the long-term *in vivo* adverse effects of administering such antibodies in NHP may be worthwhile and could provide valuable information.

With respect to the adverse events noted in the NHP studies [7] it seems that the one infant rhesus (out of 45 studied), which had neurological symptoms including hind limb paralysis, may have been predisposed to possible intracranial bleeding. Observations such as this are disturbing and could potentially be an obstruction for further development; however when such incidents are rare occurrences, a more careful and perceptive analysis needs to be done before discontinuing such approaches. Regulatory agencies such as the FDA can help or guide in interpreting such data, to enable sound decisions regarding pros and cons of pursuing potential candidates.

The polyspecificity and/or autoantigen reactivity of some of the current MABs (such as 4E10 and 2F5) need to be critically looked at. There does not appear to be a consensus that the anti-cardiolipin and other autoantigen reactivities of the currently studied MABs are associated with pathogenicity [13,17]. Anti-phospholipid antibodies have been reported in infectious disorders, such as AIDS and syphilis, without the clinical features of anti-phospholipid syndrome. Despite the prolonged clotting times, bleeding is not a typical feature associated with administration of these antibodies. Thrombocytopenia, if present, is usually mild. Patients may have either a lupus anticoagulant or an anticardiolipin antibody or they may have both antibodies. A high percentage of patients with systemic lupus erythematosus (SLE) or related autoimmune diseases have these antibodies, which may also develop in patients without an underlying disorder and transiently in association with certain medications (e.g., hydralazine, phenytoin) or infections. The human immunodeficiency virus (HIV) is commonly associated with positive tests for anti-phospholipid antibodies [14]. Infection-associated antibodies may not be associated with clinical symptoms of anti-phospholipid antibody syndrome, and they tend to recognize phospholipid rather than the phospholipid-protein complexes described [15]. In fact, it is entirely possible that broadly cross-reactive potent neutralizing MABs will intrinsically be poly-reactive, in which case the biological and biochemical properties of these antibodies need to be better studied. As a start to collecting such information, Partial Thromboplastin Time (PTT, a measure of blood clotting time) measurements were made on 4 human subjects being treated with a cocktail of 2G12, 4E10 and 2F5. It was observed that PTT was prolonged; however it was considered to be mild (symptoms causing no or minimal discomfort). One patient showed a low level of lupus anticoagulant activity. No other overt/significant clinical adverse event was reported [16].

Therapeutic considerations for *in vivo* administration of MABs

In adult therapeutic trials, resistance to MABs has been shown to develop *in vivo* in HIV infected patients who discontinue ART. In studies performed to assess the *in vivo* efficacy of passively administered pool of 2F5, 4E10 and 2G12, viral rebound upon cessation of antibody infusion was noted [17]. In general the antibody infusions were well tolerated and there were no significant adverse effects. The conclusion from clinical trials with anti-env MABs was that resistance to 2G12 developed early, and also that 2G12 had a longer elimination half life than 2F5 and 4E10. In general viral fitness was significantly decreased in the presence of 2F5 and 4E10 [18]. Passive immunization using broadly neutralizing antibodies as an adjunct to HAART needs further evaluation, using combinations of antibodies with a more favorable half life or tissue distribution profile, as well as doses that would provide neutralization responses above that achieved with autologous plasma.

An anti-CD4 MAB, TNX-355 (Tanox, Inc), is currently in Phase II trials and shows some promise of efficacy; however, resistant viruses using CD4 are still generated *in vivo*. Further work with these resistant viruses is ongoing to understand the underlying mechanisms of resistance development. MAB dB4 (United Biomedical, Inc) is another anti-CD4 MAB that has shown encouraging results in NHP challenge studies and is currently being contemplated for clinical trials. Anti-cell surface antibodies may have promise as therapeutic MABs; there are plans to use them in combinations to improve neutralizing responses. It was suggested that antibodies targeting CCR5 may have clinical utility.

Future considerations for research and development of anti-HIV MABs

Immunoglobulin class (heavy chain) is an important factor to consider in that virus neutralization can be brought about by functions such as antibody dependent cellular cytotoxicity (ADCC) and cell mediated lysis (CML) in an *in vivo* setting. It was noted that 2F5 and 4E10 were initially isolated as IgG3 isotype and it was also shown that F105, when isotype-

switched from an IgG1 to an IgG3, showed enhanced neutralizing activity. Because of the abundance of non-neutralizing antibodies present in HIV-positive individuals, studies were encouraged for investigating whether such non-neutralizing antibodies may exhibit synergistic action in mixtures with other less studied non-neutralizing or neutralizing MABs.

Although a long way off from a potential product stage, the possible use of antibodies as microbicidal agents was discussed. *In vitro* experiments looking at antibody release from vaginal rings are very promising. The use of vectors such as lactobacilli to express antibodies may be worth contemplating since they (antibodies and lactobacilli) may provide some benefits with respect to safety or decreased toxicity compared to some of the current formulations under study, and can be contemplated for counteracting perinatal transmission of some infections.

Finally, the development of a MAB based product should include an early recognition of marketability, production logistics and cost related issues, as well as regulatory and safety considerations. Discussion with the FDA (www.fda.gov/cder/ode4/preind/default.htm) and/or industry would be worthwhile if begun early at the time product oriented research is contemplated.

Table 2 lists some of the gaps and action items that were identified for consideration while revisiting issues related to the use of MABs in passive immunization studies for HIV/AIDS by the panel members who participated in the workshop, as shown in Table 3. From this brief summary of the workshop presentations and discussions, it appears that priorities should include more work on identifying the nature of the transmitted viral epitopes, isolating neutralizing MABs with breadth of reactivity, investigating the autoimmune reactivity and its relevance, and in developing consensus assays for the screening of relevant and useful MABs.

Key References

1. Safrin J, et al. Immunoprophylaxis to Prevent Mother to Child Transmission of HIV-1. *J Acquir Immune Defic Syndr* 2004 February;35(2):169–177. [PubMed: 14722451]
2. Veazey RS, Shattock RJ, Pope M, Kirijan JC, Jones J, Hu Q, Ketas T, Marx PA, Klasse PJ, Burton DR, Moore JP. Prevention of virus transmission to macaque monkeys by a vaginally applied monoclonal antibody to HIV-1 gp120. *Nat Med* 2003 Mar;9(3):343–6. [PubMed: 12579198]
3. John-Stewart G, Mbori-Ngacha D, Ekpin R, Janoff EN, Nkengasong J, Read JS, Van de Perre P, Newell ML. Ghent IAS Working Group on HIV in Women Children. Breast-feeding and Transmission of HIV-1. *J Acquir Immune Defic Syndr* 2004 February;35(2):196–202. [PubMed: 14722454]
4. Luzuriaga K, Newell ML, Dabis F, Excler JL, Sullivan JL. Vaccines to prevent transmission of HIV-1 via breastmilk: scientific and logistical priorities. *Lancet* 2006;368:511–521. [PubMed: 16890838]
5. Haigwood NL, Montefiori DC, Sutton WF, McClure J, Watson AJ, Voss G, Hirsch VM, Richardson BA, Letvin NL, Hu SL, Johnson PR. Passive immunotherapy in simian immunodeficiency virus-infected macaques accelerates the development of neutralizing antibodies. *J Virol* 2004 Jun;78(11):5983–95. [PubMed: 15140996]
6. Ferrantelli F, Rasmussen RA, Buckley KA, Li P-L, Wang T, Montefiori DC, Katinger H, Stiegler G, Anderson DC, McClure HM, Ruprecht RM. Complete protection of neonatal rhesus macaques against oral challenge with pathogenic SHIV by human anti-HIV monoclonal antibodies. *J Infect Dis* 2004;189:2167–73. [PubMed: 15181562]
7. Hofmann-Lehmann R, Vlasak J, Rasmussen RA, Smith BA, Baba TW, Liska V, Ferrantelli F, Montefiori DC, McClure HM, Anderson DC, Bernacky BJ, Rizvi TA, Schmidt R, Hill LR, Keeling ME, Katinger H, Stiegler G, Cavacini LA, Posner MR, Chou T-C, Andersen J, Ruprecht RM. Postnatal passive immunization of neonatal macaques with a triple combination of human monoclonal antibodies against oral simian-human immunodeficiency virus challenge. *J Virol* 2001;75:7470–80. [PubMed: 11462019]
8. Li M, Gao F, Mascola JR, Stamatatos L, Polonis VR, Koutsoukos M, Voss G, Goepfert P, Gilbert P, Greene KM, Bilska M, Kothe DL, Salazar-Gonzalez JF, Wei X, Decker JM, Hahn BH, Montefiori

- DC. Human immunodeficiency virus type 1 env clones from acute and early subtype B infections for standardized assessments of vaccine-elicited neutralizing antibodies. *J Virol* 2005 Aug;79(16):10108–25. [PubMed: 16051804]
9. Li M, Salazar-Gonzalez JF, Derdeyn CA, Morris L, Williamson C, Robinson JE, Decker JM, Li Y, Salazar MG, Polonis VR, Mlisana K, Karim SA, Hong K, Greene KM, Bilska M, Zhou J, Allen S, Chomba E, Mulenga J, Vwalika C, Gao F, Zhang M, Korber BT, Hunter E, Hahn BH, Montefiori DC. Genetic and neutralization properties of subtype C human immunodeficiency virus type 1 molecular env clones from acute and early heterosexually acquired infections in Southern Africa. *J Virol* 2006 Dec;80(23):11776–90. [PubMed: 16971434]Epub 2006 Sep 13
 10. Wu X, Parast AB, Richardson BA, Nduati R, John-Stewart G, Mbori-Ngacha D, Rainwater SM, Overbaugh J. Neutralization escape variants of human immunodeficiency virus type 1 are transmitted from mother to infant. *J Virol* 2006 Jan;80(2):835–44. [PubMed: 16378985]Erratum in: *J Virol*. 2006 Mar; 80(5):2585
 11. Lam GK, Liao HX, Xue Y, Alam SM, Searce RM, Kaufman RE, Sempowski GD, Haynes BF. Cardiolipin polyspecific autoreactivity in two broadly neutralizing HIV-1 antibodies. *Science* 2005 Jun 24;308(5730):1906–8. [PubMed: 15860590]Epub 2005 Apr 28
 12. Mascola JR, Lewis MG, Stiegler G, Harris D, VanCott TC, Hayes D, Louder MK, Brown CR, Sapan CV, Frankel SS, Lu Y, Robb ML, Katinger H, Birx DL. Protection of Macaques against pathogenic simian/human immunodeficiency virus 89.6PD by passive transfer of neutralizing antibodies. *J Virol* 1999 May;73(5):4009–18. [PubMed: 10196297]
 13. Sherer, EM.; Zwick, MB.; Teyton, L.; Burton, DR. The broadly neutralizing antibodies 2F5 and 4E10: specific anti-HIV antibodies or autoantibodies?. Fifth International Conference: AIDS Vaccine 2006 Abstract #P09A-26 (Antiviral Therapy); 2006. p. 134AIDS 2007 in press
 14. de Larranaga GF, Forastiero RR, Carreras LO, et al. Different Types of Anti-phospholipid Antibodies in AIDS: A Comparison with Syphilis and the Anti-phospholipid Syndrome. *Thromb Res* 1999;96(1):19–25. [PubMed: 10554081]
 15. McNally T, Purdy G, Mackie IJ, et al. The Use of an Antibeta-2-Glycoprotein-I Assay for Discrimination Between Anticardiolipin Antibodies Associated With Infection and Increased Risk of Thrombosis. *Br J Haematol* 1995;91(2):471–3. [PubMed: 8547096]
 16. Saurabh, Mehendru; Vcelar, B.; O'Neil, M.; Wrin, T.; Stiegler, G.; Shet, A.; Galovich, J.; Petropoulos, C.; Katinger, H.; Markowitz, M. Monoclonal Antibody Infusions Delay HIV-1 Rebound after Discontinuation of ART in a Cohort of HIV-1-infected Individuals Treated during Primary HIV-1 Infection. 13th Conf. on Retroviruses and Opportunistic Infections, 2006. Session 43 Abstract #178;
 17. Manrique A, Rusert P, Joos B, Fischer M, Kuster H, Leemann C, Niederöst B, Weber R, Stiegler G, Katinger H, Günthard HF, Trkola A. In vivo and in vitro escape from neutralizing antibodies 2G12, 2F5 and 4E10. *J Virol*. 2007 June 13;Epub ahead of print
 18. Joos B, Trkola A, Kuster H, Aceto L, Fischer M, Stiegler G, Armbruster C, Vcelar B, Katinger H, Günthard HF. Long-term multiple-dose pharmacokinetics of human monoclonal antibodies (MAbs) against human immunodeficiency virus type 1 envelope gp120 (MAb 2G12) and gp41 (MAbs 4E10 and 2F5). *Antimicrob Agents Chemother* 2006 May;50(5):1773–9. [PubMed: 16641449]

Table 1**Points to consider for discussion at the March 2006 workshop****Scientific:**

- a) Target population (HIV infected subjects; infants born to HIV infected mothers; breast feeding infants etc)
- b) Target epitopes for the Mabs?
- c) Single MAb or synergistic combination of MAb? How many in the combination?
- d) Synergy between different MAb combinations or synergy between drugs and MAb?
- e) Polyreactivity of antibodies and other properties that may become issues?
- f) Other emerging scientific issues?

Technical

- a) Feasibility of approach (Rationale for using MAb; therapeutic or prophylactic)
- b) Production/manufacturing and costs; Distribution; ease of administration
- c) Drugs vs MAb (Will availability of effective drugs compromise consideration of MAb)
- d) Other technical/logistical issues?

Clinical Trials

- a) Dosage in infants and children
- b) Mode of administration
- c) Formulation
- d) Adverse effects (known or unknown)
- e) Regulatory issues
- f) Other?

Table 2**Action items from the March 2006 workshop:**

1. Better understanding of the mechanism/early viral replication following mucosal transmission (including MTCT).
2. Better understanding of the genetics and biological properties of transmitted viruses (including susceptibility to entry inhibitors or Mab).
3. Better understanding of how to elicit highly neutralizing Ab using vaccination.
4. Better understanding of the clinical relevance of anti-phospholipid Abs.
5. Better understanding of the bases and clinical significance of differences in readouts from different neutralizing Ab studies.
6. Continue to develop high-throughput assays that can speed basic and clinical research; including the development of transmitted virus panels and standardized biologically relevant assay methods
7. Continue to drive clinical evaluation of promising, available products, particularly in infants and children. There is a need for generating MAbs to other Clades to define protective epitopes in those endemic regions
8. Further study synergy between non-neutralizing antibodies in mixtures with either non-neutralizing or neutralizing antibodies as well as cell-surface directed antibodies and anti-HIV antibodies.
9. Investigate the role of antibodies as microbicidal agents, and microbicides in general (with specific focus on antibody-based microbicides) as an intervention for perinatal transmission.
10. Further evaluate passive immunization using broadly neutralizing antibodies as an adjunct to HAART using combinations of antibodies with a more favorable Pk profile, as well as doses that would provide neutralization responses more than what can be achieved with autologous plasma.

Table 3**Participating Panel Members at the March 2006 workshop**

Dr. Cavacini, Lisa	Beth Israel Deaconess Medical Center
Dr. Finstad, Connie	United Biomedical Inc.
Dr. Fung, Michael	Tanox, Inc.
Dr. Hanson, Carl	California Department of Health Services
Dr. Haynes, Barton	Duke University
Dr. John-Stewart, Grace	University of Washington
Dr. Katinger, Hermann	Polymun Scientific, Austria
Dr. Koenig, Scott	MacroGenics, Inc.
Dr. Lewis, Stanley	Tanox, Inc.
Dr. Lockshin, Michael	Cornell University
Dr. Luzuriaga, Katherine	University of Massachusetts. Medical School.
Dr. Markowitz, Martin	Aaron Diamond AIDS Research Center
Dr. Markham, Richard	Johns Hopkins University
Dr. Mascola, John	Vaccine Research Center/NIH
Dr. Meade, Harry	GTC Biotherapeutics
Dr. Mynahan, Richard	Leerink Swann/MEDA Corp
Dr. Ruprecht, Ruth	Dana-Farber Cancer Institute
Dr. Safrit, Jeffrey	Elizabeth Glaser Pediatric AIDS Foundation
Dr. Trkola, Alexandra	University Hospital, Zurich