Serological comparison and haemagglutinating activity of *Mycoplasma dispar*

By C. J. HOWARD, R. N. GOURLAY AND JACQUELINE COLLINS

*Agricultural Research Council, Institute for Research on Animal Diseases,
Compton, Newbury, Berks. RG16 0NN*

(Received 3 June 1974)

SUMMARY

A comparison of twelve strains of *Mycoplasma dispar* by the metabolism inhibition and indirect haemagglutination tests has shown them to form a serologically homogeneous group of micro-organisms. The twelve strains vary in their haemagglutinating activity against erythrocytes from different animal species, and certain of the strains can be distinguished by the erythrocytes they agglutinate. Haemagglutination may thus provide a method by which certain strains can be typed. The erythrocyte receptor site does not appear to contain sialic acid and is not sensitive to proteolytic enzymes. On the mycoplasma cell two attachment sites have been demonstrated. One, by which it attaches to sheep and bovine erythrocytes, is a protein or contains a protein moiety. The chemical nature of the other attachment site, by which *M. dispar* attaches to rabbit erythrocytes, is unknown.

INTRODUCTION

Mycoplasma dispar was first isolated from pneumonic calf lungs (Gourlay, 1969; Gourlay & Leach, 1970) and subsequently from the respiratory tract of non-pneumonic calves (Thomas & Smith, 1972). Its role in calf pneumonia remains to be determined. However, the preliminary experiments of Gourlay & Thomas (1969) and St George, Horsfall & Sullivan (1973), who found lesions in the lungs of calves after endobronchial or intratracheal inoculation of broth cultures but not sterile broth, together with the observation that, of the mycoplasmas commonly found in the respiratory tract of calves in England, *M. dispar*, *M. bovirhinis*, *Acholeplasma laidlawii* and T-mycoplasma (Gourlay, Mackenzie & Cooper, 1970), only *M. dispar* had any cytopathic effect on bovine fetal tracheal organ cultures (Thomas & Howard, 1974) and the finding that certain strains of *M. dispar* cause clinical mastitis in cows experimentally inoculated (J. Brownlie, R. N. Gourlay & C. J. Howard – to be published) supports the hypothesis that this micro-organism is pathogenic for cattle, perhaps with a particular propensity for the respiratory tract.

Comparisons within certain mycoplasma species, e.g. *M. hominis* (Purcell *et al.* 1967) *M. gallisepticum* (Taylor-Robinson & Berry, 1969) and *M. pulmonis* (Forshaw & Fallon, 1972) have shown these species to be serologically heterogeneous and to contain various subtypes. No serological comparison of a group of *M. dispar*

strains has been made except for that of Gourlay & Leach (1970) who found that antiserum to strain 462/2 inhibited the growth on agar of 23 out of 36 strains examined. As little information is available on the serological heterogeneity or homogeneity of the species the first part of this study deals with this aspect of *M. dispar*.

Attachment to mucosal epithelium is considered to be the first step in the production of disease by many micro-organisms (Savage, 1972). The attachment of mycoplasmas to cells is well documented and differences between strains in their ability to attach to cells has been observed (Manchee & Taylor-Robinson, 1969*a, b*). The intimate association between mycoplasmas and cells has been proposed as a mechanism by which high local concentrations of toxic metabolites e.g. H_2O_2 , are built up which may then cause cell damage (Sobeslavsky, Prescott & Chanock, 1968). In the second part of this paper the demonstration of attachment sites on *M. dispar* for erythrocytes is reported, and these attachment sites may represent the mechanism by which this micro-organism maintains itself in the bovine respiratory tract. Besides this, a study of haemagglutinating activity of *M. dispar* provides information on its surface structure and also forms the basis of a method useful for strain identification.

METHODS

Media

Mycoplasmas were grown in glucose calf-serum (GS) broth (Gourlay & Leach, 1970) containing ampicillin (Andrews, Leach, Gourlay & Howard, 1973) except when organisms were grown for injection into rabbits, in which case rabbit digest broth and rabbit serum, heated at 56° C. for 30 min. were substituted for Hartley's digest broth and fetal calf serum.

M. dispar strains

All strains were from the lungs of calves with pneumonia. In order that we should not select serologically similar strains the criteria used to identify organisms as *M. dispar*, in the first instance, were production of acid from glucose in GS broth and colony morphology. *M. dispar* forms atypical centreless colonies on primary isolation (Gourlay & Leach, 1970). The serological studies confirmed the identity of the strains used subsequently for haemagglutination. Antisera were prepared as described by Howard & Gourlay (1972). All strains were cloned by filtering broth cultures through 650 nm. Millipore filters, and propagation of single colonies on three successive occasions (Sub-committee on the Taxonomy of *Mycoplasmatales*, 1972).

Influenza A virus

Strain SIWG/1/57 was kindly provided by Dr A. S. Beare (Common Cold Unit, Salisbury) and stored at -70° C.

Serological tests

The metabolism inhibition test was performed according to Taylor-Robinson, Purcell, Wong & Chanock (1966). The indirect haemagglutination test was performed according to Herbert (1967) using a microtitre system (Cooke Engineering Co., Flow Laboratories, Irvine, Scotland), sonicated *M. dispar* as antigen and tanned sheep erythrocytes that had been stored in Alsevers solution (Wellcome Laboratories, Beckenham, Kent).

Haemagglutination

The various strains of *M. dispar* used were grown in GS broth at 37° C., harvested by centrifugation at 15,000 g, washed and resuspended in 0.15 M-NaCl and stored at -70° C. The optical density (OD) of mycoplasma suspensions at 500 nm. was measured with a 10 mm. light path cell in a Unicam SP 800 and haemagglutination titres are given for standard suspensions with an OD of 10.

Rabbit, rat, hamster, chicken, calf and guinea-pig red cells were collected and stored in Alsevers solution. Sheep and horse red cells in Alsevers solution were obtained from the Wellcome Laboratories and human group O citrated packed cells from the Regional Transfusion Centre, Oxford. All red cells were stored at 4° C. Before use they were washed three times with 0.15 M-NaCl and made up to the required concentration by volume.

Doubling dilutions of antigen were made in 25 µl. volumes of 0.15 M-NaCl using a microtitre system and U trays (Cooke Engineering Co.) and the red cells, 25 µl. of a 1 % v/v suspension, were added to the dilutions of antigens. The results were read after overnight incubation at 4° C. All haemagglutination titres are given as the reciprocal of the highest dilution of mycoplasma suspension that caused agglutination.

Treatment applied to erythrocytes

The methods followed were mainly those of Buckland & Tyrrell (1963). Receptor-destroying enzyme (Wellcome Laboratories, from *Vibrio cholerae*) diluted 1/4 in 0.15 M-NaCl was mixed with an equal volume of 5 % v/v erythrocytes. Equal volumes of the other reagents and 10 % v/v erythrocytes were incubated together. Trypsin (bovine pancreas type III) and chymotrypsin (bovine pancreas type I) were obtained from Sigma (Kingston upon Thames). Pronase (B.D.H., Poole) and erythrocytes were incubated together for 30 min. at 37° C. After the various treatments the cells were washed once with 0.15 M-NaCl, resuspended to give a 1 % suspension, and 25 µl. volumes were added to the doubling dilutions of the mycoplasma suspension.

Treatment applied to mycoplasmas

Equal volumes of mycoplasma suspension and the reagents listed in Table 5 were incubated together. The temperatures and times of incubation were as used by Buckland & Tyrrell (1963) for treating viruses. Pronase and mycoplasmas were incubated together at 37° C. for 30 min. After incubation the mycoplasmas were centrifuged and made up to their original volume.

Table 1. *Metabolism inhibition titres* of antisera to M. dispar strains*

Strain tested against:	Antiserum prepared against:										
	F370	462/2	Vic12	Vic13	Gri 226	D2	D44	Vic7	Gri 250	Mmb 177	D52
F370	2560	1280	2560	640	2560	1280	2560	2560	1280	1280	1280
462/2	1280	640	640	640	160	80	160	40	80	160	320
Vic12	1280	640	1280	320	640	320	640	160	320	320	640
Vic13	1280	2560	1280	640	1280	640	1280	1280	640	640	640
Gri226	1280	640	1280	640	320	320	320	320	320	320	160
D2	2560	640	1280	640	1280	1280	2560	1280	1280	640	1280
D44	1280	1280	1280	1280	1280	1280	1280	1280	1280	640	640
Vic7	2560	1280	1280	1280	1280	1280	1280	1280	640	640	640
Gri250	1280	640	640	320	640	160	160	320	160	320	320
Mmb177	1280	1280	1280	640	640	1280	320	640	640	640	640
D52	2560	1280	1280	1280	1280	1280	1280	640	640	640	1280
Gri221	640	1280	640	160	320	160	80	160	160	160	160

* Reciprocal of highest dilution of antiserum inhibiting the production of acid from glucose.

Electron microscopy

M. dispar and erythrocytes were examined by electronmicroscopy after staining with ruthenium red (Springer & Roth, 1973) as previously described (Howard & Gourlay, 1974).

RESULTS

Serological comparison of twelve M. dispar strains

The results of the metabolism inhibition and indirect haemagglutination tests (Tables, 1, 2) confirmed that the strains, identified primarily on the basis of colony morphology and ability to produce acid from glucose, were *M. dispar*. The twelve strains studied form a serologically homogeneous group, only minor differences in antigenicity being indicated by the two tests. It was considered that these differences might be largely due to the variation in potencies of the different antisera, the state of growth of antigen in the metabolism inhibition test and the antigen concentration used to sensitize the red cells in the indirect haemagglutination test. Normal rabbit serum had titres of < 20 in the metabolism inhibition and indirect haemagglutination tests. No factor was observed analogous to that in rabbit sera which prevents the growth of human T-mycoplasmas (Howard & Gourlay, 1973).

Antisera were raised against organisms grown in medium containing rabbit digest broth and serum. However, in order to ensure that antibodies to bovine serum were not being detected in the indirect haemagglutination test, the dilutions of antisera and the final suspensions of tanned-sensitized sheep red cells were made in phosphate buffered saline containing 5% v/v fetal calf serum. Besides this, absorption of antiserum with bovine erythrocytes had no effect on the titre of antisera for sensitized red cells.

Table 2. *Titres* in indirect haemagglutination test of antisera to M. dispar strains*

Strain used as antigen	Antiserum prepared against:										
	F370	462/2	Vic12	Vic13	Gri226	D2	D44	Vic7	Gri250	Mmb177	D52
F370	5120	5120	5120	5120	1280	2560	1280	640	640	640	320
462/2	10240	5120	320	1280	1280	320	640	80	160	640	320
Vic12	5120	2560	10240	20480	2560	2560	2560	1280	2560	640	1280
Vic13	10240	5120	2560	5120	5120	2560	640	640	320	2560	640
Gri226	20480	10240	5120	10240	10240	2560	2560	2560	1280	2560	1280
D2	20480	20480	5120	20480	5120	20480	10240	10240	5120	2560	1280
D44	10240	2560	1280	5120	1280	2560	1280	640	160	1280	320
Vic7	10240	5120	5120	10240	2560	5120	2560	2560	2560	1280	640
Gri250	40960	20480	10240	20480	10240	10240	10240	2560	2560	1280	1280
Mmb177	40960	40960	20480	40960	10240	10240	10240	10240	5120	5120	2560
D52	20480	5120	10240	20480	20480	5120	10240	10240	2560	2560	2560
Gri221	20480	20480	20480	20480	10240	5120	5120	10240	2560	1280	640

* Reciprocal of highest dilution of antiserum causing agglutination of sensitized erythrocytes.

Haemagglutination by M. dispar

Strains 462/2, Gri226, Gri221, F370, Mmb177 and Vic13 were tested for haemagglutinating activity against sheep, rat, rabbit, human, guinea-pig, hamster, chicken, horse and calf erythrocytes at 4°, 22°, and 37° C. There was little variation in agglutination titres at the different temperatures. However, the more concentrated mycoplasma suspensions caused haemolysis at 37° and 22° C. and subsequent tests were therefore performed at 4° C.

The growth of *M. dispar* strains F370 and Vic12 was studied over a 7-day period, the number of viable organisms per ml. measured (Gourlay & Leach, 1970) and the pH of the medium recorded. No difference was found in haemagglutinating activity of standard suspensions of these organisms measured after 2, 3, 4 and 7 days' growth.

The haemagglutinating activity of twelve strains for erythrocytes from nine animal species was measured. Samples from at least three animals of each species were tested on at least two occasions and the average haemagglutinating titre is shown in Table 3. The titre range was within one doubling dilution of the average given. The types of erythrocytes agglutinated by strain Mmb177 distinguished it from the eleven other strains examined. These other eleven strains showed a varied haemagglutinating activity. Strain Gri250 appeared the least active while, at the other end of the range strain Vic12 was the most active. Some of these eleven strains appeared rather similar but others could be distinguished on the basis of their haemagglutinating activity.

Effects of various treatments of red blood cells on their susceptibility to agglutination by M. dispar

Various treatments of red cells were carried out in an attempt to characterize the receptor site on the red cell surface. Strains F370 and Vic12 were chosen because they gave high haemagglutination titres and because their haemagglutinating

Table 3. *Haemagglutination titre* of M. dispar strains with erythrocytes from various animal species*

<i>M. dispar</i> strain	Source of erythrocytes								Guinea- pig
	Rabbit	Sheep	Hamster	Calf	Horse	Chicken	Rat	Human	
Vic12	32	16	16	8	16	16	8	16	8
D44	64	16	8	8	4	8	8	8	8
Vic13	32	4	4	8	2	2	1	2	1
F370	64	16	8	8	8	8	—	—	—
Vic7	32	8	4	4	1	1	1	1	1
D52	16	8	4	2	2	—	1	—	—
462/2	32	4	4	2	2	—	1	—	—
D2	32	4	4	1	1	1	—	—	—
Gri226	32	8	2	—	2	—	—	—	—
Gri221	32	2	—	—	—	—	—	—	—
Gri250	8	1	—	—	—	—	—	—	—
Mmb177	—	32	—	64	1	—	—	—	—

* Reciprocal of highest dilution of standard suspensions of *M. dispar* strains which agglutinated erythrocytes.

Table 4. *Effects of various treatments of erythrocytes on the haemagglutination titre of M. dispar strain F370*

Treatment	Titre* against erythrocytes from:		
	Rabbit	Calf	Sheep
Phosphate-buffered saline	64	16	32
Tris buffer	64	16	16
Pronase (0.02 %)	128	64	64
Trypsin (0.01 %)	128	64	32
Chymotrypsin (0.01 %)	64	32	32
Periodate (0.02 %)	32	16	8
Bisulphite (0.5 %)	64	16	32
RDE†	32	16	16

* Reciprocal of highest dilution of mycoplasma suspension causing haemagglutination.

† Receptor-destroying enzyme.

activity was representative of the majority of strains. The results obtained with both strains were essentially the same and those for strain F370 are given in Table 4. Erythrocytes from sheep, calves and rabbits were affected in essentially the same way by the reagents. Pronase and trypsin caused some red cells to become slightly more susceptible to agglutination. The other reagents had very little effect. The change with receptor-destroying enzyme was negligible compared with the change in susceptibility of erythrocytes to the stock suspension of influenza A virus, which had haemagglutination titres of < 4, 32 and 256 against calf, rabbit and sheep erythrocytes respectively before receptor-destroying-enzyme treatment and < 4 against all three types of erythrocytes subsequently.

Table 5. *Effects of various treatments of Mycoplasma dispar F370 on haemagglutinating activity*

Treatment	Titre* against erythrocytes from:		
	Rabbit	Calf	Sheep
Phosphate-buffered saline	32	8	16
Tris buffer	64	8	16
Pronase (0.02 %)	64	< 2	2
Trypsin (0.01 %)	64	< 2	< 2
Chymotrypsin (0.01 %)	64	8	16
Periodate (0.1 %)	16	8	16
Bisulphite (1.0 %)	32	8	4

* Reciprocal of the highest dilution of mycoplasma suspension causing haemagglutination.

Effect of various treatments of mycoplasmas on haemagglutinating activity

In an attempt to characterize the attachment site on *M. dispar* two strains, F370 and Vic12, were treated with various compounds. The treatments affected both in essentially the same way and the results obtained for strain F370 are given in Table 5. The effect of treatments on the agglutination of bovine and sheep erythrocytes seemed similar but was different from the effect of the treatments on the agglutination of rabbit erythrocytes. Pronase and trypsin reduced the haemagglutination titre for bovine and sheep erythrocytes, but not for rabbit erythrocytes. On the other hand periodate, bisulphite and chymotrypsin had little effect.

Electron-microscopy

M. dispar strain F370 was incubated with sheep and rabbit erythrocytes at 4° C. for 1 hr. The mixture was then examined by electron-microscopy. The capsular material previously described by Howard & Gourlay (1974) can be seen outside the membrane (Plate 1). Fine threads of material seem to bridge the gap between the dense extramembranous material and the erythrocyte membrane.

DISCUSSION

The twelve strains examined formed a serologically homogeneous group and no serological heterogeneity was observed to the extent that exists amongst certain other mycoplasma species, e.g. *M. hominis* (Purcell *et al.* 1967), *M. pulmonis* (Forshaw & Fallon, 1972) and *M. gallisepticum* (Taylor-Robinson & Berry, 1969). The metabolism inhibition test was chosen because of its sensitivity and specificity, and indirect haemagglutination because of its sensitivity (Purcell, Chanock & Taylor-Robinson, 1969).

Considerable strain variation in haemagglutinating activity was observed. Strain Mmb177 appeared distinct from the other eleven strains examined. The other strains exhibited a graded haemagglutinating potency. Although serological methods do not appear to be useful for distinguishing strains haemagglutination may be. Haemagglutinating activity appears unrelated to serological structure for *M. dispar* as is the case with *M. gallisepticum* (Manchee & Taylor-Robinson, 1969*b*).

In contrast to certain other mycoplasmas (Manchee & Taylor-Robinson, 1968) neither the incubation temperature for the haemagglutination reaction nor the final pH of the growth medium appear to affect the haemagglutinating potency of *M. dispar*.

M. dispar strain 462/2 has been reported to have no haemagglutinating activity for bovine and guinea-pig erythrocytes (Erno & Stipkovits, 1973). The slight difference between the previously reported results and ours for this strain probably does no more than reflect the use of different methods. The variation observed between strains of the same species of mycoplasma emphasizes the necessity to study more than one representative before comparisons between different mycoplasma species are made.

The nature of the receptor site for mycoplasmas on erythrocytes and other cells and the nature of the attachment site on mycoplasmas have been investigated by other workers using haemagglutination and haemadsorption. However, it should be noted that haemagglutination and haemadsorption are not necessarily correlated (Manchee & Taylor-Robinson, 1968). Strains of *M. dispar* rarely form centres (Gourlay & Leach, 1970). It is therefore difficult to perform haemadsorption experiments because colonies wash off the agar.

M. gallisepticum, *M. synoviae* and *M. pneumoniae* have been shown to attach to various cells by means of sialic-acid-containing cell receptor sites (Gesner & Thomas, 1966; Manchee & Taylor-Robinson, 1969*a, b*). It has been suggested that in the case of *M. hominis* and *M. salivarium* the receptor sites are protein (Manchee & Taylor-Robinson, 1969*a*).

The experiments performed here to determine the nature of the receptor sites on erythrocytes for *M. dispar* indicate that they do not contain sialic acid since they are resistant to receptor-destroying enzyme (cf. influenza A virus). Furthermore the observation that bovine red cells are not agglutinated by influenza A virus but are agglutinated by *M. dispar* supports the hypothesis that *M. dispar* does not attach to a similar sialic-acid-containing erythrocyte receptor. The erythrocyte receptor site is resistant to bisulphite and periodate at the concentrations used. Treatment with pronase and trypsin perhaps makes erythrocytes slightly more susceptible to agglutination by *M. dispar*, possibly by altering the surface charge on the cell, although there is no relation between either the zeta potential or electrophoretic mobility of red cells from different animal species (values taken from Spector, 1956) and susceptibility to agglutination by *M. dispar*.

The results reported by Sobeslavsky *et al.* (1968) indicate that the attachment site of *M. pneumoniae* contains a lipid or glycerophosphate hapten. In contrast *M. salivarium* and *M. hominis* have been reported to have protein attachment sites (Manchee & Taylor-Robinson, 1969*a*; Hollingdale and Manchee, 1972). The attachment site on *M. dispar* for bovine and sheep erythrocytes is sensitive to pronase and trypsin. Since neither of these proteolytic enzymes destroys the mycoplasma attachment site for rabbit erythrocytes at least two attachment mechanisms must exist on the mycoplasma. The attachment site for bovine and sheep erythrocytes is therefore either a protein or contains a protein moiety. The chemical nature of the attachment site for rabbit erythrocytes is unknown.

Savage (1972) suggested that surface components similar to capsules may be involved in cell attachment by lactobacilli and the possibility exists that the capsule of *M. dispar* observed by electron-microscopy after staining with ruthenium red (Howard & Gourlay, 1974) may take part in cell attachment.

As has been stated by Manchee & Taylor-Robinson (1969*a*) the ability to attach to cells should not be regarded as necessarily indicating pathogenicity *in vivo*, but as a valuable property for an invading mycoplasma which might assist in the disease process. Thus the demonstration of attachment sites on *M. dispar* for erythrocytes may indicate the mechanism by which this micro-organism maintains itself in the bovine respiratory tract.

We would like to thank Mrs P. Bland for the electron-micrograph.

REFERENCES

- ANDREWS, B. E., LEACH, R. H., GOURLAY, R. N. & HOWARD, C. J. (1973). Enhanced isolation of *Mycoplasma dispar* by substitution of ampicillin for benzyl penicillin in growth media. *Veterinary Record* **93**, 603.
- BUCKLAND, F. E. & TYRRELL, D. A. J. (1963). A comparative study of virus haemagglutinins. The stability of haemagglutinins and red cell receptors to certain physical and chemical treatments. *Journal of General Microbiology* **32**, 241-53.
- ERNO, H. & STIPKOVITS, L. (1973). Bovine mycoplasmas: cultural and biochemical studies, II. *Acta Veterinaria Scandinavica* **14**, 450-63.
- FORSHAW, K. A. & FALLON, R. J. (1972). Serological heterogeneity of *Mycoplasma pulmonis*. *Journal of General Microbiology* **72**, 501-10.
- GESNER, B. & THOMAS, L. (1966). Sialic acid binding sites: role in haemagglutination by *Mycoplasma gallisepticum*. *Science, New York* **151**, 590-1.
- GOURLAY, R. N. (1969). Isolation of a Mycoplasma-like organism from pneumonic calf lungs. *Veterinary Record* **84**, 229.
- GOURLAY, R. N. & LEACH, R. H. (1970). A new mycoplasma species isolated from pneumonic lungs of calves (*Mycoplasma dispar* sp. nov.) *Journal of Medical Microbiology* **3**, 111-23.
- GOURLAY, R. N., MACKENZIE, A. & COOPER, J. E. (1970). Studies of the microbiology and pathology of pneumonic lungs of calves. *Journal of Comparative Pathology* **80**, 575-84.
- GOURLAY, R. N. & THOMAS, L. H. (1969). Experimental pneumonia in calves produced by inoculation of mycoplasmas. *Veterinary Record* **85**, 583.
- HERBERT, W. J. (1967). Passive haemagglutination. In *Handbook of Experimental Immunology* (ed. D. M. Wier), pp. 720-44. Oxford and Edinburgh: Blackwell Scientific Publications.
- HOLLINGDALE, M. R. & MANCHEE, R. J. (1972). The role of mycoplasma membrane proteins in the adsorption of animal cells to *Mycoplasma hominis* colonies. *Journal of General Microbiology* **70**, 391-3.
- HOWARD, C. J. & GOURLAY, R. N. (1972). Serology of bovine T-mycoplasmas. *British Veterinary Journal* **128**, xxxvii-xl.
- HOWARD, C. J. & GOURLAY, R. N. (1973). Inhibition by normal rabbit sera of the growth of T-mycoplasma strains isolated from different animal species. *Journal of General Microbiology* **78**, 277-85.
- HOWARD, C. J. & GOURLAY, R. N. (1974). An electron-microscopic examination of certain bovine mycoplasmas stained with ruthenium red and the demonstration of a capsule on *Mycoplasma dispar*. *Journal of General Microbiology*. In the press.
- MANCHEE, R. J. & TAYLOR-ROBINSON, D. (1968). Haemadsorption and haemagglutination by mycoplasmas. *Journal of General Microbiology* **50**, 465-78.
- MANCHEE, R. J. & TAYLOR-ROBINSON, D. (1969*a*). Studies on the nature of receptors involved in attachment of tissue culture cells to mycoplasmas. *British Journal of Experimental Pathology* **50**, 66-75.

- MANCHEE, R. J. & TAYLOR-ROBINSON, D. (1969*b*). Utilization of neuraminic acid receptors by mycoplasmas. *Journal of Bacteriology* **98**, 914-19.
- PURCELL, R. H., CHANOCK, R. M. & TAYLOR-ROBINSON, D. (1969). Serology of the mycoplasmas of man. In *The Mycoplasmatales and the L-phase of Bacteria* (ed. L. Hayflick), pp. 221-64. Amsterdam: North-Holland Publishing Company.
- PURCELL, R. H., WONG, D. G., CHANOCK, R. M., TAYLOR-ROBINSON, D., CANCHOLA, J. & VALDESUSO, J. (1967). Significance of antibody to mycoplasma as measured by metabolic-inhibition techniques. *Annals of the New York Academy of Sciences* **143**, 664-75.
- ST GEORGE, T. D., HORSFALL, N. & SULLIVAN, N. D. (1973). A subclinical pneumonia of calves associated with *Mycoplasma dispar*. *Australian Veterinary Journal* **49**, 580-86.
- SAVAGE, D. C. (1972). Survival on mucosal epithelia, epithelial penetration and growth in tissues of pathogenic bacteria. *Symposium of the Society for General Microbiology* **22**, 25-57.
- SOBESLAVSKY, O., PRESCOTT, B. & CHANOCK, R. M. (1968). Adsorption of *Mycoplasma pneumoniae* to neuraminic acid receptors of various cells and possible role in virulence. *Journal of Bacteriology* **96**, 695-705.
- SPECTOR, W. S. (1956). *Handbook of biological data*. Philadelphia and London: W. B. Saunders.
- SPRINGER, E. L. & ROTH, I. L. (1973). The ultrastructure of the capsules of *Diplococcus pneumoniae* and *Klebsiella pneumoniae* stained with ruthenium red. *Journal of General Microbiology* **74**, 21-31.
- SUBCOMMITTEE ON THE TAXONOMY OF MYCOPLASMATALES (1972). Proposal for minimal standards for descriptions of new species of the order *Mycoplasmatales*. *International Journal of Systematic Bacteriology* **22**, 184-8.
- TAYLOR-ROBINSON, D. & BERRY, D. M. (1969). The evaluation of the metabolic-inhibition technique for the study of *Mycoplasma gallisepticum*. *Journal of General Microbiology* **55**, 127-37.
- TAYLOR-ROBINSON, D., PURCELL, R. H., WONG, D. C. & CHANOCK, R. M. (1966). A colour test for the measurement of antibody to certain mycoplasma species based upon the inhibition of acid production. *Journal of Hygiene* **64**, 91-104.
- THOMAS, L. H. & HOWARD, C. J. (1974). The effect of *Mycoplasma dispar*, *Mycoplasma bovirhinis*, *Acholeplasma laidlawii* and T-mycoplasmas on organ cultures of bovine trachea. *Journal of Comparative Pathology* **84**, 193-201.
- THOMAS, L. H. & SMITH, G. S. (1972). Distribution of mycoplasmas in the non-pneumonic bovine respiratory tract. *Journal of Comparative Pathology* **82**, 1-4.

EXPLANATION OF PLATE

Electron-micrograph of *M. dispar* following incubation with rabbit erythrocytes. Mm, *M. dispar* membrane; Em, erythrocyte membrane; C, capsular material; bar marker, 100 nm.