Published in final edited form as:

Insect Syst Evol. 2006 June ; 37(2): 197–212.

Phylogenetic position of the ant genus *Acropyga* Roger (Hymenoptera: Formicidae) and the evolution of trophophoresy

JOHN S. LAPOLLA,

John S. LaPolla, Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20013-7012 USA (lapollaj@si.edu)

TED R. SCHULTZ,

Ted R. Schultz, Department of Entomology, National Museum of Natural History, Smithsonian Institution, Washington, D.C. 20013-7012 USA (schultz@lab.si.edu)

KARL M. KJER, and

Karl M. Kjer, Department of Ecology and Evolution, Rutgers University, New Brunswick, NJ 08901 (kjer@aesop.rutgers.edu)

JOSEPH F. BISCHOFF

Joseph F. Bischoff, National Institutes of Health, National Center for Biotechnology Information Bethesda, Maryland 20892-6510 (bischoff@ncbi.nlm.nih.gov)

Abstract

Trophophoresy is exhibited in two ant genera: *Acropyga* (Formicinae), in which all 37 species are thought to be trophophoretic, and *Tetraponera* (Pseudomyrmecinae), in which it has been observed in only one species, *T. binghami*. This study analyses a dataset comprised of both morphological and molecular (D2 region of 28S rRNA and EF1- α) data. Evidence is presented in favor of *Acropyga* being monophyletic, hence trophophoresy has evolved only once within the Formicinae and twice within the ants overall. The data further suggests that *Acropyga* belongs within a clade containing *Anoplolepis*, *Aphomomyrmex*, and *Petalomyrmex*. *Aphomomyrmex* and *Petalomyrmex* were found to be the sister group to *Acropyga*. The results indicate that the Lasiini and Plagiolepidini are not monophyletic and are in need of reexamination. Given the extant pantropical distribution of *Acropyga* it is speculated that *Acropyga* maybe of Gondwanan origin and that trophobiosis was the first form of agriculture to evolve in the ants.

Introduction

The evolution of agriculture in ants conforms to two broad patterns: “farming” as practiced by fungus-growing ants and “herding” as practiced by trophobiont-tending ants. Fungiculture, practiced by the attine ants (tribe Attini), has evolved only once within the ants (for reviews, see Hölldobler and Wilson 1990; Mueller et al. 1998; Schultz et al. 2005). In a complex agricultural symbiosis that appears to be the product of ancient coevolutionary interactions, attine fungiculture is the product of a mutualism between the ants, their fungal cultivars, and an antibiotic-producing filamentous bacterium (Schultz et al., 2005).

Trophobiosis, in contrast, has evolved multiple times in ants, most commonly in the subfamilies Formicinae, Dolichoderinae, and Myrmicinae subfamilies, and more rarely in the Ponerinae, Ectatomminae, and Pseudomyrmecinae (Way 1963; Hölldobler and Wilson 1990). In trophobiosis, ants acquire honeydew, the sugar-rich fluid excreted by sap-feeding insects (the trophobionts: aphids and other sternorrhynchan insects in the order Hemiptera or certain lycaenid and riodinid caterpillars in the order Lepidoptera); in return, the ants protect these insects from parasites and predators (Hölldobler and Wilson 1990). For most ants, trophobiosis is facultative (Hölldobler and Wilson 1990). In a very few ant species, however, trophobiosis

is obligate. Some obligately trophobiotic ant species have acquired particular adaptations for ensuring the transfer of trophobionts from parent to daughter colonies. Among the nomadic herding *Dolichoderus* species of Southeastern Asia, for example, colonies reproduce by budding (Dill and Maschwitz 1998), in which a portion of the colony, including a newly mated daughter queen, workers, and mealybugs (Hemiptera: Pseudococcidae), separate from the parent colony and thereafter lead an independent existence. Colony budding may also occur in some *Pseudolasius* spp. that also depend on trophobionts (Acleridae, Aphidae, and Pseudococcidae) (Malsch et al. 2001; Kaufmann et al. 2003). In other obligately trophobiotic ant species, virgin queens carry trophobionts when they depart from the nest to mate and found new colonies. In the case of the plant-ant, *Aphomomyrmex afer*, mealybugs of the species *Paraputo anomala* cling to the virgin queens body when it leaves its birth nests, and these mealybugs subsequently serve as foundresses for a new mealybug “herd” in the new ant colony (Gaume et al. 2000). In another form of carrying behavior, the virgin queen departs on its mating flight carrying the trophobiont between its mandibles.

This latter form of trophobiosis has been termed trophophoresy (LaPolla et al. 2002). Only two ant genera are known to contain trophophoretic species: *Acropyga* (Formicinae), in which all 37 species are thought to be trophophoretic, and *Tetraponera* (Pseudomyrmecinae), in which trophophoresy has been observed in only one species, *T. binghami*. The details of the behavior remain largely unknown (Ward 2001). These two genera are members of demonstrably distantly related subfamilies (Baroni Urbani et al. 1992; Ward & Brady 2003). In both *Acropyga* and *T. binghami*, virgin queens depart from their parent nests carrying between their mandibles a trophobiont (in both genera a mealybug) that will serve as the progenitor of a new “herd” in their newly founded nests (Klein et al. 1992; LaPolla et al. 2002).

Acropyga are small ants (typically between 1–2 mm in total length), and live almost entirely underground where they place rhizoecine mealybugs (Hemiptera: Pseudococcidae) on roots to feed. The ants and mealybugs are apparently mutually dependent on each other for survival (Flanders 1957; LaPolla et al. 2002). The degree of species specificity between ant and mealybug remains unclear, but there is evidence that co-evolution has occurred between the two groups (LaPolla 2004). Fossil *Acropyga* are known from Dominican amber, indicating that trophophoresy is at least 15–20 mya old (fig. 1; Johnson et al. 2001; LaPolla 2004; LaPolla 2005).

The taxonomy of *Acropyga* has until recently been confused (LaPolla 2004), and the monophyly of the genus has been questioned. It has been suggested the genus was either paraphyletic or polyphyletic (Agosti 1991; Bolton 1994), with either scenario having important implications for the evolution of trophophoresy (i.e., implying that trophophoresy may have arisen more than once within the Formicinae and more than twice within the ants overall). While LaPolla (2004) presented seven putative morphological synapomorphies supporting *Acropyga* monophyly, that study did not examine the placement of the genus within the subfamily Formicinae. Here we provide evidence for the phylogenetic position of *Acropyga* drawn from both morphological and molecular data.

Materials and Methods

Source of Material

Twenty-six taxa were examined (table 1) for analysis. The sampling of the ingroup, *Acropyga* spp., was constrained by the availability of specimens for DNA sequencing. However, of the four species included in the analysis, *A. arnoldi* is the putative sister group to the rest of the genus, and the remaining species represent different, distantly related clades in the phylogeny of *Acropyga* (LaPolla 2004). The sampling of the diversity of the genus is therefore considered sufficient for the purposes of the present study.

Specimens were examined from a number of research collections, and below follows the list of the institutions and individuals' collections that contributed to this study. Voucher specimens for every species used in DNA work are deposited in the National Museum of Natural History (USNM).

INBC: Instituto Nacional de Biodiversidad, Santo Domingo de Heredia, Costa Rica

JSLC: J.S. LaPolla Collection, private collection

JTLC: J.T. Longino Collection, Evergreen State College, Olympia, WA, USA

LACM: Natural History Museum of Los Angeles County, Los Angeles, CA, USA

MCZC: Museum of Comparative Zoology, Harvard University, Cambridge, MA, USA

PSWC: P.S. Ward Collection, University of California at Davis, CA, USA

SAMC: South African Museum, Cape Town, South Africa

SYPC: S. Yamane Collection, Kagoshima University, Kagoshima, Japan

UGBC: Centre for the Study of Biological Diversity, University of Guyana, Georgetown, Guyana

USNM: National Museum of Natural History, Washington DC, USA

Morphological Data

Characters were examined using a light microscope (Nikon SMU-Z) or a scanning electron microscope (Hitachi S-510). Morphological terminology employed throughout follows Bolton (1994), with modifications where noted.

Morphological Characters

The morphological characters and their alternative states are listed below. Table 2 lists the states assigned to each taxon. Character state designations (0, 1, 2) have no significance with respect to polarity.

1. Antennal segment number: (0) 12 segments; (1) 11 or fewer segments.
2. Antennal torulae (fig. 2a–b): (0) not tubular; (1) tubular.
3. Eye size: (0) 30 or more ommatidia; (1) 30 or fewer ommatidia.
4. Eye position: (0) posterior end of eye fails to reach the midline; (1) posterior end of eye surpasses approximate midline of head.
5. Palp formula: (0) 6:4; (1) less than 6:4.
6. Mandibular teeth number: (0) 6 teeth or greater; (1) fewer than 6 teeth.
7. Propodeal spiracle: (0) small and round; (1) large and round; (2) oval-shaped or slit-like.
8. Metapleural gland: (0) present; (1) absent.
9. Petiole muscle orifice (fig. 2c–d): (0) round; (1) oval.

10. Petiole foramen (adapted from Bolton, 2003) (fig. 2e–g): (0) fails to reach anterior of metacoxal cavities; (1) surpasses anterior of metacoxal cavities; (2) reaches anterior of metacoxal cavities.
11. Petiole ventral surface (fig. 3a–b): (0) with ventral extension; (1) without ventral extension.
12. First gastral sternite (third abdominal) (fig. 3c–d): (0) surpasses petiole articulation; (1) does not surpass petiole articulation.
13. Proventriculus: (0) asepalous; (1) sepalous.
14. Protibial spur (fig. 3e–f): (0) long; (1) short.
15. Acidopore: (0) absent; (1) present

Molecular Data

For DNA extraction a whole ant was selected from among several nestmates in a vial; the remaining specimens are deposited as vouchers in the USNM. When possible specimens were identified to species. Samples were placed in labeled Eppendorf tubes and ground under liquid nitrogen, using microtissue grinders (Phoenix Research). DNA was extracted with sodium dodecyl sulfate, Proteinase-K, and phenol/chloroform, as described in Kjer et al. (2001). Dried DNA pellets were resuspended in 50–250 µl of Tris-EDTA buffer. Material was separated out into two tubes, one of which was added to a stock DNA collection and kept at –70°C, and the other kept in a freezer for amplification by polymerase chain reaction (PCR).

D2 region of 28S ribosomal RNA Protocol

D2 sequences were amplified on a thermal cycler using the reaction conditions described in Kjer et al. (2001). Primer sequences for the D2 were: 5'-GAGTTCAAGAGTACGTGAAACCG-3' paired with 5'-CCTTGGTCCGTGTTTCAAG-AC-3'. PCR conditions were 95°C, 30s; 52°C, 60s; 72°C, 60s for 35–45 cycles. Amplified DNA was separated on a 1.5% low-melting-point agarose gel (NuSieve 3:1; FMC Bioproducts). Bands of DNA were cut from the agarose gel, purified with GeneClean (Bio101), and sequenced on an ABI 377 automated sequencer (Applied Biosystems). Sequences were completed in both strands and edited manually with the assistance of Sequence Navigator (Applied Biosystems). During the editing of each strand, nucleotides that were readable but either showed irregular spacing between peaks or had some important competing background peak were coded in lowercase letters. These letters were: 1) converted to uppercase if the complementary strand strongly confirmed them; 2) left lowercase when both strands were lowercase; 3) changed to Ns (nucleotide undermined) when strands were contradictory. Sequences are deposited at Genbank under accessions DQ 226030–DQ 226054.

EF 1-alpha

EF 1-alpha sequence data (F1 copy) were generated for an approximately 1.2KB fragment (representing 364 amino acid residues and an intron of variable size, approximately 0.1KB in length). Extracted DNA was generally amplified in two steps: In the first step, a fragment approximately 2KB in length was amplified using the primers M3 (forward) and TRS10R (reverse) with a 55°C annealing temperature. Then, using the product of the initial amplification as template, 4 shorter overlapping fragments were reamplified with a variable (58°C to 64°C) annealing temperature, using as primers (i) For1 and Rev1.1, (ii) U377.1 and TRS1R, (iii) Cho7.1 and L53 or U52.1 and L53, and (iv) TRS4F and TRS9.1R. Primer sequences are listed in Table 3. Sequences are deposited at Genbank under accessions DQ 226013–DQ 226029.

Phylogenetic Analyses Methods

After eliminating the intron sequence, alignment of the 1092 bp of EF 1- α nucleotide sequence was trivial, as EF 1- α is a protein-coding gene and amino acid number is highly conserved across all ants (TRS, unpublished). DNA sequence alignments are available upon request. The D2 region of the 28S rDNA gene was aligned according to secondary structure (Gutell 1994; annotation following Kjer 1995). Our alignment is reproduced in appendix 1.

Six data subsets were separately analyzed: (1) the D2 region of the 28S rDNA gene (25 species), (2) EF 1- α (26 species), (3) morphology (23 species), (4) combined 28S, EF 1- α , and morphology (26 species), (5) maximum combined EF 1- α and 28S (26 species, 10 incomplete for one or the other gene), and (6) minimum combined EF 1- α and 28S (16 species, complete for both genes).

Parsimony (MP) analyses

MP analyses were conducted in PAUP 4.0b10 (Swofford 2002) using the heuristic search option with TBR branch-swapping and 1000 random-taxon-addition replicates; successive-approximations weighting (SW) analyses used 500 replicates. Heuristic-search bootstrap analyses (Felsenstein 1984) used TBR branch-swapping and consisted of 1000 pseudoreplicates, with 10 random-taxon-addition replicates per pseudoreplicate. In all cases gaps are treated as missing. MP analyses of the 28S data set and the combined (28S, EF 1- α , and morphology) data utilized four step-matrix characters representing four D2 variable regions as coded in the INNASE computer application (Lutzoni et al. 2000).

Maximum-likelihood (ML) analyses

Nucleotide substitution models for ML analyses were evaluated with the relevant nucleotide data and the MP (EF 1- α) or SW (28S, combined) tree using the Akaike Information Criterion (AIC) calculated in ModelTest 3.06 (Posada and Crandall 1998). ML analyses were conducted in PAUP 4.0b10 (Swofford 2002). Heuristic searches employed the adopted model and an optimal MP tree as the branch-swapping starting tree and consisted of five iterative sub-searches, each iteration employing updated model parameter values and topologies based on the results of the preceding iteration and each using successively more intensive branch-swapping regimes (Currie et al. 2003; Sallum et al. 2002; Mueller et al. 1998; Villesen et al. 2004). Heuristic ML bootstrap analyses employed TBR branch-swapping and consisted of 1000 pseudoreplicates in which parameter values were constrained to those estimated on the optimal ML tree.

Bayesian analyses

Bayesian analyses were conducted in MrBayes 3.0b4 (Huelsenbeck and Ronquist 2001). Analyses of the EF 1- α data and the combined data employed multiple character partitions governed by separate, unlinked models based on the model-fitting analyses described above and subject to the limitations of models available in MrBayes. Two character partitions were employed in analyses of EF 1- α : (i) first and second codon positions and (ii) third codon positions. Four character partitions were employed in analyses of the combined data set: (i) 28S D2, (ii) EF 1- α first and second codon positions, (iii) EF 1- α third codon positions, and (iv) morphology. Analysis of each data set included four separate runs, each consisting of one million MCMC generations and four simultaneous MCMC chains (three heated), and each with a “burn-in” of 200K generations. For each analysis, post-burn-in trees from all four runs were pooled to calculate posterior probabilities.

Results and Discussion

Phylogenetic Analyses

Morphology—MP analyses of the morphology data produced 90 equally parsimonious trees (MPTs) with parsimony-informative length = 38, C.I. = 0.474, R.I. = 0.804; successive approximations weighting favored a subset of 12 of these trees (fig. 4).

28S D2: MP analyses of the 28S D2 nucleotide data produced 2510 MPTs with parsimony-informative length = 220, C.I. = 0.668, R.I. = 0.610; successive approximations weighting favored a subset of three of these trees. For one of these successive approximations weighting trees, the AIC favored the TVM+G model. ML analysis with this model identified a single tree with log likelihood of -1928.44657 . Bayesian analyses using the GTR+G model (the closest model available in the MrBayes application) identified a tree similar to the ML tree.

EF 1-alpha: MP analyses of the EF 1-alpha data produced a single MPT with parsimony-informative length = 590, C.I. = 0.595, R.I. = 0.493. For this tree and the complete EF 1-alpha sequence data, the AIC favored the TrN+I+G model. ML analysis using this model identified a single tree with log likelihood of -4280.97199 . For this ML tree and the subset of EF 1-alpha sequence data consisting of codon positions 1 and 2, the AIC favored the TrN+I model; for this ML tree and codon positions 3, the AIC favored the TVM+G model. Bayesian analyses using character partitions governed by the closest available models in the mrbayes application, GTR+I and GTR+G, respectively, identified a nearly identical tree, differing only with regard to the relationships of BRA, CMM, and PRE.

Combined morphology, EF 1-alpha, and 28S D2, 26 species—MP analyses of the full 26-taxon data set, including all four MP character data partitions (morphology, 28S D2, EF 1-alpha, and the four INNASe step matrices), produced 176 MPTs with parsimony-informative length = 749, C.I. = 0.526, R.I. = 0.533; SW favored a subset of 7 of these trees. For one of these trees and the combined 28S D2 and EF 1-alpha sequence data, the AIC favored the GTR+I+G model. ML analysis using this model identified a single tree with log likelihood of -6331.4013 . Based on the preceding analyses, Bayesian analyses of the 26-species data set utilized the following character partitions and models: (i) morphology with Tuffley-Steele parsimony; (ii) 28S D2 with GTR+G; (iii) EF 1-alpha first and second codon positions with GTR+I; and (iv) EF 1-alpha third codon positions with GTR+G (fig. 5).

Combined morphology, EF 1-alpha, and 28S D2, 16 species—Because the 26-species data set contained 10 species lacking sequences for one of the two genes, and because these missing data adversely affected branch support in the resulting trees, analyses were carried out for the 16-species data set for which all sequence data were complete. MP analyses of the 16-species data set, including all four MP character data partitions (morphology, 28S D2, EF 1-alpha, and the four INNASe step matrices) produced 3 MPTs with parsimony-informative length = 643, C.I. = 0.533, R.I. = 0.502 (fig. 6A); SW favored one of these trees. For this tree and the combined 28S D2 and EF 1-alpha sequence data, the AIC favored the GTR+I+G model. ML analysis using this model identified a single tree with log likelihood of -5839.43443 . Based on the preceding analyses, Bayesian analyses of the 16-species data set utilized the following character partitions and models: (i) morphology with Tuffley-Steele parsimony; (ii) 28S D2 with GTR+G; (iii) EF 1-alpha first and second codon positions with GTR+I; and (iv) EF 1-alpha third codon positions with GTR+G (fig. 6B).

The combined analyses of both morphological and molecular (D2 and EF-1a; Fig. 6) datasets (with a reduced number of taxa, excluding those without both molecular datasets [16-taxa]) recovered a well-supported monophyletic *Acropyga* (100% bootstrap values [parsimony and

likelihood] and a 100% posterior probability). *Acropyga* was found within a clade containing *Anoplolepis* and *Petalomyrmex*. The sister group for *Acropyga* was found to be *Petalomyrmex*, with high support (98% MP bootstrap, 96% ML bootstrap, and 100% posterior probability).

Our combined morphological and molecular dataset of three *Acropyga* (for the 16-taxon dataset), while relatively limited, is significant because it contains *A. arnoldi*, the sister taxon to all other *Acropyga* (LaPolla 2004), and a member of the derived *decedens* species-group, *A. epedana* (LaPolla 2004), thereby providing phylogenetic depth to the analysis. We can therefore state that *Acropyga* is monophyletic, corroborating the morphology-based study of LaPolla (2004). We can also state that trophophoresy has probably evolved only once among the formicine ants (there is always the possibility the behavior will be observed in another group), and possibly twice among the ants overall (Klein et al., 1992; Ward, 2001).

In all analyses, except morphology only, *Petalomyrmex phylax* was found to be the sister group of *Acropyga*. The only exception is the result of the morphology-only analysis, in which all conclusions about genus-level relationships lacked statistical support. *Petalomyrmex* is a monotypic genus whose sister group is *Aphomomyrmex* (Snelling 1979; Chenuil and McKey 1996), another monotypic genus. Both genera are plant-ants restricted to the Afrotropical region. A close relationship between *Petalomyrmex* (and, by implication, *Aphomomyrmex*) and *Acropyga* is intriguing because, whereas *Petalomyrmex phylax* is not trophobiotic, *Aphomomyrmex afer* is obligately trophobiotic (Gaume et al. 2000) and practices an unusual method for acquiring mealybugs. Mealybugs of the species *Paraputo anomala* cling to the bodies of virgin *Aphomomyrmex afer* queens when those queens depart from the natal nest to mate and found new colonies. This phoretic behavior on the part of the mealybug is thought to benefit the ants because, first, it provides a mealybug that will serve as a “seed individual” for producing a mealybug “herd,” and, secondly, because the queen does not have to carry the mealybug in her mandibles, she is free to use her mandibles to excavate an opening for her nest chamber (Gaume et al. 2000). Gaume et al. (2000) speculated that, because the mealybug clings to the queen’s body, there is little possibility that it will wander off and become separated from the queen. It bears noting that *Acropyga* queens, which carry the mealybug in their mandibles, are presumably presented with this same problem, and that perhaps the mealybug does not wander when put down by the queen after all. It is known that *Acropyga* queens sometimes return to their birth nests after mating (Bünzli 1935); in such cases they do not need to excavate a nest cavity. But independent nest founding has also been observed in *Acropyga* (Bünzli 1935; LaPolla et al. 2002) and independently-founding queens obviously need to excavate nest cavities. Morphological characters of male genitalia (the placement of the penis valve apodemal ridge) of *Acropyga*, *Aphomomyrmex*, and *Petalomyrmex* also support a close relationship between these genera (LaPolla and Longino, 2006).

Origins of *Acropyga*

Acropyga is at least 20 million years old based on the presence of Dominican amber fossils (Johnson et al. 2001). However, the genus is certainly much older than these fossils. For example, the current distribution of *Acropyga* is pantropical. This is at least consistent with a Gondwanan origin of the genus, especially because the species-level phylogeny of LaPolla (2004) found all Old World species to be paraphyletic from which arises a clade of New World species. This suggests a vicariant pattern, preceded by a Gondwanan distribution predating the complete separation of the southern continents, especially because *Acropyga* species are poor dispersers (Bünzli 1935). Under a scenario in which extant *Acropyga* distribution is due to vicariance a more ancient origin for *Acropyga*, perhaps in the mid to late Cretaceous, is plausible. Additional data are required to confidently test this hypothesis.

Certainly, honeydew from trophobionts was readily available in the Cretaceous, with scale insects making up nearly 10% of all insect samples from New Jersey and Canadian amber (Grimaldi and Agosti 2001). The recent discovery of a fossil formicine ant from New Jersey amber (92 mya), *Kyromyrmex neffi*, suggests that the subfamily originated fairly early in ant evolution (Grimaldi and Agosti 2001). An ancient origin for *Acropyga* is thus not impossible. Contrasting *Acropyga* agriculture with other forms of ant agriculture may provide insight into general patterns of agricultural evolution. *Acropyga* fossils from Dominican amber represent the oldest definitive record of trophobiosis (Johnson et al. 2001; LaPolla 2005). Therefore, we can infer that *Acropyga* are certainly among the oldest “agricultural ants,” however, the question of just how old still remains to be answered with the collection of more data. Trophophoresy is probably older than fungiculture in ants. Attine ants are known only from the New World, and there is no evidence that they have ever occurred outside of this region. There are fossil attines from Dominican amber. Thus, *Acropyga* could have entered trophobiotic relationships very early in the course of ant evolution. Quaintly put, this could mean that ants were herdsman before they were farmers.

Comments on the Lasiine Tribe Group

Bolton (2003) recently proposed the lasiine tribe group which consists of three Formicinae tribes: Lasiini, Myrmoteratini, and Plagiolepidini. While Bolton acknowledged that the Lasiini and Plagiolepidini might need to be combined in the future, this study indicates that even combination of the two would not result in monophyly. With strong support for the *Acropyga/Anoplolepis/Petalomyrmex* clade at the apex of the tree, recovery of the two tribes as monophyletic seems very unlikely even with the addition of taxa. A thorough review of the Lasiini and Plagiolepidini is needed in order to redefine the tribes in light of the current lack of monophyly. In fact, across the Formicinae a phylogenetic review of tribal classification is much needed. We recovered a monophyletic Camponotini, but our dataset cannot address the status of the other tribes. Perhaps as Shattuck (1992) concluded after analyzing the Dolichoderinae, the time has come to seriously reconsider the use of the tribal rank within the Formicinae as well, but this can only be considered after a well supported phylogeny of generic relationships within the subfamily has been proposed.

Acknowledgements

TRS was funded by National Science Foundation awards DEB 0110073 and DEB 0431330. Field work for JSL was funded in part by the Smithsonian NMNH's Biological Diversity of the Guiana Shield Program and is #100 in BDG's publication series. For collection permits we are grateful to The Office of the President of Guyana and the Guyana Environmental Protection Agency. Specimens were graciously provided by: R. Crozier, B. Fisher, J. Longino, A. Moog, H. Robertson, P. Ward, and S. Yamane. Sequencing of EF 1-alpha was carried out in the Smithsonian NMNH's Laboratories of Analytical Biology. We thank B. Danforth, T. Friedlander, and J. Regier for guidance and primers; M. Braun, K. Carol, C. Huddleston, S. Rehner, J. Sweere Sakamoto, and L. Zimmer for generous advice in the lab; and F. Dahlan and E. Okonski for critical technical support.

References

- Agosti D. Revision of the ant genus *Cladomyrma*, with an outline of the higher classification of the Formicinae (Hymenoptera: Formicidae). *Systematic Entomology* 1991;16:293–310.
- Bolton, B. Identification guide to the ant genera of the world. Harvard University Press; Cambridge, Mass: 1994. p. 222
- Bolton, B. A new general catalogue of the ants of the world. Harvard University Press; Cambridge, Mass: 1995. p. 504
- Bolton B. Synopsis and classification of Formicidae. *Memoirs of the American Entomological Institute* 2003;71:1–370.
- Baroni Urbani C, Bolton B, Ward PS. The internal phylogeny of ants (Hymenoptera: Formicidae). *Systematic Entomology* 1992;17:301–329.

- Bünzli GH. Untersuchungen über coccidophile Ameisen aus den Kaffeefeldern von Surinam. Mitteilungen der Schweizerischen Entomologischen Gesellschaft 1935;16:455–593.
- Chenuil A, McKey DB. Molecular phylogenetic study of a myrmecophyte symbiosis: did *Leonardoxa*/ant associations diversify via cospeciation? Molecular Phylogenetics and Evolution 1996;6:270–286. [PubMed: 8899728]
- Currie CR, Wong B, Stuart AE, Schultz TR, Rehner SA, Mueller UG, Sung GH, Spatafora JW, Straus NA. Ancient Tripartite Coevolution in the Attine Ant-Microbe Symbiosis. Science 2003;299 (5605): 386–388. [PubMed: 12532015]
- Dill M, Maschwitz U. The nomadic herdsmen of Kinabalu Park: A unique lifestyle in ants. Sabah Parks Nature Journal 1998;1:61–67.
- Felsenstein J. Distance methods for inferring phylogenies: a justification. Evolution 1984;38:16–24.
- Flanders SE. The complete interdependence of an ant and a coccid. Ecology 1957;38:535–536.
- Gaume L, Matile-Ferrero D, McKey D. Colony foundation and acquisition of coccid trophobionts by *Aphomyrmex afer* (Formicinae): co-dispersal of queens and phoretic mealybugs in an antplant-homopteran mutualism? Insectes Sociaux 2000;47:84–91.
- Grimaldi D, Agosti D. A formicine in New Jersey Cretaceous amber (Hymenoptera: Formicidae) and early evolution of the ants. Proceedings of the National Academy of Sciences 2001;97(25):13678–13683.
- Gutell RR. Collection of small subunit (16S- and 16S-like) ribosomal RNA structures. Nucleic Acids Research 1994;22:3502–3507. [PubMed: 7524024]
- Hölldobler, B.; Wilson, EO. The ants. Harvard University Press; Cambridge, Mass: 1990. p. 732
- Huelsenbeck JP, Ronquist F. MrBayes: Bayesian inference of phylogeny. Bioinformatics 2001;17:754–755. [PubMed: 11524383]
- Johnson C, Agosti D, Delabie JH, Dumpert K, Williams DJ, von Tschirnhaus M, Maschwitz U. *Acropyga* and *Azteca* ants (Hymenoptera: Formicidae) with scale insects (Sternorrhyncha: Coccoidea): 20 million years of intimate symbiosis. American Museum Novitates 2001;3335:1–18.
- Kaufmann E, Malsch AKF, Williams DJ, Maschwitz U. *Pseudolasius* spp. (Formicinae) and their trophobionts - codispersal by colony budding. Sociobiology 2003;41:429–436.
- Kjer KM. Use of rRNA secondary structure in phylogenetic studies to identify homologous positions: an example of alignment and data presentation from the frogs. Molecular Phylogenetics and Evolution 1995;4(3):314–330. [PubMed: 8845967]
- Kjer KM, Blahnik RJ, Holzenthal R. Phylogeny of Trichoptera (Caddisflies): characterization of signal and noise within multiple datasets. Systematic Biology 2001;50:781–816. [PubMed: 12116634]
- Klein RW, Kovac D, Schellerich A, Maschwitz U. Mealybug-carrying by swarming queens of a Southeast Asian bamboo-inhabiting ant. Naturwissenschaften 1992;79:422–423.
- LaPolla JS. *Acropyga* (Hymenoptera: Formicidae) of the World. Contributions of the American Entomological Institute 2004;33(3):1–130.
- LaPolla JS. Ancient trophophoresy: A fossil *Acropyga* (Hymenoptera: Formicidae) from Dominican Amber. Transactions of the American Entomological Society 2005;131 (12):21–28.
- LaPolla JS, Cover SP, Mueller UG. Natural history of the mealybug-tending ant *Acropyga epedana*, with descriptions of the male and queen castes. Transactions of the American Entomological Society 2002;128(3):367–376.
- LaPolla JS, Longino JT. An Unusual New *Brachymyrmex* Mayr (Hymenoptera: Formicidae) from Costa Rica, with Implications for the Phylogeny of the Lasiine Tribe Group. Proceedings of the Washington Entomological Society 2006;108(2):297–305.
- Lutzoni F, Wagner P, Reeb V, Zoller S. Integrating ambiguously aligned regions of DNA sequences in phylogenetic analyses without violating positional homology. Systematic Biology 2000;49:628–651. [PubMed: 12116431]
- Malsch AKF, Kaufmann E, Heckroth HP, Williams DJ, Maryati M, Maschwitz U. Continuous transfer of subterranean mealybugs (Hemiptera, Pseudococcidae) by *Pseudolasius* spp. (Hymenoptera, Formicidae) during colony fission. Insectes Sociaux 2001;48(4):333–341.
- Mueller UG, Rehner SA, Schultz TR. The evolution of agriculture in ants. Science 1998;281:2034–2038. [PubMed: 9748164]

- Posada D, Crandall KA. Modeltest: testing the model of DNA substitution. *Bioinformatics* 1998;14:817–818. [PubMed: 9918953]
- Sallum MAM, Schultz TR, Foster PG, Aronstein K, Wirtz RA, Wilkerson RC. Phylogeny of Anophelinae (Diptera: Culicidae) based on nuclear ribosomal and mitochondrial DNA sequences. *Systematic Entomology* 2002;27:361–382.
- Schultz, TR.; Mueller, UG.; Currie, CR.; Rehner, SA. Fernando Vega and Meredith: Ecological and Evolutionary Advances in Insect-Fungal Associations. Oxford University Press; New York: 2005. Reciprocal illumination: A comparison of agriculture in humans and ants; p. 149-190.
- Shattuck SO. Higher classification of the ant subfamilies Aneuretinae, Dolichoderinae, and Formicinae (Hymenoptera: Formicidae). *Systematic Entomology* 1992;17:199–206.
- Snelling RR. *Aphomomyrmex* and a related new genus of arboreal African ants. *Contributions in Science* 1979;316:1–8.
- Swofford, DL. PAUP*. Phylogenetic Analysis Using Parsimony (*and Other Methods). Sinauer Associates; Sunderland, Massachusetts: 2002.
- Villesen P, Mueller UG, Schultz TR, Adams RMM, Bouck MC. Evolution of ant-cultivar specialization and cultivar switching in *Apterostigma* fungus-growing ants. *Evolution* 2004;58:2252–2265. [PubMed: 15562688]
- Ward PS. Taxonomy, phylogeny, and biogeography of the ant genus *Tetraponera* (Hymenoptera: Formicidae) in the Oriental and Australian regions. *Invertebrate Taxonomy* 2001;15:589–665.
- Ward PS, Brady SG. Phylogeny and biogeography of the ant subfamily Myrmeciinae (Hymenoptera: Formicidae). *Invertebrate Systematics* 2003;17:361–386.
- Way MJ. Mutualism between ants and honeydew-producing Homoptera. *Annual Review of Entomology* 1963;8:307–344.

Fig. 1. Ancient trophophoresy. *Acropyga glaesaria* queen in Dominican amber with a mealybug (*Electromyrmococcus abductus*) held in her mandibles (see LaPolla 2005).

Fig. 2. Morphological characters used in phylogenetic analysis. Antennal torulae: A) not tubular, B) tubular; petiolar muscle orifice: C) round, D) oval; Petiole foramen: E) fails to reach anterior of metacoxal cavities, F) surpasses anterior of metacoxal cavities, G) reaches anterior of metacoxal cavities.

Fig. 3. Morphological characters used in phylogenetic analysis. Petiolar ventral surface: A) with ventral extension, B) without ventral extension; first gastral sternite: C) surpasses petiole articulation, D) does not surpass petiole articulation; protibial spur: E) long; F) short.

Figure 4.

Cladogram of strict consensus tree resulting from a heuristic search of morphological data summarizing the 90 shortest trees to estimate relationships of Formicinae (tree length= 38 steps).

Fig. 5. Phylogram of Formicinae resulting from a Bayesian analysis of the full 26-taxon dataset. Asterisks (*) indicate taxa for which data were available from both genes (EF1-alpha and D2), i.e. those taxa constituting the “16-taxon dataset.” Support values from the analysis of the 16-taxon dataset are indicated. The break in the line for *Acropyga acutiventris* represents a long branch that has been shortened for image size constraints.

Fig. 6.

Analysis of the 16-taxon data set; arrows indicate taxa that differ in position from the 26-taxon dataset. A) Consensus of 3 MPTs resulting from the parsimony analysis; numbers above branches are bootstrap values. B) Phylogram resulting from the Bayesian analysis; numbers above branches are Bayesian posterior probabilities, below are bootstrap values from the maximum likelihood analysis.

Table 1
Species used in analysis with GenBank accession numbers.

	28S rRNA	Ef1-alpha	Origin of Specimen
<i>Acropyga acutiventris</i>	DQ226030		Malaysia
<i>Acropyga arnoldi</i>	DQ226031	DQ226013	South Africa
<i>Acropyga donisthorpei</i>	DQ226032	DQ226014	Brazil
<i>Acropyga epedana</i>	DQ226033	DQ226015	USA
<i>Amblyopone pallipes</i>	DQ226034	DQ226016	USA
<i>Anoplolepis gracilipes</i>	DQ226035	DQ226017	USA
<i>Anoplolepis steingroeveri</i>		DQ226018	South Africa
<i>Brachymyrmex depilis</i>	DQ226036	DQ226019	USA
<i>Camponotus chromaiodes</i>	DQ226037	DQ226020	USA
<i>Cladomyrma maryatae</i>	DQ226038	DQ226021	Malaysia
<i>Dorymyrmex</i> sp.	DQ226039	DQ226022	USA
<i>Formica subsericea</i>	DQ226040	DQ226023	USA
<i>Gigantiops destructor</i>	DQ226041		Guyana
<i>Lasius</i> sp.	DQ226042		USA
<i>Lepisiota</i> sp.	DQ226043		South Africa
<i>Linepithema humile</i>	DQ226044		USA
<i>Melophorus</i> sp.	DQ226045	DQ226024	Australia
<i>Myrmecocystus mimicus</i>	DQ226046		USA
<i>Oecophylla smaragdina</i>	DQ226047		Malaysia
<i>Opithopsis respiciens</i>	DQ226048	DQ226025	Australia
<i>Paratrechina longicornis</i>	DQ226049		USA
<i>Petalomyrmex phylax</i>	DQ226050	DQ226026	Cameroon
<i>Plagiolepis alluaudi</i>	DQ226051		USA
<i>Pogonomyrmex</i> sp.	DQ226052	DQ226027	USA
<i>Polyrhachis</i> sp.	DQ226053	DQ226028	Equatorial Guinea
<i>Prenolepis imparis</i>	DQ226054	DQ226029	USA

Appendix 1

Alignment of Taxa

Secondary structural symbols are as follows: “()” represent hair-pin stem loops; “[]” represent simple stems; “[]” represent unalignable regions.

Large subunit rRNA		2 (1)	D2	D2-2a	2b	2c
<i>Camponotus chromoides</i>	[CAA]	????????????????????????????????????nnnGAGG (---agaUUAUCUGUACGCGCGUGGCUUCnGUGUGGCGGCAU/GUCCCGGGAG				
<i>Anolepis gracilipes</i>	[ANL]	?????????????????CCUUAAGAACCCAAAG-AUGGAAG (GGGAGUUAUCUGUACGCGCGUGGCUUCnGUGUGGCGGCAU/GUCCCGGGAG				
<i>Petalomyrmex phylax</i>	[PBT]	cgcuuacaggGUAUAACCUAAGAAACCCAAAG-AUGGAAG (GGGAGUUAUCUGUACGCGCGUGGCUUCnGUGUGGCGGCAU/GUCCCGGGAG				
<i>Formica subsericea</i>	[FOR]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Polyrhachis</i> sp.	[POR]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Otiphiopsis respiciens</i>	[PRE]	?? (ggagAUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Atropis arnoldi</i>	[ARN]	?? (ggagAUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Acropyga donisthorpei</i>	[ART]	?? (GnnnAUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Acropyga epedana</i>	[ARE]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Acropyga acutiventris</i>	[ARA]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Cladomyrmex mayriatae</i>	[CMB]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Brachymyrmex depilis</i>	[BDA]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Melophorus</i>	[MEL]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Paratrechina longicornis</i>	[PAL]	CGCUUACGGGGUAACCUAAGAAACCCAAAG-AUGGAAG (GGGAGUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Lasius</i> sp.	[LAS]	?? (GGGAGUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Gigantiopsis destructor</i>	[GITG]	cgcuuacaggguuaaaccuaAgAAACcAAAG-AUGGAAG (GGGAGUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Lepisiota</i> sp.	[LBP]	?? (777A-AuAUcAggUCA-CGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Plagiopsis alluaudi</i>	[PLA]	?????????????????ACCUAAGACACCCAAAG-AUGGAAG (gg-AuAUcAUcUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Myrmecocystus mimicus</i>	[MRC]	?? (????????????????????GCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Oecophylla smaragdina</i>	[OCB]	?? (GGAGA-UUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Dorymyrmex</i> sp.	[DOR]	?? (GGGAGUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Amblyopone pallipes</i>	[AMB]	CGCUUACGGGGUAACCUAAGAAACCCAAAG-AUGGAAG (GGGAGUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				
<i>Pogonomyrmex</i> sp.	[POG]	CGCUUACGGGGUAACCUAAGAAACCCAAAG-AACGAAG (GGGAGUUAUCUGUACGCGCGUGGCUUCGCGUGUGGCGGCAU/GUCCCGGGAG				

	2c'	2b'	D2-2a'	D2-3		
[CAA]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUnGmG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[ANL]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[PBT]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[FOR]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[POR]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[OPR]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[PRE]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[ARN]	-UCCUUGGUGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[ART]	-UCCUUGGUGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	sAC
[ARE]	-uccuueggg-cueg-cgccccgggacAGcGuaacccg-cgaugucggggcgucgugugugugacuu-ucuccc			caaguagaa	gugcgc	gac
[ARA]	-UcnUGUGUG-CUGG-CGCGGGGACGUGnGUGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	AGUGCG	Gms
[CMB]	??			caugcgc	gugcgc	gac
[BDA]	gucueggg-cueg-cgccccgggacgucacccg-cgacgucggggcgucgugugugugacuu-ucuccc			CAAGUAAAA	GUGCnn	nnC
[MEL]	-UCCUUGCGG-CUGG-CGCGGGGACGUGnGUGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[PAL]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[LAS]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[LBP]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[GITG]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[PLA]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[MRC]	-uccuungag-cueg-cgccccgggacgucacccg-cgaugucggggcgucgugugugug					
[OCB]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[DOR]	-UCCUUGCGG-CUGG-CGCGGGGACGUGACGUGGUGG-UcUAUGUGGnGUGUGUGGnGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[AMB]	---U-U-UcUGU-CGCGGGGACGUGACGUGGUGG-AGCAGUCCGAGC-UCCUGUGUGGACU-UCUCCG			CAAGUAGAA	GUGCG	GAC
[POG]	-CCCUUGCGG-CUGG-CGCGGGG-UACGCGCGCGUGG-CGGAUGUGGCGGUGUGGAGUGUGGACU-UCUCCG			CUAGUAGAA	GUGCG	GAC

	4a	4b	4c	4d	4d'	4c'	4b'	4a'
[CAA]	(CGCGUGGUGUGUGGUGUACGCGCGCGGUGGnGnGnGACUGUGGUG-UCCGCG-UUAUAAAG-GnCGCGGGAACAAACCCGUGGUGCGCGCGGCGUGUGGCGGCGG)							
[ANL]	(CGCGUGGUGUGUGGUGUACGCGCGCGGUGGUGUACUGUGUGUG							

	5a'		D2-3'					
[CAA]	GggGAGC	CG	AUACC	GtGtG	GCAGC	CGANU	GCUUU----	Neu
[ANL]	GCAGAC	CG	AUACC	GUUCG	GCAGC	CUACU	CGUUUGGUAU	
[PTT]	GAGGAGC	CG	AUACC	GUUCG	GCAGC	CUACU	GCUUUGGUAU	
[POR]	nnnnnAU	CU	AUACC	GUUCG	GCAGC	CUAC?	????????????	
[FOR]	GAGGAGC	CG	AUACC	GUUCG	GCAGC	CUACU	GCU???	???????
[OPR]	GGGUGAC	CG	AUACC	GUUCG	GCAGC	CUU??	??????????	
[GAG]	GGGAGC	CG	AUACC	GUUCG	GCAGC	CUU??	??????????	
[ARN]	GGAAGAC	CG	AUACC	GUUCU	GCAGC	CUU??	??????????	
[ART]	gga?????	??	?????	?????	??????	?????	??????????	
[ARE]	ggagagac	CG	?????	?????	??????	?????	??????????	
[ARA]	AAACG?	??	?????	?????	??????	?????	??????????	
[CMN]	cgagag?	??	?????	?????	??????	?????	??????????	
[BRA]	???????	??	?????	?????	??????	?????	??????????	
[GAG]	GGAGC	CG	AUACC	GUUCG	GCAGC	CGU	g-----	
[PAL]	GGAGC	CG	AUACC	GUUCG	GCAGC	CGU	g-----	
[LAS]	GGGAGC	CG	AUACC	GUUCG	GCAGC	CUACU	CGuugGg???	
[GIG]	GGGAGC	CG	AUACC	GUUCG	GCAGC	CUACU	GCUUUGGUAU	
[LEP]	GGGAGC	CG	AUACC	GUUCA	GCAGC?	?????	??????????	
[PLA]	GGGAGC	CG	AUACC	GUUCA	??????	?????	??????????	
[MRC]	GGGAGC	CG	AUACC	GUUCG	GCAGC	CUACU	??????????	
[OCG]	AGGAGC	CG	AUACC	GUUCG	GCAGC	CUAan	??????????	
[GAG]	GGGAGC	CG	AUACC	GUUCG	GCAGC	CUU??	??????????	
[AMB]	?????????	??	?????	?????	??????	?????	??????????	
[POG]	----GAC	CG	AUACC	GUUCG	GCAGC	CUUUGGUAU	?	

Table 2

Morphological character matrix. Numbers of characters and character states correspond to those found in the Material and Methods.

	000000000111111 123456789012345
<i>Acropyga acutiventris</i>	111011100100011
<i>Acropyga arnoldi</i>	11101(01)100100011
<i>Acropyga donisthorpei</i>	111011100100011
<i>Acropyga epedana</i>	111011100100011
<i>Anoplolepis gracilipes</i>	100100000111001
<i>Brachymyrmex depilis</i>	100001000100111
<i>Camponotus chromaiodes</i>	000100211011101
<i>Cladomyrma maryatae</i>	100100000200111
<i>Dorymyrmex</i> sp.	00000000010000
<i>Formica subsericea</i>	100100201011101
<i>Gigantiops destructor</i>	000100200011101
<i>Lasius</i> sp.	000110000100111
<i>Lepisiota</i> sp.	100101000102011
<i>Linepithema humile</i>	01000000010000
<i>Melophorus</i> sp.	000101201211001
<i>Myrmecocystus mimicus</i>	000100200100101
<i>Oecophylla smaragdina</i>	000110210011101
<i>Opithopsis respiciens</i>	000101201011101
<i>Paratrechina longicornis</i>	000101000102111
<i>Petalomyrmex phylax</i>	100110100200?11
<i>Plagiolepis alluaudi</i>	100000000102011
<i>Polyrhachis</i> sp.	000100211011101
<i>Prenolepis imparis</i>	000100000102111

Table 3

EF 1-alpha primer sequences.

M3	5' CAT ATW AAC ATT GTS GTS ATY GG 3'
TRS10R	5' ACG GCS ACK GTT TGW CKC ATG TC 3'
For1	5' GCA TCG ACA AGC GTA CCA TCG 3'
Rev1.1	5' CGT CTT ACC ATC GGC ATT GCC 3'
U377.1	5' TTG GCG TGA AGC AGC TGA TCG 3'
TRS1R	5' ACC TGG TTT YAA GAT RCC GGT 3'
Cho7.1	5' CTT CAG GAT GTC TAT AAG ATT GG 3'
U52.1	5' CCG CTT CAG GAT GTC TAT AA 3'
L53	5' CC GCG TCT CAG TTC YTT CAC 3'
TRS4F	5' GCG CCK GCG GCT CTC ACC ACC GAG G 3'
TRS9.1R	5' GGA AGG CCT CGA CGC ACA TMG G 3'