FOR THE RECORD

Extending the diffraction limit of protein crystals: The example of glutamine synthetase from *Salmonella typhimurium* in the presence of its cofactor ATP

SHWU HUEY LIAW, GYO JUN, AND DAVID EISENBERG

Molecular Biology Institute and Department of Chemistry and Biochemistry,
University of California, Los Angeles, California 90024-1570

(RECEIVED November 18, 1992; REVISED MANUSCRIPT RECEIVED December 21, 1992)

Glutamine synthetase (GS) plays a central role in cellular nitrogen metabolism in bacteria, catalyzing the ATP-dependent condensation of ammonia with glutamate to yield glutamine in the presence of divalent cations. GS is regulated by product feedback, by oxidative modification, and by adenylation (Rhee et al., 1989). The 3.5-Å structure of completely unadenylylated GS was determined by Almasy et al. (1986) and refined by Yamashita et al. (1989). To improve the atomic model, and our understanding of the regulation and reaction mechanism, it has been necessary to grow GS crystals that diffract to higher resolution. We find that the cofactor ATP affects the size, growth rate, and diffraction quality of GS crystals.

Two crystal forms grown in different concentrations of ATP or the reaction products ADP and Pi are found to have distinct morphologies, stabilities, space groups, and resolution limits. Form XVI crystals, grown at high ATP concentration (greater than 500 μ M), grow to greater than 1 mm, belong to space group C22₂, and have unit-cell constants $a = 235.1$ Å, $b = 138.7$ Å, and $c = 205.6$ Å. They diffract to no better than 3.2 Å resolution and have a short lifetime in the X-ray beam. These crystals tend to redissolve.

Form XIV crystals are grown at low ATP concentrations (less than 200 μ M). They are routinely $0.8 \times 0.6 \times 0.2$ mm and belong to space group C2, with unit-cell dimensions $a = 235.5$ Å, $b = 134.5$ Å, $c = 200.1$ Å, and $\beta = 102.8^\circ$. They diffract to 2.4 Å resolution, are suitable for data collection, and are stable for at least 6 months. Both Form XIV and XVI crystals have been observed earlier (Heidner et al., 1978; Janson et al., 1984), and Form XIV crystals have been used to determine the GS structure at 3.5 Å resolution. However, before we determined

the importance of ATP, crystals grew irregularly, were rarely large enough for X-ray diffraction studies, and had limited resolution. The present study reveals that the ATP concentration is a crucial variable in obtaining large, better diffracting crystals.

The two differences in the growth conditions between crystal Forms XIV and XVI are the concentrations of ATP and of spermine tetrahydrochloride. The ATP:GS monomer ratio is 0.5 in the best Form XIV crystals, and it is equal to or greater than 1 in Form XVI crystals.

Two lines of evidence suggest that high ATP concentration makes GS more soluble, accounting for the redissolving of Form XVI crystals. The first is that increased concentrations of positively charged spermine tetrahydrochloride stabilize the growth of GS containing high ATP. This is presumably because of charge compensation. The other line of evidence is that addition of 2 mM ATP to crystallization drops dissolves Form XIV crystals. High ATP may change the packing of GS molecules, either because of an additional ATP binding site on the GS monomer, or because of the added negative charges of ATP. Whether ATP actually alters the packing of GS molecules we will learn only when we determine the structure of Form XVI crystals.

It may be significant that the best crystals are obtained under the condition in which the ATP concentration is 6 times the GS dodecamer concentration. This suggests that half of the active sites are occupied by ATP in the best quality crystals. This could imply that glutamine synthetase shares some characteristics of a half-of-the-sites reactive protein (Fersht, 1977). In fact, the studies on the inactivation of rat liver GS by [³⁵S]- or [methyl-¹⁴C]-L-methionine-S-sulfoximine indicated that about four molecules of methionine sulfoximine phosphate were bound per molecule of totally inactivated GS, which has eight subunits. This result might imply that rat liver GS nor-

Reprint requests to: David Eisenberg, Molecular Biology Institute, University of California, Los Angeles, California 90024-1570.

mally uses only four subunits for catalysis; the remaining four subunits may function in regulation (Tate & Meister, 1971). However, "half-of-the-sites" behavior was not observed in the inactivation of sheep brain GS or *Escherichia coli* GS (Ronzio et al., 1969; Weisbrod & Meister, 1973). Also, studies by equilibrium dialysis (Ginsburg, 1969) show that *E. coli* dodecamer GS binds 12 molecules of two feedback inhibitors, AMP and tryptophan. Nevertheless, the question "Does dodecamer *S. typhimurium* GS have some half-of-the-sites character?" may deserve further investigation.

References

- Almassy, R.J., Janson, C.A., Hamlin, R., Xuong, N.-H., & Eisenberg, D. (1986). Novel subunit-subunit interactions in the structure of glutamine synthetase. *Nature* 323, 304-309.
- Fersht, A. (1977). *Enzyme Structure and Mechanism*, p. 216. W.H. Freeman and Company, San Francisco.
- Ginsburg, A. (1969). Conformational changes in glutamine synthetase from *Escherichia coli*. II. Some characteristics of the equilibrium binding of feedback inhibitors to the enzyme. *Biochemistry* 8, 1726-1740.
- Heidner, E.G., Frey, T.G., Held, J., Weissman, L.J., Fenna, R.E., Lei, M., Harel, M., Kabsch, H., Sweet, R.M., & Eisenberg, D. (1978). New crystal forms of glutamine synthetase and implications for the molecular structure. *J. Mol. Biol.* 122, 163-173.
- Janson, C.A., Almassy, R.T., Westbrook, E.M., & Eisenberg, D. (1984). Isolation and crystallization of unadenylylated glutamine synthetase from *Salmonella typhimurium*. *Arch. Biochem. Biophys.* 228, 512-518.
- Rhee, S.G., Chock, P.B., & Stadtman, E.R. (1989). Regulation of *Escherichia coli* glutamine synthetase. *Adv. Enzymol. Related Areas Mol. Biol.* 62, 37-92.
- Ronzio, R.A., Rowe, W.B., & Meister, A. (1969). Studies on the mechanism of inhibition of glutamine synthetase by methionine sulfoximine. *Biochemistry* 8, 1066-1075.
- Tate, S.S. & Meister, A. (1971). Regulation of rat liver glutamine synthetase: Activation by α -ketoglutarate and inhibition by glycine, alanine and carbamyl phosphate. *Proc. Natl. Acad. Sci. USA* 68, 781-785.
- Weisbrod, R.E. & Meister, A. (1973). Studies on glutamine synthetase from *Escherichia coli*. *J. Biol. Chem.* 248, 3997-4002.
- Yamashita, M.M., Almassy, R.J., Janson, C.A., Cascio, D., & Eisenberg, D. (1989). Refined atomic model of glutamine synthetase at 3.5 Å resolution. *J. Biol. Chem.* 264, 17681-17690.