RECOLLECTIONS

Growing up in the golden era of intermediary metabolism

MORRIS E. FRIEDKIN

Department of Biology and the Center for Molecular Genetics,
University of California at San Diego, La Jolla, California 92093-0634

In this article I emphasize recollections of encounters with gifted scientists who impressed me with their ingenuity and humanity during the early years of my career. Growing up scientifically when the knowledge of intermediary metabolism flowered, my thinking and approach to research were profoundly influenced and molded by a milieu that generously provided the means for discovery. I hope this account of my experiences captures the atmosphere of an era in biochemistry that preceded the discoveries of molecular and cellular biology.

The extraordinary ability of great biochemists to determine the nature of individual steps of intermediary metabolism was surpassed by the insight that enabled them to elucidate the molecular logic of an entire pathway.¹ This imprimatur of genius was clearly revealed by the knack of scientists such as Warburg, Krebs, Szent-Györgyi, and Lipmann to recognize the sense of metabolic pathways.

Warburg admired a pathway that supplies its own hydrogen acceptor; in anaerobic glycolysis the hydrogen released during the oxidation of 3-phosphoglyceraldehyde is utilized downstream in the metabolic pathway to reduce either pyruvate or acetaldehyde. Krebs comprehended that the pathway of the tricarboxylic acid cycle was an unusual catalytic process; i.e., oxaloacetate

in a sequence of steps regenerated oxaloacetate, thus providing a way of oxidizing many molecules of pyruvate. The Krebs urea cycle employs an analogous cyclic process for removing many molecules of ammonia. Szent-Györgyi fathomed the implications of the catalytic effects of dicarboxylic acid metabolism on the oxidation of carbohydrates. Lipmann illuminated the relationship between energy-yielding reactions coupled to the biosynthesis of ATP and the subsequent use of energy-rich ATP for biological syntheses and function

Iowa State College in Ames, 1938–1941

In the late 1930s an undergraduate at Iowa State College could gain entree into a variety of research projects and earn a few dollars as well. Beans harvested by hand from experimental soybean plots needed to be packaged and tagged. Although a day in the fresh air of an Iowa farm could be invigorating, a more exciting place was the Agricultural By-Products Laboratory, a facility designed during the Great Depression to develop new uses for crops.

At the Ames laboratory, particulate masses of *Aspergillus niger* were grown in a rotary aluminum fermentor with a capacity of 140 gallons. Gluconic acid, as soon as it was formed from glucose by the mold, was neutralized with calcium carbonate, yielding calcium gluconate; furthermore, the mold could be recovered by filtration and recycled to ferment many batches of medium. This provided an economical source of gluconic acid, in contrast to an expensive chemical method employed previously. The procedures developed at Ames were utilized thereafter in the commercial production of millions of pounds of gluconic acid. A few years later I would again encounter a similar submerged culture technique for growing molds of the *Penicillium notatum-chrysogenum* group in the large scale production of penicillin.

During the period 1940–1941, graduate work under the guidance of Harvey Diehl, an inorganic chemist, led to a Master's degree in analytical chemistry. I studied the performance of an instrument designed to separate metals (such as tin and copper) by graded cathode potential control of electrodeposition from aqueous solutions of their respective salts. Although I understood the electrochemical concepts underlying the separation process, I was not stimulated by the project.

Reprint requests to: Morris E. Friedkin, Center for Molecular Genetics, 0634, University of California at San Diego, 9600 Gilman Drive, La Jolla, California 92093-0634; e-mail: mfriedkin@ucsd.edu.

Morris E. Friedkin is Professor Emeritus of Biology at the University of California, San Diego, where he taught biochemistry to medical students as well as undergraduates from 1969 to 1989. After receiving his Ph.D. from the University of Chicago in 1948, he was a postdoctoral fellow at the University of Copenhagen in Herman Kalckar's laboratory. In 1949–1958, Dr. Friedkin served on the faculty of the Washington University School of Medicine and then became chairman of the Department of Pharmacology and later headed the combined Departments of Biochemistry and Pharmacology at Tufts University School of Medicine at Boston. After moving to California he became Provost of Revelle College from 1974 to 1976. Dr. Friedkin is a member of the National Academy of Sciences and a fellow of the American Academy of Arts and Sciences.

¹“Molecular logic” was a term coined by Lehninger to indicate the coherence of a metabolic pathway consisting of sequential reactions that provide for channeling along specific routes and products. This provides the means of maintaining dynamic relationships between structure and function of biomolecules as well as the underlay of energetics that makes life possible.

Harvey Diehl and Clifford C. Hach's experiments on organic cobalt compounds, however, aroused my curiosity. By bringing together cobalt acetate, ethylene diamine, and salicylaldehyde, a cinnamon-colored oxygen-carrying compound was formed. My participation in this project was to help prepare large quantities of the organic cobalt complex named Co-Ox. The reversible absorption and release of oxygen from Co-Ox made possible the isolation of pure oxygen from the atmosphere. Clifford Hach's parents allowed us to carry out pilot runs for the manufacture of oxygen at their farm not far from the campus. When exposed to air, Co-Ox absorbed oxygen at room temperature and the residual nitrogen was removed; then upon heating, oxygen was released in pure form. The process could be recycled by removing the oxygen, cooling Co-Ox, and exposing it again to the atmosphere.

The process appeared to have a bright future. After I left Ames this belief was reinforced when I was informed by the government that the Co-Ox experiments had been classified secret and that I was not to divulge the nature of Co-Ox's properties to anyone. I speculated that Co-Ox was being tested for use on submarines to replenish the oxygen stores. After the end of the war, in 1948 details of the unusual properties of Co-Ox were published in a series of papers. Then I learned about the demise of the Co-Ox project (Hach & Diehl, 1948). After several cycles, Co-Ox, either in the oxygenated or deoxygenated state, when suspended in an atmosphere of oxygen could cause violent dust explosions!

The impact of teachers at Iowa State College

Henry Gilman, a world authority on metallo-organic compounds, published 1,000 papers over a span of 59 years. He taught elementary organic chemistry in a manner that terrified his students. Most of his lectures consisted of question-answer inquisitions. However, on a one-to-one basis as my faculty advisor he was most helpful and pleasant.

Frank H Spedding taught a course in physical chemistry. His eyes seemed to dart about constantly. His students attributed this to the long hours he spent in a darkroom, where he carried out spectroscopic research related to his studies of rare earth elements. During the war years he shuttled between Chicago and Ames. He and his colleagues had developed a method of producing pure uranium metal. When the reactor constructed under the stands of Stagg Field at the University of Chicago became sustaining on December 2, 1942, Spedding was present for good reason. His group at Ames had supplied the largest portion of pure uranium metal used in assembling the pile. In his lectures to us before the war he had described the fission of uranium and emphasized the inherent difficulties in achieving the successful use of atomic energy.

Northern Regional Research Laboratory in Peoria, Illinois, 1941–1945

On a Sunday afternoon a day before I started my new job at the Northern Regional Research Laboratory, I was listening to a radio broadcast of the New York Philharmonic Orchestra, when suddenly the program was interrupted by an announcement that Pearl Harbor had been bombed by the Japanese. The next day, December 8, 1941, the United States declared war. I entered a laboratory that overnight had switched to a war footing.

I was assigned initially to two projects in the Fermentation Division. One involved a fermentation that yielded itaconic acid, a precursor of useful plastics. The second explored the conversion of carbohydrates to 2,3-butanediol, a precursor of butadiene. With the capture of all natural sources of rubber by the Japanese, the production of synthetic butadiene rubber became a high priority. It soon became apparent that the production of butadiene from ethyl alcohol and later from petroleum would be more efficient. The fermentation of wheat became the source of most of the ethyl alcohol used to produce butadiene for synthetic rubber (Ward, 1970).

Unaware that the Fermentation Division was poised to expand its research on the production of penicillin and that 8 days after the United States entered the war Robert D. Coghill, the head of the group, had met with four major drug companies to encourage them to participate in the penicillin project, I was switched from the butanediol effort to a division supervised by Dr. Frank Stodola. The research was part of a national mission: to make penicillin available to all of our forces and by secrecy to prevent its use by Germany and Japan. Within a year, 17 drug companies were working on penicillin. The military goal was achieved. By D-Day, June 6, 1944, when the Allies began the invasion of Normandy, penicillin was available in large amounts. The humanitarian aspects had been downgraded throughout the war. The Germans and the Japanese were dependent on sulfa drugs, which were not as effective as penicillin, and even in the United States the diversion of penicillin for civilian use was forbidden.

To explain why the Peoria laboratory launched a major endeavor to produce penicillin and to convey the sense of urgency that characterized this effort, certain events, prior to and after the outbreak of war, are worth recalling.

A crucial experiment was a protection experiment carried out by Florey and Heatley in Oxford on May 25, 1940. Seven weeks of work had yielded enough penicillin (only 1% pure) to carry out an experiment with eight mice inoculated with streptococci. Four infected animals were injected with penicillin; four served as controls. Heatley kept the death watch and wrote in his diary "The controls were looking very sick but the treated animals seemed very well. I stayed at the lab until 3:45 (a.m.) by which time all four control animals were dead. It really looks as if P. may be of practical importance" (Moberg, 1991).

The procedure Heatley had used to obtain the historic penicillin preparation involved extraction of penicillin from acidified mold cultures into ether. The back extraction from ether with slightly alkaline water yielded upon freeze-drying a crude sodium salt of penicillin. During the years 1942–1945 in Peoria, I used the extraction procedure many times, not knowing that Heatley had devised this simple method.

Among many types of containers used to grow *Penicillium* in surface culture, Heatley found that an enamel bedpan with a side arm through which cultures could be inoculated was most expedient. On Christmas Day 1940, Heatley inoculated 174 bedpan-type containers of culture with the mold. It was a cumbersome and time-consuming procedure (Moberg, 1991).

Howard Florey and Norman Heatley came to the United States in July of 1941 seeking help in the mass production of penicillin. The structure of penicillin was unknown, the yield of penicillin in surface culture was quite low, and the isolation procedures were primitive. The English scientists were directed to the Northern Regional Research Laboratory in Peoria, where

a general program in fermentation and the growth of molds was in progress. They described their attempts to prepare large quantities of penicillin by surface culture of Fleming's *Penicillium* mold, identified by Charles Thom, a world authority on fungi, as *P. notatum* (Stodola, 1968). Florey returned to England and Heatley stayed on for a few months to work with Andrew Moyer, whose forte was the nutrition of molds. Heatley had developed a simple biological assay for penicillin at Oxford. Thereafter, the amount of penicillin produced by mold in culture was expressed as Oxford Units. In order to study the effects of various nutritional factors, Moyer designed his experiments in checkerboard arrays, varying conditions of growth and additions of different substances. Although not very creative in concept, it was a surefire way of collecting data that might provide interesting leads. One of the nutrients Moyer added to the mold cultures was corn steep water (referred to initially as "corn steeping liquor" (non alcoholic!), a by-product of the wet corn milling process. The results were most encouraging, yielding marked increases in penicillin formation. Lactose also improved the *Penicillium* growth.

Working for Dr. Frank Stodola, an adept organic chemist, was an experience of inestimable value. For the first time I saw clearly that an independent career in research depended on having a doctorate degree.

The role of serendipity in Stodola's research is revealed in the following account. Stodola wished to prepare a crystalline derivative of penicillin as the benzylammonium salt. He added benzylamine to the free acid of penicillin in diethyl ether. An oil formed, which he set aside with some disappointment. At the end of the day, Stodola would wash his hands at the sink where he also placed discarded preparations. He happened to glance at the tube containing the oiled-out benzylamine-treated penicillin and noted with some surprise that a mass of crystals had formed after some hours had elapsed. This was a fortuitous observation that led to the discovery that the penicillin prepared in Peoria was different from Fleming's penicillin. The benzylamine had reacted with the beta lactam structure of penicillin to form a crystalline benzylamide derivative. X-ray powder diffraction patterns of the derivative from Peoria penicillin were found to be different from those of Fleming penicillin. The corn steep water used to increase the production of penicillin in fermentation vats contained phenylethylamine, a decarboxylation product of phenylalanine, one of the many amino acids released during the steeping process used in a commercial process to obtain starch from corn. Fleming's penicillin contained an aliphatic side chain, whereas the American product had an aromatic moiety. The British penicillin was named penicillin F; the American penicillin, penicillin G. The mold probably converted phenylethylamine to phenylacetic acid, which then combined with 6-aminopenicillanic acid to form penicillin G. Many semisynthetic penicillins have been made in recent years by utilizing an amidase-catalyzed reaction.

On May 7, 1943, Robert Coghill, head of the Fermentation Division in Peoria, wrote to Florey: "looks as though we shall be making at least 1,000,000,000 units per week now — we're beginning to harvest the fruit you and Dr. Heatley planted two years ago" (Roeder, 1980). Several pharmaceutical companies were turning out massive quantities of penicillin by the deep fermentation process developed in Peoria. Years later, Coghill commented, "We had the problem of the century presented to us on a silver platter" (Lambert, 1980).

Members of the original penicillin team were honored in various ways. In 1948, Coghill received The Medal for Merit from the War Department for planning and directing the team effort that led to the discovery of high-yielding strains of penicillin molds as well as methods of recovery and purification of penicillin. In 1987, Moyer was inducted posthumously into the National Inventors Hall of Fame for his findings that corn steep water and lactose in submerged shaken cultures were key ingredients in the mass production of penicillin. Although Heatley had been involved in Moyer's early experiments, the process was patented in the United States but not in England. The British at first did not favor applying for a patent on penicillin, a natural product. According to Sir Edward Abraham, the commitment of the British Medical Research Council to an ideal "that the fruits of such work should be freely available to all" was a strongly held view that was changed by the end of the war (Abraham, 1985).

Postscript on penicillin

Whereas Fleming, Florey, and Chain had received the Nobel Prize for the discovery and clinical use of penicillin, Heatley was not recognized for his contributions until 50 years later. Heatley, at the age of 80, was awarded an honorary Doctorate of Medicine from Oxford, the only such award in 800 years of the university's existence (Moberg, 1991).

There are several ironies in the penicillin story. Florey wrote Coghill in December of 1942: "I am thoroughly disillusioned about all commercial firms now. We have told them all we know, but seem to get neither information nor materials in return. . . . Now that these firms see the possibilities of large profits in this material, they seem to be behaving like a lot of tigers" (Roeder, 1980). The humanitarian aspects of penicillin gave way to the cruel demands of modern warfare. The project files were not declassified until 1980. As late as February 1945, Coghill considered the possibility that publication of the penicillin research could give the enemy an advantage. By June 1945, penicillin production had risen to 647 billion units. At the low price of 6 dollars per million units the "tigers" referred to by Florey in 1942 were doing pretty well.

Moyer, later extolled as developing "the use of precursors to produce more effective penicillin" (quoted from a penicillin historical marker at the Northern Regional Research Laboratory), was not comfortable in the role of a team player. He tended to be secretive about his research, as shown by his reluctance to make his notebooks available to Robert Coghill, his chief. Moyer apparently did not choose to emulate Heatley, who had revealed all of his expertise freely and without hesitation to the Peoria scientists. Even the idea of submerged fermentation with molds was not an original concept, as shown by a US Department of Agriculture process patented by O.E. May and others at Arlington Farms, Virginia, in 1935. As mentioned earlier, the formation of gluconic acid from glucose by *Aspergillus niger* was carried out at Ames, Iowa, in the late thirties by submerged culture techniques. The production of penicillin in deep vat fermentors, as much as 12,000 gallons in capacity, was therefore not an entirely new approach.

The last irony was that despite an intensive program in which the Army Transport Command delivered hundreds of new strains of penicillin-producing molds from soils obtained all over the world, Kenneth Raper, who was in charge of isolating new

strains, found the highest-yielding culture from a moldy cantaloupe picked up in a fruit market in Peoria—a tale told on the Penicillin Historical Marker in which “the discovery of a mold strain more productive of penicillin” is cited. The Peoria isolate had important properties significant for successful commercial production of penicillin. Raper’s group found that many previous *Penicillium* isolates unfortunately produced less penicillin per cc in submerged culture than in surface culture. Merck had tried to use submerged culture to no avail. The Raper isolate from a moldy Peoria cantaloupe turned out to be an excellent penicillin producer in submerged culture. With further mutations induced by irradiation, the Peoria cantaloupe-derived strain became the workhorse of the penicillin industry.

Over the ensuing years I have often thought about the good luck of working on the penicillin program. Had I stayed in Ames, Iowa, I would have probably become involved in the Manhattan Project.

Graduate studies at the University of Chicago, 1945–1948

In an advanced course in biochemistry at the University of Chicago in 1946, Albert Lehninger gave lectures on Otto Warburg’s discovery of cytochrome oxidase in the respiratory chain and how Warburg had used the strategy of employing an action spectrum to determine the actual spectrum of the enzyme in intact cells. Not allowing factual material to obscure the big picture, Lehninger captured the broad scope of biochemistry. Some years later his famous textbook reflected the same kind of elegant thinking. Inspired by his lectures, I approached Lehninger and asked if he would consider supervising my doctoral studies. Fortunately, he agreed to become my mentor.

Lehninger’s emotional encounter with Otto Warburg is described in the boxed section entitled “Lehninger Meets Otto Warburg.”

In 1946, I was excited when I first understood the significance of Otto Warburg’s discovery of 1,3-diphosphoglycerate as an intermediate that conserves the energy of the oxidative step of anaerobic glycolysis. In a similar vein, Fritz Lipmann had theorized that in the Krebs tricarboxylic acid cycle a phosphorylated intermediate might be formed when malate is converted to oxaloacetate, i.e., phosphomalate. Upon oxidation, the postulated

phosphomalate intermediate would give rise to phosphoenol-oxaloacetate, an energy-rich compound. Lehninger suggested that phosphomalate should be synthesized and tested according to Lipmann’s scheme. It seemed like an approachable first project. Fortunately, the synthesis went smoothly. In the spring of 1947, Lehninger introduced me to Lipmann at a meeting of the Society of Biological Chemistry. I told Lipmann that I had synthesized his postulated compound. Lipmann’s reply dashed my enthusiasm. “Oh, I don’t believe that anymore!” I was surprised that he had discarded an idea so soon after publishing it. Lipmann was right. Phosphomalate turned out to be inert; it had no future.

During my graduate days with Lehninger, we found that water-treated liver mitochondria (we didn’t know in our earlier studies that the active component of our liver preparations was indeed the mitochondrion demonstrated by Eugene Kennedy’s research on fatty acid oxidation) would support oxidative phosphorylation, as shown by the incorporation of inorganic ^{32}P into ATP when NADH was present. We believed our results were due to respiratory-chain electron transport coupled to ATP formation.

In addition to our experiments with NADH and inorganic ^{32}P , Lehninger also demonstrated that when mitochondria oxidize 3-hydroxybutyrate with the concomitant reduction of intramitochondrial NAD to NADH, a coupled phosphorylation of ADP to ATP occurs. Although much information has accumulated supporting the existence of respiratory-chain phosphorylations starting with intramitochondrial NADH, our early inorganic ^{32}P experiments with added NADH were revisited in 1981, as described in the boxed section entitled “Lehninger’s response. . .”

Eugene Kennedy and I were in synchrony during the early years of our scientific careers. During World War II both of us were engaged in medical aspects of the war effort: Gene worked on the large scale production of discrete serum protein fractions, I on the production of penicillin. When the war ended, we both entered the graduate program in biochemistry at the University of Chicago. The campus still showed the vestiges of military activities. The gates to the historic nuclear reactor under the stands of Stagg Field were guarded by armed soldiers, and the Manhattan Project was still operating. President Hutchens, a mythic

Fig. 1. Albert Lehninger at the Federation Meetings in Atlantic City, circa 1948.

Lehninger Meets Otto Warburg

Lehninger's admiration for Otto Warburg became reciprocal, as evidenced by a letter Lehninger sent to me during my postdoctoral year in Copenhagen in 1948:

"—Had quite a thrill a month ago. Was invited down to Urbana to spend a day with Warburg and give a seminar (on the DPN-linked phosphorylation). He's quite a boy—starchy, one-track mind, no small talk, monocle, and all brain. I told him about the work in the morning. He was skeptical at first but after asking some questions thought we really had something. After he got interested in it he refused to talk about anything else. In the afternoon I gave the seminar—huge crowd. He made some very interesting comments—one of which was that due to some structural changes occurring during electron transport, ATP-ase might be reversed. This sounded very much like Szent-Györgyi and I was surprised to hear it from Warburg. He was also tickled about the similarity of mitochondria and the chloroplast granules he has been studying. In the evening he called me from his apartment to tell me that he thought this was one of the most significant developments of recent years in this racket. They all think he is a genius down here but he is very uncooperative about lecturing etc. The U of C [University of Chicago] tried to make arrangements for him to spend a year or two here but I'm afraid they are going to fall through."

figure who made a significant imprint on the unique curriculum of the University of Chicago, was often seen on campus.

Gene and I became close friends when he too was accepted by Lehninger to direct his research. Our mentor did not lord over

a laboratory empire. There were just four in the group: Lehninger, two graduate students, and a laboratory assistant. We had no difficulty making daily contact with our leader. Close by were several clinical units involved in research on cancer of

Lehninger's Response to Ernster's Interpretation of Our ^{32}P Results with NADH

Many years later, Ernster and colleagues attributed the NADH effect we had reported in 1949 to the reduction of cytochrome *c* via a cytochrome *c* reductase that was not a part of the respiratory chain (Ernster et al., 1981). Respiratory-chain phosphorylations require an intact mitochondrial inner membrane known to be impermeable to NADH. The malate shuttle is generally believed to provide a way of reducing intramitochondrial NAD. It is difficult in retrospect to understand how extramitochondrial NADH alone without added malate could support respiratory-chain phosphorylations. If Ernster's interpretation was valid, i.e., electrons from extramitochondrial NADH were not entering the respiratory chain at Site I but rather at Site III, then we had demonstrated for the first time phosphorylations coupled to electron transport via terminal oxidation–reduction events in which reduced cytochrome *c* and oxygen participate.

Writing to Lehninger in 1981, I mentioned Ernster's view. Lehninger replied:

"Yes, I have seen the article by Ernster. The occurrence of a cytochrome reductase in the outer membrane has been known for a very long time and was even suspected in the early 1950s. I think Ernster's conclusion may well apply to some of the experiments we did with the ^{32}P method. However I don't think he used the same conditions I used in later experiments showing that water-treated mitochondria would oxidize NADH with net oxidative phosphorylation up to a P/O ratio of 2. The function of the outer membrane cytochrome reductase is not entirely clear but it is known that cytochrome *c* can shuttle back and forth between the outer and inner membranes, within the inter-membrane space."

Lehninger's comments about a P/O ratio of "up to 2" were significant because the NADH–cytochrome reductase–cytochrome *c*–oxygen pathway emphasized by Ernster should not yield a P/O ratio above 1.

Fig. 2. Adelaide and Gene Kennedy, our dear friends for half a century.

the prostate. Charles Huggins, a surgeon who in 1966 received a Nobel Prize for his research on hormonal treatment of prostatic cancer, had a group of young clinical researchers, including Paul Talalay and Elwood Jensen. Overheard one day in an elevator was a conversation between an elegantly attired surgeon and Huggins, who at that time spent most of his time at the laboratory bench: "Charles, you sure have a chemical air these days," this addressed to Huggins, who not in his Sunday best was informally clothed for working with dogs.

Gene and I both worked with insoluble particles that were easily centrifuged and capable of oxidizing fatty acids as well as sustaining oxidative phosphorylation. Lehninger was intrigued by the relationship between the Krebs tricarboxylic acid cycle, fatty acid oxidation, and ketone body formation. Except for a DU Beckman spectrophotometer, our equipment was simple but ad-

equate: an Elvehjem-Potter pestle homogenizer, a clinical centrifuge, Warburg flasks to measure microliter uptakes of oxygen, a colorimeter, and a pH meter.

Gene focused his efforts on fatty acid oxidation, I on oxidative phosphorylation. Gene and Lehninger believed that fatty acid oxidation was due to mitochondria because of the osmotic behavior of the insoluble particle preparations from liver. They isolated mitochondria using the hypertonic sucrose method developed at Rockefeller Institute and showed for the first time that essentially all the measurable fatty acid oxidation activity of the liver is due to mitochondria and that the integrated reactions of the Krebs tricarboxylic acid cycle as well as the esterification of inorganic phosphate can be attributed also to the mitochondria.

Earl A. Evans, Jr., our departmental chairman, was a bachelor who wore well-made tailored suits. He had become chairman in 1942 at age 32. His luxurious apartment in the near Northside of Chicago had a living room two stories high with a balcony overlooking two back-to-back grand pianos, as well as rare paintings. Once a year, wide-eyed graduate students were invited to a party in these sumptuous surroundings. The message: someday we too could live that way once we obtained our Ph.D.'s in biochemistry.

In preparation for his lectures, Evans stopped off at the John Crerar Library, an excellent private noncirculating library in downtown Chicago, where he read original articles on anaerobic glycolysis. His accounts of intermediary metabolism, with emphasis on the contributions of Embden, Meyerhof, and Warburg, captured the exuberant thrust of biochemical research during the prewar years in Europe.

During a typical day in the UC graduate student laboratory, political discussions held sway between Lloyd Kozloff, Irving Zabin, Gene Kennedy, Robert Johnston, Eugene Goldwasser, Eric Conn, Ray Koppelman, Miriam Banks, and me; night work was imperative to make up the hours lost in hot debate.

Quite often we would see Konrad Bloch, then a faculty member, with a tray of radioactive samples from experiments on cholesterol synthesis. He was headed for the counting room adjacent to the graduate student laboratory. We didn't know that he was also headed for a Nobel Prize.

Fig. 3. An outing near the University of Chicago, circa 1947. From left to right: Jan Lehninger, Frank Putnam, Albert Lehninger, Konrad Bloch, and Earl A. Evans Jr.

I liked go to one of the lecture halls to hear Enrico Fermi discuss elementary aspects of physics. He demonstrated basic physical laws with verve and ebullience and with an Italian passion that left a deep impression. The undergraduates sat entranced listening to a brilliant physicist who had directed the construction of the world's first self-sustaining chain reacting pile just a block away on the floor of a squash court under the West Stands of Stagg Field; yet here he was as excited about making basic concepts of classical physics clear to his students as he was about atomic physics.

Gene and I had common interests beyond the activity of mitochondria: Fred Hoyle's steady-state concept of cosmology in 1948 occupied our conversations, especially over beer and lobster at Hackney's, a popular seafood restaurant in Atlantic City, where the Federation Meetings were held. Atlantic City during the pre-Taj Mahal casino era was a great place for biochemistry meetings. Most of the important figures in intermediary biochemistry could be gathered in one meeting room. For graduate students it was a thrill to see Fritz Lipmann, Severo Ochoa, Carl and Gerty Cori, Otto Meyerhof, Erwin Chargaff, David Shemin, Harland Wood, and others. It was also a place to become acquainted with young biochemists such as Nate Kaplan and Sidney Colowick.

For relaxation and escape from the heat of midsummer Chicago, we walked to the shores of Lake Michigan, where we could plunge off the rocks into clear water not yet polluted. Playing chess with Gene on the rocks was a wonderful pastime and remained a paradigm for a friendship that has endured for 50 years.

I did not read Gene's scientific odyssey, *Sailing to Byzantium*, until I had finished my recollections. Describing the atmosphere of biochemistry during his graduate days at the University of Chicago, Gene wrote: "It is not easy to convey the heady sense of excitement and adventure that caught up those entering biochemistry at that time. The feeling was that of an intellectual gold rush" (Kennedy, 1992). It is not surprising that Gene concluded his autobiography with a quotation from Yeats, the great Irish poet. Gene has always loved poetry and without text could hold forth spontaneously.

State of biochemical knowledge and methodology in the late 1940s

Although important gaps remained to be filled in the field of biochemistry, advances were coming fast.

1. The nature of the two-carbon unit that fed into the Krebs tricarboxylic acid cycle was yet to be elucidated. Although Lipmann had discovered coenzyme A, Feodor Lynen showed that acetate and coenzyme A combined to form acetylCoA and that this was the critical two-carbon unit at the biochemical intersection of carbohydrate and fatty acid metabolism. Lipmann's liver extracts lacked phosphate transacetylase, the enzyme that transfers the acetyl group of acetyl phosphate to CoA; otherwise, he would have probably discovered acetyl CoA.

2. Linus Pauling correctly surmised that complementary units must exist in the genes to explain the duplication of genetic material, but initially he guessed mistakenly that the hereditary stuff was protein.

3. Pauling's discovery of the alpha-helix structure in proteins paved the way for Watson and Crick. Because the alpha-helix

did not contain an integral number of amino acid residues, Pauling's model was not accepted at first, especially in England. After Perutz read Pauling's paper, he took a new X-ray diffraction picture and found a reflection that confirmed the alpha-helix structure.

4. In our advanced biochemistry lectures we were told about Avery's pneumococcal transforming factor, in which DNA appeared to be the hereditary material. This experiment was the prototype for the present-day era of gene transfer and gene therapy.

5. There were no biotechnology companies with controversies about patent priorities. ATP as well as other nucleotides were not commercially available. Radioisotopes and labeled compounds (with the exception of inorganic ^{32}P from Oak Ridge), growth factors, antibodies, and restriction enzymes could not be ordered from thick catalogs.

6. Biochemists could detect metabolic events by use of a spectrophotometric instrument that became generally available (see boxed section entitled "Saga of the DU Spectrophotometer").

Copenhagen, 1948–1949

Deciding to spend a postdoctoral year with Herman Kalckar in Copenhagen was a natural consequence of my thesis work on oxidative phosphorylation. Kalckar was one of the pioneers in this field, having demonstrated oxidative phosphorylation in respiring kidney cortex extracts in which malate or fumarate in the presence of inorganic phosphate gave rise to phospho-enol-pyruvate. In 1949, Kalckar published a landmark review on energetic coupling in biological syntheses. The conceptual foundations emphasized by Kalckar in his article had a profound impact on my thinking about biochemical events.

Kalckar's scientific adventures before my postdoctoral year with him in Copenhagen are described in the boxed section entitled "Kalckar and the Coris."

During my postdoctoral year in Copenhagen, I cycled everyday to the Institute of Medical Physiology at 28 Juliane Maries Vej. Herman Kalckar and his first wife Vips lived in an apartment above the laboratories. Herman's kindness to me was beyond expectation. Quite often at the end of the day we would sip some sherry in Herman's living quarters. There we discussed not only science but the arts and the humanities. Sometimes we would interrupt our research and go on strolls during which Kalckar told me about Nordic mythology.

Undoubtedly Kalckar's interest in the humanities was influenced by the intellectual environment provided by his parents. His mother read and spoke French and German as well as Danish. She was familiar with the writings of Flaubert, Proust, Goethe, Lessing, and Heine. Herman was proud of his father, a broker, who refused to earn money during World War I on stocks that in Herman's words involved "sunken ships and lost lives."

I remember two noteworthy lectures during my stay in Copenhagen. One was by Karl von Frisch, who had spent the war years studying the language of honeybees. A bee having discovered a new source of food returns to the hive, where, by means of body language, a figure eight waggle dance, it informs the rest of the bees the whereabouts of the food source. Frisch's talk in Copenhagen was the first given in German since the end of the war.

Saga of the DU Spectrophotometer

The penicillin project in Peoria during World War II utilized an early model of the Beckman DU spectrophotometer, a battery-operated instrument, for studies of the structure of penicillin. I remember the first spectrum of penicillin G consistent with the presence of a phenyl ring.

Later, in Albert Lehninger's laboratory, I used the instrument to measure reduced cytochrome *c*. One day Lehninger informed me that he had discovered a treasure-trove of Beckman DU spectrophotometers in one of the hospital rooms—left over from a stockpile of instruments used in the extensive malaria projects carried on during World War II.

In Copenhagen, Kalckar introduced me to differential spectrophotometry. In studies of purine metabolism involving the conversion of hypoxanthine to uric acid, absorbency changes at 293 nm were easily detected. In Kalckar's laboratory I made preparations of xanthine oxidase from fresh, warm, unpasteurized cow's milk. The laboratory group eagerly devoured the by-product: whipped cream. Food in Denmark was still rationed as an aftermath of World War II.

During my years at Washington University, I inherited a Beckman with an unusual history. The instrument had been shipped from New York City to St. Louis and stored in the basement of Robert Shank's home (Bob was chairman of the Department of Preventive Medicine at the School of Medicine). Unfortunately, Bob's basement was flooded during a rain storm. Presumably, the instrument was irreversibly damaged. Oliver Lowry persuaded Shank to let him try to repair it, and Ollie successfully restored the instrument. A plastic dial facing had lifted off during the immersion in water. Ollie etched wavelength values into a brass dial by hand. In later years, technicians from the Beckman Company could not understand the exquisite sensitivity and stability of that miraculous instrument risen like Lazarus from the dead. The new Beckman models were constantly breaking down; my Old Stalwart worked like a charm.

When I moved to the Tufts University School of Medicine, I bought a \$15,000 UV recording spectrophotometer to follow the action of dihydrofolate reductase by the change of absorbency of NADPH at 340 nm. I also followed the activity of thymidylate synthase based on the difference spectrum between 5,10-methylene-tetrahydrofolate and dihydrofolate. I remember well how I was asked to write a letter to the NIH to justify the need for such an expensive instrument. I recalled that Otto Warburg had used a spectrophotometer built in his own laboratory to follow changes in the ratio of NADH to NAD in his fundamental studies of anaerobic glycolysis. In my letter of justification I cited the decisive studies of Jesse C. Rabinowitz in which he used differential spectrophotometry in his research on folate metabolism. Fortunately, NIH awarded me the funds to buy the instrument.

Fig. 4. Herman Kalckar, whose kindness to me was beyond expectation.

Kalckar and the Coris

Kalckar's first encounter with Carl and Gerty Cori in St. Louis shows how even a young scientist can teach his elders a thing or two. As described by Kalckar:

"I arrived in the bleak and sooty city of St. Louis in February 1939, and was well received by the Coris. Unexpectedly they were actively interested in my articles on oxidative phosphorylation but were unable to repeat the experiments. Gerty wanted me to make a few demonstrations, but Carl cut it short and simply asked a few questions, and so did I. I saw that the Coris used the old-fashioned Meyerhof extract technique, using test tubes. I explained to them that kidney cortex extracts show a very high oxygen consumption, and in order to see oxidative phosphorylation I always aerated the cortex extracts by shaking them in Warburg vessels.

A bright young researcher in the Cori department, Dr. Sidney Colowick, witnessed my discussion with the Coris. He asked to compare my technique with theirs. Colowick soon confirmed that the simple shaking technique was indeed needed in order to observe oxidative phosphorylation in cortex preparations. The Coris wrote to me with the good news and with their best thanks." (Kalckar, 1991).

Kalckar was convinced that the episode described above was an important factor that led to an invitation by the Coris to come to their lab when Herman's first-year fellowship was terminated. He returned to St. Louis in late February 1940, where, in his own words, he spent some of his happiest months in research.

The other lecture was by John Edsall, a pioneer in the fractionation of proteins, especially those present in human blood plasma. Although I cannot remember the topic of his talk, I do recall how impressed I was with his bearing, that of a noble man incapable of any artifice, a figure right out of the eighteenth century.

Asked to give a seminar on recent advances in the field of photosynthesis, I was honored by the presence of Niels Bohr, who in his usual fashion struck many matches to fire up a pipe that never seemed to remain lit. In my presentation, I took note of Martin Kamen's clever use of radioisotopes to probe the mechanism of photosynthesis; Kamen had shown that light photons caused the release of oxygen from water during the photosynthetic process. Being questioned about the surface area of a leaf undergoing photosynthesis, I could not adequately answer the query, but Niels Bohr immediately entered the fray, assuring the audience that everything was OK. He asserted that the efficiency of photosynthesis was such that enough light photons were captured per square centimeter of leaf surface to explain the rate of carbon dioxide utilization and the amount of oxygen released. How Bohr knew that was beyond me. He seemed very comfortable with his explanation. After reading Pais's biography of Niels Bohr, I finally understood why Bohr had attended my seminar on Kamen's use of radioisotopes in photosynthesis. In 1989, Hilde Levi (who had collaborated with Hevesy in Bohr's Institute on the application of radioisotopes in biology) was asked by Pais how Bohr responded to biological work with tracers. She replied, "He was exceedingly fascinated, electrified" (Pais, 1991).

One evening Kalckar presented a lecture at a meeting of the Royal Danish Academy. He invited me to come along as his

guest. The talks were held in a magnificent room with a huge painting of illustrious Danish scientists of the past. Most of the presentations were given in Danish. Soon it became apparent that the listeners were nodding off. Suddenly, during one of the talks the audience became unusually alert. Kalckar whispered to me that the topic being discussed was pornography in Greek art. Later, we moved to an adjacent area surrounded by large casks of Carlsberg beer. Beer and Danish-style smorgasbord were served by attractive young ladies. Those scholars who had nodded off earlier in the evening were now quite wide awake.

In Copenhagen we discovered that the phosphorolysis of guanine deoxynucleoside with deoxynucleoside phosphorylase yielded deoxyribose-1-phosphate, a highly labile phosphate ester. The favored reverse reaction was used in the biosynthesis of deoxynucleosides analogous to the Cori synthesis of glycogen with glucose-1-phosphate.

St. Louis, 1949–1959

After my postdoctoral year in Copenhagen, I moved to St. Louis as an instructor in Oliver Lowry's Department of Pharmacology. When I arrived in 1949, the Medical School at Washington University had already hatched four Nobel Laureates: Joseph Erlanger, Herbert S Gasser, Carl Cori, and Gerty Cori. The university with its abiding devotion to basic research served as a way station for many gifted investigators. Louis F. Leloir, Christian R. de Duve, and Edwin G. Krebs had come and gone, later to receive Nobel Prizes. The regal scientific roster would be further enlarged in subsequent years with Nobel Prizes to Arthur Kornberg, Severo Ochoa, Earl Sutherland, Alfred Hershey, Paul Berg, Rita Levi-Montalcini, Stanley Cohen, and Dan

Nathans—all with careers that included stays at Washington University.

Oliver Lowry had replaced Carl Cori as chairman of the Department of Pharmacology in 1947. Cori, who had other attractive offers, elected to remain in St. Louis as chairman of the Department of Biological Chemistry with commitments from the School of Medicine to provide more space for basic research.

I was happy to be at a center of learning where research and scholarship were valued and where the faculty was top-notch. In the 1950s, I learned enough pharmacology to teach medical students. Lowry felt I should have more knowledge of the clinical use of drugs by joining him in Barnes Hospital Rounds headed by the eminent physicians Carl Moore and W. Barry Woods, Jr. I also attended two unique seminars, one weekly and one daily. The weekly seminar was held on Friday afternoons and was presided over by Gerty and Carl Cori in the Department of Biochemistry at the Medical School. The topics covered in the Friday seminars were not the basis of attendance. To be there was the thing to do! Although the room was quite crowded, Carl Cori refused to move the seminar to a larger conference room; he preferred the intimacy of a small meeting place. Faculty from three Medical School departments (Biochemistry, Pharmacology, and Microbiology) as well as Bill Daughaday from Medicine attended regularly. The daily seminars were held at lunch and covered recent journal articles. These sessions were presided over by Arthur Kornberg in the Department of Microbiology.

At first, Carl Cori appeared to be unapproachable. That my perception of him was erroneous was made clear to me as the result of a memorable encounter during a summer vacation. I mentioned to him that my family would be camping at Bar Harbor, Maine. Cori urged me to call him, since he and Gerty would be renting a house at Bar Harbor. After we arrived in Maine,

we set up our tent on a beautiful campsite. After about a week, I ventured with some trepidation to call the telephone number Cori had given me. "Where have you been? Where are you camping? I will be right over." He astonished me with his friendliness. We rushed back to the campsite, frantically cleaning up and putting things in order. Shortly, Dr. Cori found our tent and insisted that he be allowed to inspect it inside and out. On hands and knees he crawled into our tent, came out, and nodded approval. From that moment on, my attitude toward Cori changed to one of deep devotion. Not only was he a great scientist but a wonderful human being. For years thereafter, we had many enjoyable conversations. He had become my scientific father figure and a dear friend. On several occasions over the years he shared with me his broad interests in the arts and the humanities.

In 1976, Cori published an article in which he recounted an episode that had occurred in June 1958 and that reflected his sensitive nature:

"I had climbed this mountain [the Faulhorn in Switzerland] as a sort of scientific pilgrimage on previous occasions and have climbed it since, so there was no way of holding me. The going was hard through the deep snow, but I got up without difficulty and finding myself alone in a sheltered spot, I took a little nap. When I woke up, the sun was low and the mountain completely deserted. . . . I then had a most exhilarating experience watching the mountain marmots cavorting in the snow. They must have been just up from their hibernation and they were the most playful animals imaginable. They gave me a real feeling of *joie de vivre*" (Cori, 1976).

My experiments continued to have a phosphate motif. One of my early research goals was the enzymatic synthesis of [^{14}C]thymidine with deoxyribose-1-phosphate (the compound I had crystallized as the cyclohexylammonium salt in Denmark), [^{14}C]thymine, and nucleoside phosphorylase. I was intrigued by Peter Reichard's findings that [^{15}N]thymidine was uniquely incorporated into thymine and not cytosine of DNA; furthermore, it was not utilized for RNA synthesis. My concept was that the replication of DNA could be easily followed with thymidine labeled with a radioisotope such as ^{14}C and that this could open up an avenue to the mechanism of DNA biosynthesis.

Although my concept was right, the design of experiments was flawed. With [^{14}C]thymidine I could easily demonstrate incorporation into newly formed DNA of intact cells from bone marrow, thymus, chick chorioallantoic membranes, and onion rootlet meristems. I had no trouble isolating beautiful preparations of labeled DNA from bone marrow and thymus, yet despite many attempts I could not detect in vitro incorporation of [^{14}C]thymidine into DNA with cell-free extracts.

With hindsight, my failure could be explained easily. The specific activity of [^{14}C]thymidine was too low to detect incorporation into DNA of cell-free extracts of mammalian cells that divided every 12–24 hours. Kornberg had wisely chosen *Escherichia coli* cells, which doubled within 20–40 minutes. In my studies, I was off by a factor of about 25. Even with cell-free extracts of *E. coli*, Arthur found less than 20 counts per minute above background when [^{14}C]thymidine was used—but they were real counts—not contamination!

In other experiments with H.A. McQuade and Alice Atchison, we found that chromosomal breaks were produced when

Fig. 5. My family camping at Bar Harbor, Maine, in 1954. From left to right: my wife Bert, Noah, Sukey, Debbie, Morris, and Lucky, our Airedale.

onion root tips were immersed in solutions of [^{14}C]thymidine. Meristematic cells have high numbers of mitotic figures. We attributed the fragmentation to the incorporation of radioactive thymidine into DNA determined by chemical and autoradiographic means. In controls in which [^{14}C]thymine was used to irradiate the meristematic cells from extrachromosomal loci ([^{14}C]thymine enters the plant cells but is not incorporated into DNA), the amount of chromosomal damage was significantly lower.

I turned my attention to the biosynthesis of thymidylate in cell-free extracts of *E. coli*. [^{32}P]deoxycytidylate, isolated from the DNA of ^{32}P -labeled *E. coli*, was deaminated to [^{32}P]deoxyuridylate. In the presence of thymidylate kinase, tetrahydrofolate, and formaldehyde or serine as sources of one-carbon units, [^{32}P]deoxyuridylate was converted to thymidine triphosphate by extracts of *E. coli* in collaborative experiments with Arthur Kornberg.

Encouraged by Arthur, I spent the summer of 1956 in van Niel's course on microorganisms at the Hopkins Marine Station at Pacific Grove, California. Arthur had gone there before taking up his new position at Washington University as head of the Department of Microbiology. He extolled the course for its total immersion in classical bacteriology. On my first day in class, van Niel handed out a bouquet of flowers to each of the students. It was not, however, to celebrate the start of his course but to supply us with a source of wild yeast. Our first project was to isolate individual yeast cells from the flowers. Later, we isolated slime molds from deer dung as well as a host of other microorganisms from a variety of sources. After an hour or two at the bench we rearranged our seats and prepared to listen to several hours of lecture by van Niel, who with intensity and enthusiasm regaled us with fascinating tales about the early history of bacteriology, his facts buttressed by a stack of journals close at hand. David Shemin also took the course that summer. I had admired the way he had unraveled the pathway of porphyrin synthesis. His broad smile and enthusiastic love of life could not help but bring cheer to all who were fortunate to know him. Our interactions gave me much pleasure. Years later, W.D. McElroy told me that during his undergraduate days at Stanford he too had taken van Niel's course and had learned more biochemistry during one summer at Pacific Grove than was offered at the main campus in Palo Alto.

Brookhaven National Laboratory, summer 1957 . . . and many summers thereafter

Invited by W.L. Hughes (Pete) to spend part of the summer of 1957 at Brookhaven, located about midway up Long Island, I was happy to accept for several reasons. Pete had prepared [^3H]thymidine by catalytic exchange between thymidine and tritiated acetic acid. [^3H]thymidine had advantages over the [^{14}C]thymidine I had synthesized. The lower-energy beta-particles released by radioactive decay resulted in a smaller range in tissues, making possible autoradiographs with high definition, properties that were skillfully employed to demonstrate semi-conservative replication in mitotic cells of *Vicia* root tips (Taylor et al., 1957). Pete had an early model scintillation counter, making quantitation of tritium very convenient. Lastly, in contrast to St. Louis, beaches were close by on Long Island Sound as well as on the Atlantic Ocean shore.

The Medical Division at Brookhaven was housed in wooden barracks dating back to World War I. Long corridors connected the units in a labyrinthine fashion. The overall effect, evoking an earlier era at Camp Yaphank, was unexpectedly more pleasing than the modern building built later to house the expanding activities of the medical group.

The pattern was set. After moving to Tufts in Boston, I returned again to Brookhaven many times with my family. Pete's interests ranged from the use of exotic radioisotopes in medical research to astronomy and to politics. He and his wife Eleanor were gracious hosts at parties given in their lovely Bellport home close to the Atlantic shore. David Shemin also collaborated with Pete. The three of us had many lively conversations.

Another unusual aspect of summers at Brookhaven was the opportunity to meet the physicists, especially those involved in particle acceleration research. Lee and Yang, who disproved the principle of conservation of parity, were some of the notable members of the colony of physicists. One summer, a close neighbor in the apartment complex was Leon Lederman. I enjoyed telling Lederman about the latest advances in molecular biology, while he staggered me with accounts of the intricacies of particle physics. Later, in 1988, Lederman won a Nobel Prize for his research on elementary particles. He also became the director of a huge facility, the Fermi National Accelerator Laboratory, in Batavia, Illinois. In 1994, the Fermi Lab announced the discovery of evidence for the top quark, a predicted and much sought-after subunit of matter.

In autoradiographic studies of tritiated thymidine uptake into the DNA of mouse tissues, Pete and his associates demonstrated cellular migration in the gastrointestinal epithelium from the crypts to the tips of villi. Experiments of this type showed the great utility of tritiated thymidine as a probe in a variety of physiological and pathological situations. Pete also used ^{131}I , a gamma-emitting isotope of iodine, to label iododeoxyuridine for in vivo studies of DNA synthesis. This provided a simple way of locating and evaluating regions of DNA synthesis in the intact mammal. Cell death revealed by loss of ^{131}I -labeled DNA could also be ascertained by this technique.

Pete alerted me to the unusual properties of carbon-11. Kamen had used this isotope in his early studies at Berkeley, but with his discovery of carbon-14, an isotope with a half-life of over 5,000 years, he discarded carbon-11 because of its very short half-life of 20 minutes. Syntheses and metabolic studies with carbon-11 were too hectic. Pete was very knowledgeable about the decay products of various radioisotopes. In contrast to carbon-14, an electron emitter, carbon-11 gives rise to a positron. Within a centimeter or two from the point of emission, positrons come to rest, and upon contact with electrons, matter and antimatter self-destruct. This annihilation gives rise to two powerful back-to-back gamma-rays that can be detected by an array of scintillation crystals designed to single out the reception of simultaneous photons 180° apart. A computer program provides the exact location of the initial emission of positrons. This technology (an in vivo imaging method), now called positron emission tomography (PET), has many applications. Carbon-11, fluorine-18, oxygen-15, and nitrogen-13—all positron emitters—have been incorporated into a large variety of organic molecules for probing biochemical transformations in humans and in animals.

The Brookhaven group headed by Alfred P. Wolf, a pioneer and a key player in the PET field, combined forces with Eliza-

both Crawford and me to prepare carbon-11-labeled thymidine by chemical and enzymatic means. The labeled deoxynucleoside was then used to study DNA synthesis in intact animals.

My interactions with Pete became more frequent when I induced him to move to Tufts University School of Medicine to become head of the Department of Physiology. Years later, both Pete and I moved from Boston, he back to Brookhaven and I to California, but we maintained our contacts with each other.

Tufts University School of Medicine in Boston, 1958–1969

When I moved to Tufts as chairman of the Department of Pharmacology, Alton Meister, head of the Department of Biochemistry, greeted me cordially and offered any help I needed. Al, true to his word, and experienced in the ways of the medical school, gave me good advice.

Al was indefatigable. After a full day at Tufts, he worked on the second edition of his book on amino acids between midnight and two in the morning. It was important to have an adequate supply of cigars to bolster his spirits and clarity of thought. He treasured three collections: good cigars, vials of organic compounds he had prepared during his career, and photographic equipment. During his many travels he usually carried several cameras to document the sights and experiences of every trip. He enjoyed developing his own film, including color as well as black and white. He photographed models of organic compounds, which he used in his numerous publications on amino acid metabolism. His lectures to medical students as well as his scientific talks were characterized by careful preparation and lucidity. I was sorry when Al Meister moved to Cornell Medical School, because I enjoyed our almost daily conversations. He became the first Jewish chairman of a medical department at Cornell. His father had been the first head of the Bronx High School of Science.

One of Al's colleagues, Gerhard Schmidt, had a background that reflected the turmoil in Europe during the 1930s. Although he was trained as a clinician, Gerhard opted for basic research in the Frankfurt laboratory of Gustav Embden. In 1928, he discovered that muscle deaminase distinguishes between adenine nucleotides that contain phosphate at positions 5', 3', and 2'. Only the 5'-nucleotide, formed from ATP during muscle contraction, could serve as a substrate. In 1933, he was forced to flee from Nazi Germany, moving from country to country until 7 years later he finally obtained a secure position at Tufts. With his direct link to European prewar biochemistry, Gerhard was an unforgettable figure at Federation Meetings, where he played the cello with his chamber music cronies. As a graduate student, I had used his excellent method for determination of DNA and RNA (published with Thannhauser).

Our two departments, Biochemistry and Pharmacology, meshed well, for good reason. When I came to Tufts I hoped that I could recreate the coherence of Oliver Lowry's department at Washington University. This meant that pharmacology at Tufts would have a biochemical flavor. Naturally, I recruited a group that was biochemically oriented. They included Louis Shuster, 1958; Walter McNutt, 1959; Roy Kisliuk, 1960; Norman Krinsky, 1960; Nico van Gelder, 1962; David Stollar, 1964; and Len Corman, 1965—all devoted to research and responsive to the needs of the students, the department, the medical school, and the dental school. Aside from biochemistry, the breadth of

interests in our group was strengthened by the expertise of van Gelder in neurophysiology, Stollar in immunology, and Corman in dentistry. My only requirement was that the faculty should learn enough pharmacology to teach it to the students. Clinical aspects of pharmacology were presented to the students in the form of seminars with invited clinicians. This arrangement worked quite well, since my colleagues were serious about their teaching responsibilities. Our form of biochemical pharmacology evolved to what is now known as molecular pharmacology.

Dean Joe Hayman ran the medical school, not with an iron hand but with a little black book he kept secure in a safe near his desk. Since his handwriting was concise and minuscule, his database could fit into his book. In response to a plea for more research funds, Hayman would open his book and shortly come up with a helpful suggestion. Perhaps \$5,000 would suffice? In a lower drawer of his desk he kept a bottle of whiskey for medicinal purposes. Both Joe Hayman and I had the gout. "Take a swig and put your feet up," he would say. "It always works for me and should do the same for you. I find the pain of gout is quickly relieved." Of course, a small dose of colchicine could also be helpful.

At Tufts, the motif of my research changed from phosphate to folic acid. Phosphate no longer played a significant role. The conversion of deoxyuridylate to thymidylate could be directly observed by spectrophotometric changes. The interplay between one-carbon units and tetrahydrofolate became my major focus.

The conversion of 5,10-methylene-tetrahydrofolate to dihydrofolate during thymidylate synthesis could be followed by changes in absorbency that were easily detected, thus providing a convenient assay for thymidylate synthase in experiments with Albert Wahba. Dihydrofolate, one of the products of the reaction, was isolated by chromatographic separation.

Tritium-labeled tetrahydrofolate, prepared by enzymatic reduction of dihydrofolate with tritium-labeled NADPH, was shown to be the source of hydrogen in the newly formed methyl group of thymidylate in experiments with Edward Pastore. The location of hydrogen in the reduced pyrazine ring of tetrahydrofolate that is specifically utilized in the synthesis of the methyl group of thymidylate could be pinpointed by the use of nuclear magnetic resonance in experiments with Edward Pastore and Oleg Jardetsky.

In collaborative experiments with a group headed by Abraham Goldin at the National Cancer Institute, marked increases of dihydrofolate reductase were detected in livers and spleens of leukemic mice that had become resistant to chemotherapy with various aminopterin derivatives. Computer models were constructed that related the development of resistance to the increase of dihydrofolate reductase.

Together with Laurence Plante and Elizabeth Crawford, attempts were made to modify folic acid so as to take advantage of increased dihydrofolate reductase activity in leukemic mice that have become resistant to antifolates. Analogues were screened by their capacity to be reduced to the tetrahydro form, which in turn exhibited the property of inhibiting thymidylate synthase.

Favorite lecturers

Over the years, four individuals stand out as my favorite lecturers: Cornelius B. van Niel, Enrico Fermi, Albert Lehninger,

and Richard Feynman. I have already described the impact of Lehninger, Fermi, and van Niel.

Richard Feynman, my favorite lecturer *in absentia*, based on television interviews as well as his published *Lectures on Physics*, always spoke with humor and wrote with an originality seldom equaled. He seemed incapable of making a trivial statement. His approach to any problem was always fresh and daring. He could articulate his thoughts without dressing them up with fancy words.

The four masters of communication left a deep impression on me. I tried to emulate these marvelous individuals, seeking to excite my students by describing the illuminations of insight and discovery that are the foundations of great science.

My unusual friend Samuel Bessman

Sam Bessman, chairman for many years of the Department of Pharmacology and Nutrition at the University of Southern California School of Medicine, has had a single-minded pursuit during most of his career: to elucidate the mechanism of action of insulin. His auto license plate spells out INSULIN.

Spurned by the diabetes establishment because of his maverick ideas, Sam, now retired, continues his unrelenting quest. In the thick of battle he is a formidable opponent who enjoys demolishing arguments that from his way of thinking reflect widely held but unfounded beliefs.

At this time, 1995, the mechanism of insulin is not completely understood. The initial binding of insulin to specific surface receptors activates transmembrane tyrosine-specific protein kinases. The trail ends there. Sam's ideas extend beyond to a biochemical arena that involves mitochondrial functions. For many years he has published articles in which he has concluded that the anabolic effects of insulin can be attributed to its effect on the association of hexokinase with mitochondria, i.e., insulin compartmentalizes hexokinase. Sam has also emphasized that an analogous compartmentalization occurs in muscle with creatine kinase bound to mitochondria. Thus, in both cases, hexokinase and creatine kinase can regulate mitochondrial functions by direct binding to the most important cellular organelle involved in oxidative metabolism.

In recent years, Sam has used specifically labeled intermediates in the Krebs tricarboxylic acid cycle to demonstrate that the TCA cycle is the ultimate target of insulin action. When I sent Sam's paper describing these effects to a well-known and highly qualified biochemist for review, the response was, "Although I question the conclusions, I can't see anything wrong with the data; therefore I recommend acceptance of the paper for publication."

In the absence of definitive experiments that elucidate the molecular mechanism of insulin action, I hope the diabetes establishment will not continue to dismiss Sam's ideas as unworthy of any consideration. If proved right by other investigators, Sam will receive the long-delayed recognition he deserves.

My special friend Nate Kaplan

The phosphate motif played an important role in Nate Kaplan's career, as it did in mine. As a graduate student at Berkeley, he studied phosphate utilization obtaining ^{32}P from the Lawrence cyclotron with the help of Martin Kamen, who became a life-

long friend. With Doudoroff and Hassid, he showed that glucose-1-phosphate was formed during the breakdown of sucrose.

Stimulated by Lipmann's monumental paper on the coupling of ATP formation and its use in biological syntheses, Nate joined Lipmann's group at Massachusetts General Hospital in 1945. There he worked on the identification of a heat-stable factor required for the acetylation of sulfanilamide. The factor, renamed coenzyme A, contained pantothenate as well as adenine, ribose, and phosphate. Lipmann's group was elated that most of the pantothenate in tissues was in the form of coenzyme A.

Nate moved to Johns Hopkins University, where, together with Sidney Colowick, he helped W.D. McElroy to establish a strong group of biochemists at the McCollum-Pratt Institute. With Colowick he organized important symposia, Phosphorus Metabolism I and II, and started the *Methods of Enzymology* series of volumes that would have a great influence on biochemical research for years to come. (The series in 1994 contained more than 240 volumes.)

Together with Martin Kamen, his old friend from Berkeley days, Nate founded a new department of graduate biochemistry at Brandeis, a small university barely 10 years old, located near Boston. He induced several former members of the Lipmann group (William Jencks, Mary Ellen Jones, Morris Soodak, and W. Farnsworth Loomis) to join him. It is not surprising that the graduate biochemistry program not only expanded rapidly but remained cohesive because of the spirit of cooperation he engendered as a benevolent and caring chairman.

Because of our similar interests, Nate and I became good friends. Initially our contacts were at yearly meetings in Atlantic City and Chicago. Later, when both of us were in Boston, the scientific and social aspects of our friendship brought us into even closer interactions.

Nate was like an uncut diamond which when cleaved reveals new facets of brilliance. He could wed chaos and order with his

Fig. 6. Nate Kaplan was like an uncut diamond which when cleaved reveals new facets of brilliance.

miraculous touch. Although he appeared to be unorganized, any decision involving planning and coordination of effort usually led to an effective outcome. Part of his effectiveness could be attributed to the wholehearted support of his colleagues, graduate students, postdoctoral fellows, and friends. His generosity was reciprocated by the loyalty of those he helped. An unusual example of reaching out to people was Nate's successful and unpublicized endeavor to help Frank Chytil, now a professor at Vanderbilt University, to escape with his family from communist Czechoslovakia.

Foremost was Nate's abiding love for biochemistry. As pointed out by W.D. McElroy, Nate's studies on the isozymes of lactic dehydrogenase in various species and tissues led to his interest in evolution as well as cancer research. Analogues of NAD (he preferred the classical DPN name) were also a subject of many articles. He and his students and colleagues published over 400 papers and books.

Attracted by a new medical school at the University of California in San Diego, Nate left Brandeis with hopes of bolstering the research base of the new clinical teaching entity. I followed him to La Jolla a year later, happy to be able to teach biochemistry to medical students and undergraduates as well.

It wasn't long before the departmental structure at UCSD caused him to regret his move from Boston. Embedded in the departments of Chemistry and Biology, biochemistry as a discrete graduate entity was nonexistent. The freedom of decision that prevailed at Brandeis was not possible at UCSD. The complexities of the academic structure were frustrating. When a controversy arose on campus involving W.D. McElroy, Nate was an outspoken advocate on behalf of his old friend. The intimacy of the Brandeis days could not be recaptured. Nate missed the respect he had received at Brandeis—after all, there were dozens of professors at UCSD who were equally powerful. However, for one moment in 1982, Nate had an uplifting experience, when at a festive Brandeis celebration he received an Honorary Degree, with a great outpouring of warmth and esteem.

Nate's enthusiasm for biochemistry never waned despite his discomfort at UCSD. He utilized the most modern techniques in his research. He was daring in his approach to science. He

Fig. 7. My brother David Friedkin, who, by suggesting biochemistry as a career option, set the compass of my life.

kept up with the most recent advances in molecular biology. As an elder statesman he was sought after for advice from foundations, institutes, companies, and other universities in the United States, Europe, and Japan. He thought it ironic that UCSD seemed less interested in his judgment than those beyond the campus.

In an autobiographical account of his career he wrote:

"The biochemist is changing; he is leaning more towards biological problems. This is an admirable turn since the new fields

Fig. 8. Morris Enton Friedkin.

are challenging and exciting. — Many believe that biochemistry of the past decades is ‘old hat’ but it should be remembered that the principles of biochemistry are what have led to the advances in biology. Students should not lose sight of the eloquence of the experiments of Warburg because it is the same eloquence which is inherent in the isolation, characterization, and manipulation of genes” (Kaplan, 1985).

When Nate died in 1986, I lost a dear friend with whom I had shared decades of happy experiences and conversations. Whenever something exciting happens in biochemistry, I imagine what Nate’s enthusiastic reaction would be.

Epilogue

During the Great Depression, David, an actor who was unemployed most of the time in New York City, showered his kid brother with letters of praise, encouragement, advice, and loving care.

In a typewritten letter to me, circa 1936 (I was 18 years old then), David put biochemistry at the top of a list of scientific fields I might enter, including chemistry, physics, and the biological sciences (bacteriology, medicine, etc.), with an interesting handwritten annotation after biochemistry: “This is my pet — for some reason.” Certainly he was not conversant with the discoveries being made in biochemistry. During a period of 9 years I worked my way down my brother’s list, skipping biochemistry. It was not until the war years, when I was a technician on the penicillin project, that my interest in biochemistry caught fire after reading articles published by a group that was working on anaerobic glycolysis in malaria parasites, a war project at the University of Chicago. In 1945, I was accepted for

graduate studies at the University of Chicago and received a doctorate in biochemistry in 1948.

David became a successful writer and director in Hollywood for radio, television, and films. He followed my career closely and with pride. I dedicate this article in memory of my brother, a wonderful human being.

References

- Abraham E. 1985. Penicillin and its successors: A personal view. *Bull Am Acad Sci* 39:8–27.
- Cori CF. 1976. The role of lactic acid in the development of biochemistry. In: Kornberg A, Horecker BL, Oro J, eds. *Reflections on biochemistry in honour of Severo Ochoa*. Oxford, UK: Pergamon Press. pp 17–26.
- Ernster L, Bharaj BS, Nordenbrand K. 1981. Origin of phosphorylation coupled to the oxidation of extramitochondrial NADH. *FEBS Lett* 132:6–9.
- Hach CC, Diehl H. 1948. Studies on oxygen-carrying cobalt compounds. XIV. The design of apparatus for the manufacture of oxygen using Co-Ox and its derivatives. *Iowa State Coll J Sci* 22:165–176.
- Kalckar HM. 1991. 50 Years of biological research — From oxidative phosphorylation to energy requiring transport regulation. *Annu Rev Biochem* 60:1–37.
- Kaplan NO. 1985. Experiences in biochemistry. *Comprehensive Biochem* 36:255–296.
- Kennedy EP. 1992. Sailing to Byzantium. *Annu Rev Biochem* 61:122–127.
- Lambert M. 1980. Northern regional lab honored for work with penicillin. *Journal Star, Peoria, September* 6:B-12.
- Moberg CL. 1991. Penicillin’s forgotten man: Norman Heatley. *Science* 253:734–735.
- Pais A. 1991. *Niels Bohr’s times, in physics, philosophy and polity*. Oxford, UK: Clarendon Press.
- Roeder M. 1980. The penicillin chronology. *Journal Star, Peoria, August* 31:A-10.
- Stodola FH. 1968. Penicillin: Breakthrough to the era of antibiotics. *Yearb Agric* 1968:339–344.
- Taylor JH, Woods PS, Hughes WL. 1957. The organization of chromosomes as revealed by autoradiographic studies using tritium-labeled thymidine. *Proc Natl Acad Sci* 43:122–127.
- Ward GE. 1970. Some contributions of the U.S. Department of Agriculture to the fermentation industry. *Adv Appl Microbiol* 13:363–382.